

Effect of Verapamil on Serum Level of Salinomycin in Diabetic Rats

¹Hossein Najafzadeh, ¹Said Reza Fatemi and ²Amir Ashna

¹Faculty of Veterinary Medicine, Shahid Chamran University, Ahvaz, Iran

²Department of Veterinary Medicine, Faculty of Veterinary Medicine,
Shahid Chamran University, Ahvaz, Iran

Abstract: Problem statement: The aim of present study was evaluation function of P-glycoprotein with or without verapamil in normal and diabetic rats; which the function of P-glycoprotein was indirectly evaluated by detection of serum salinomycin concentration with HPLC method. **Approach:** This study was carried in 4 groups of rats including normal rats which received salinomycin and verapamil together salinomycin; and diabetic rats which received salinomycin and verapamil together salinomycin. Serum concentration of salinomycin was measured by HPLC after 3 h from its administration. **Results:** Results show the serum concentration of salinomycin significantly elevated in diabetic rats which received verapamil together salinomycin; while this concentration did not significantly change in other groups. **Conclusion:** Since the p-glycoprotein activity decreases in diabetic conditions and verapamil inhibits it; probably transport of salinomycin from blood to tissues or its elimination was decreased that caused its elevated serum concentration.

Key words: Salinomycin significantly, verapamil inhibits, diabetes induction, HPLC method, serum concentration, p-glycoprotein, diabetic rats, serum level, tumor cells

INTRODUCTION

P-glycoproteins (P-gp) are transmembrane carrier that prevents the cellular accumulation of some xenobiotics and endogenous compounds and is responsible for certain multidrug resistance mechanism in tumor cells (Aller *et al.*, 2009). P-gp is belonged to ATP-binding cassette transport protein superfamily (ABCB1/MDR1) (De Lange, 2004). This pump is mainly expressed in organs that have an excretion (liver, kidney), absorption (intestine) or barrier function, such as the blood-brain barrier (Mercier *et al.*, 2004; Hsiao *et al.*, 2008). For example, P-gp inhibition at the rat BBB has been determined, when the rat is pretreated with cyclosporine A (as p-gp inhibitor) the brain/plasma ratio of verapamil is increased (Hsiao *et al.*, 2006; Hendrikse and Vaalburg, 2002). Based on these data and others, it has been widely postulated that P-gp plays a vital role in limiting drug distribution at blood-brain barrier and drug interactions will result in an important increase in brain concentrations of the affected drugs and, therefore, their CNS efficacy or toxicity (Neuhaus *et al.*, 2010).

Salinomycin is a polyether antibiotic belonging to the group of ionophores. Salinomycin is extensively used as a coccidiostat in poultry and other livestock

(Rajaian *et al.*, 2009). Severe human poisoning with salinomycin has also been reported (Story and Doube, 2004). P-gp activity may affect the toxic exposure to salinomycin. Individuals with reduced or absent P-gp activity could therefore be more susceptible to salinomycin toxicity (Lagas *et al.*, 2008). Verapamil, a calcium channel blocker, is substrate and inhibitor of p-gp (Sulova *et al.*, 2008).

This study was carried to evaluation function of P-glycoprotein with or without verapamil in normal and diabetic rats; which the function of P-gp was indirectly evaluated by detection of serum salinomycin concentration with HPLC method.

MATERIALS AND METHODS

This study was carried in 4 groups of rats (10 rats in each group). The male Wistar rats were purchased from laboratory animal center Jundishapour of Ahvaz. The age of rats was 12 weeks. The animals access to food and water *ad libitum* under 12h light and 12 dark conditions. Two groups of normal rats received salinomycin (1mg kg⁻¹ orally) and verapamil (25mg kg⁻¹ orally) (Bansal *et al.*, 2009) and 1 h later salinomycin (1 mg kg⁻¹ orally).

Corresponding Author: H. Najafzadeh, Faculty of Veterinary Medicine, Shahid Chamran University, Ahvaz, Iran




Fig. 1: Mean \pm S.E. serum concentration of salinomycin: 1: normal rats received salinomycin, 2: normal rats received salinomycin+ verapamil, 3: diabetic rats received salinomycin, 2: diabetic rats received salinomycin+ verapamil. n=8, * represents significant difference between group4 and other groups ($p<0.05$)

Two groups of rats received streptozocin (45mg kg^{-1} intravenously) for diabetes induction. One group of diabetic rats (those had blood glucose more than 600mg dl^{-1}) received salinomycin (1mg kg^{-1} orally) and another group received verapamil (25mg kg^{-1} orally) and 1 h later salinomycin (1mg kg^{-1} orally). Serum concentration of salinomycin was measured by HPLC after 3 hrs from its administration in all rats.

HPLC system was used from Shimadzu model (Japan) with column C₁₈. Column temperature was set at 40C° . HPLC mobile phase A was water/acetonitrile (95:5, v/v), containing 0.1% formic acid; mobile phase B was acetonitrile containing 0.1% formic acid. The flow rate was 0.6 ml min^{-1} and the injection volume was 20 ml.

RESULTS

Results show minimum level of salinomycin was detected in serum of diabetic rats. Although this level did not significantly differ with normal group. The serum concentration of salinomycin significantly was enhanced in diabetic rats which received verapamil together salinomycin; while this concentration did not significantly change in other groups. The mean of serum concentration of salinomycin (\pm standard error) was illustrated at Fig. 1.

DISCUSSION

Our study shows the interaction salinomycin and verapamil changes in diabetic condition and increases serum level of salinomycin. It seems the some parts of

this interaction are related to p-gp transporting. P-gp function alters in diabetic status (Wu *et al.*, 2009). Nawa *et al.* (2011) evaluated p-gp expression and function in diabetic rats. They demonstrated the expression and function of p-gp significantly decreased after 9 days induction of diabetes (Nawa *et al.*, 2011). Also Liu *et al.* (2006) and Maeng *et al.* (2007) demonstrated the function of p-gp decreased in the blood-brain barrier of streptozotocin-induced diabetic rats (Liu *et al.*, 2006; Maeng *et al.*, 2007). Insulin therapy restores impaired function and expression of P-gp in blood-brain barrier of experimental diabetes (Liu *et al.*, 2008). But we did not demonstrate significant change at serum concentration of salinomycin in diabetic and normal rats. This may be related diuresis in diabetic rats and more excretion of salinomycin.

Salinomycin is a p-gp inhibitor (Riccioni *et al.*, 2010). This drug inhibited p-gp in leukemic cells and increased cell death (Fuchs *et al.*, 2010). In other hands, Verapamil is P-gp inhibitor (Bansal *et al.*, 2009; De Klerk *et al.*, 2010). Administration of verapamil enhanced uptake of dextromethorphan in the CNS of rats (Marier *et al.*, 2005). We demonstrated co-administration of salinomycin and verapamil significantly increased serum level of salinomycin in diabetic rats. We think this enhancing may be related inhibition of salinomycin transport into kidney and CNS by verapamil and diabetic condition. But exact conclusion needs more detailed studies.

CONCLUSION

Since the p-glycoprotein activity decreases in diabetic conditions and verapamil inhibits it; probably transport of salinomycin from blood to tissues or its elimination was decreased that caused its elevated serum concentration.

ACKNOWLEDGMENT

The researchers wish to express their gratitude to the research council of Shahid Chamran University for their financial supports.

REFERENCES

- Aller, S.G., J. Yu, A. Ward, Y. Weng and S. Chittaboina *et al.*, 2009. Structure of P-glycoprotein reveals a molecular basis for poly-specific drug binding. *Science*, 323: 1718-1722. PMID: 19325113
- Bansal, T., G. Mishra, M. Jaggi, R.K. Khar and S. Talegaonkar, 2009. Effect of P-glycoprotein inhibitor, verapamil, on oral bioavailability and pharmacokinetics of irinotecan in rats. *Eur. J. Pharm. Sci.*, 36: 580-590. PMID: 19135530
- De Klerk, O.L., A.T. Willemsen, F.J. Bosker, A.L. Bartels and N.H. Hendrikse *et al.*, 2010. Regional increase in P-glycoprotein function in the blood-brain barrier of patients with chronic schizophrenia: A PET study with [(11)C]verapamil as a probe for P-glycoprotein function. *Psychiatry Res.*, 183: 151-156. PMID: 20620031
- De Lange, E.C., 2004. Potential role of ABC transporters as a detoxification system at the blood-CSF barrier. *Adv. Drug Deliv. Rev.*, 56: 1793-1809. PMID: 15381334
- Fuchs, D., V. Daniel, M. Sadeghi, G. Opelz and C. Naujokat, 2010. Salinomycin overcomes ABC transporter-mediated multidrug and apoptosis resistance in human leukemia stem cell-like KG-1a cells. *Biochem. Biophys. Res. Commun.*, 39: 1098-1104. PMID: 20350531
- Hendrikse, N.H. and W. Vaalburg, 2002. Dynamics of multidrug resistance: P-glycoprotein analyses with positron emission tomography. *Methods*, 27: 228-233. PMID: 12183111
- Hsiao, P., T. Bui, R.J. Ho and J.D. Unadkat, 2008. In vitro-to-in vivo prediction of P-glycoprotein-based drug interactions at the human and rodent blood-brain barrier. *Drug Metab. Dispos.*, 36: 481-484. PMID: 18057117
- Hsiao, P., L. Sasongko, J.M. Link, D.A. Mankoff and M. Muzi *et al.*, 2006. Verapamil P-glycoprotein transport across the rat blood-brain barrier: Cyclosporine, a concentration inhibition analysis and comparison with human data. *J. Pharmacol. Exp. Ther.*, 317: 704-710. PMID: 16415090
- Lagas, J.S., R.W. Sparidans, R.A. van Waterschoot, E. Wagenaar and J.H. Beijnen *et al.*, 2008. P-glycoprotein limits oral availability, brain penetration and toxicity of an anionic drug, the antibiotic salinomycin. *Antimicrob. Agents Chemother.*, 52: 1034-1039. PMID: 18195061
- Liu, H., X. Liu, L. Jia, Y. Liu and H. Yang *et al.*, 2008. Insulin therapy restores impaired function and expression of P-glycoprotein in blood-brain barrier of experimental diabetes. *Biochem. Pharmacol.*, 75: 1649-1658. PMID: 18299117
- Liu, H., X. Xu, Z. Yang, Y. Deng and X. Liu *et al.*, 2006. Impaired function and expression of P-glycoprotein in blood-brain barrier of streptozotocin-induced diabetic rats. *Brain Res.*, 1123: 245-252. PMID: 17074306
- Maeng, H.J., M.H. Kim, H.E. Jin, S.M. Shin and T. Tsuruo *et al.*, 2007. Functional induction of P-glycoprotein in the blood-brain barrier of streptozotocin-induced diabetic rats: Evidence for the involvement of nuclear factor-kappaB, a nitrosative stress-sensitive transcription factor, in the regulation. *Drug Metab. Dispos.*, 35: 1996-2005. DOI: 10.1124/dmd.107.015800 PMID: 17664251
- Marier, J.F., J.L. Deschenes, A. Hage, E. Seliniotakis and A. Gritsas *et al.*, 2005. Enhancing the uptake of dextromethorphan in the CNS of rats by concomitant administration of the P-gp inhibitor verapamil. *Life Sci.*, 77: 2911-2926. PMID: 15964599
- Mercier, C., C. Masseguin, F. Roux, J. Gabrion and J.M. Scherrmann, 2004. Expression of P-glycoprotein (ABCB1) and Mrp1 (ABCC1) in adult rat brain: Focus on astrocytes. *Brain Res.*, 1021: 32-40. PMID: 15328029
- Nawa, A., W. Fujita-Hamabe and S. Tokuyama, 2011. Regulatory action of nitric oxide synthase on ileal p-glycoprotein expression under streptozotocin-induced diabetic condition. *Biol. Pharmacol. Bull.*, 34: 436-438. PMID: 21372399
- Neuhaus, W., M. Stessl, E. Strizsik, B. Bennani-Baiti and M. Wirth *et al.*, 2010. Blood-brain barrier cell line PBMEC/C1-2 possesses functionally active P-glycoprotein. *Neurosci. Lett.*, 469: 224-228. PMID: 19963040

- Rajaian, H., J. Sadjedianfard, S. Nazifi and A.M. Kian, 2009. Drug Interaction between salinomycin and exogenous adrenaline in dogs. *Am. J. Pharmacol. Toxicol.*, 4: 17-21. DOI: 10.3844/ajptsp.2009.17.21
- Riccioni, R., M.L. Dupuis, M. Bernabei, E. Petrucci and L. Pasquini *et al.*, 2010 . The cancer stem cell selective inhibitor salinomycin is a p-glycoprotein inhibitor. *Blood Cells Mol. Dis.*, 45: 86-92. PMID: 20444629
- Story, P. and A. Doube, 2004. A case of human poisoning by salinomycin, an agricultural antibiotic. *N. Z. Med. J.*, 117: 799-799. PMID: 15107902
- Sulova, Z., D. Macejova, M. Seres, J. Sedlak and J. Brtko *et al.*, 2008. Combined treatment of P-gp-positive L1210/VCR cells by verapamil and all-trans retinoic acid induces down-regulation of P-glycoprotein expression and transport activity. *Toxicol. In Vitro*, 22: 96-105. PMID: 17920233
- Wu, K.C., H.J. Pan, H.S. Yin, M.R. Chen and S.C. Lu *et al.*, 2009. Change in P-glycoprotein and caveolin protein expression in brain striatum capillaries in New Zealand Obese mice with type 2 diabetes. *Life Sci.*, 85: 775-781. PMID: 19891976