

## Seroprevalence of *Neospora Caninum* in Stray Dogs

Garedaghi Yagoob

Department of Parasitology, Tabriz Branch,  
Islamic Azad University, Tabriz, Iran

---

**Abstract: Problem statement:** To investigate anti-*Neospora caninum* antibodies in stray dogs living in Tabriz city, 100 blood samples were collected. Serum samples were screened for detection of anti-*N. caninum* IgG antibodies using Indirect Fluorescent Antibody Test (IFAT;  $\geq 50$ ). Antibodies were seen in 31 (31%) of 100 dogs. **Approach:** The IFAT antibody titers were as follows: 1:50 in 11 dogs, 1:100 in ten dogs, 1:200 in six dogs, 1:400 in one dog, 1:800 in two dogs and 1:1,600 in one dog. There were no significant differences in seroprevalence of *Neospora* infection between different genders ( $p > 0.05$ ). The seropositive results were increased with age and differences were statistically significant ( $p < 0.05$ ). **Results:** The results confirm the presence and exposure of stray dogs to *N. caninum* in Tabriz city and importance of this protozoan as a cause of disease in dogs of the region. **Conclusion:** Based on the present study, this is the first report on seroprevalence of *N. caninum* in stray dogs of the Tabriz northwest of Iran. This finding confirms the presence of *Neospora* infection and the important role of dogs in the region.

**Key words:** Higher seroprevalence, *Neospora caninum*, blood samples, serum samples, statistically significant, buffered saline, fluorescent antibody test

---

### INTRODUCTION

*Neospora caninum* is an apicomplexan protozoan which was described by Dubey *et al.* (1988a, 1988b). The agent can infect domestic dogs (*Canis familiaris*) (McAllister *et al.*, 1998) as well as an ample variety of hosts including ruminants (Anderson *et al.*, 1991). It has been regarded as the major cause of abortion with great economical impact in dairy cattle industry and neurologic signs in puppies and older dogs (Ruehlmann *et al.*, 1995; Dubey and Lindsay 1996; Dubey, 2003). Although the parasite can be transmitted transplacentally in several hosts and the vertical route is the major mode of its transmission in cattle, the role of the definitive host in spreading the infection through shedding of oocyst is important. Carnivores can acquire infection by ingestion of infected tissues (McAllister *et al.*, 1998; Dijkstra *et al.*, 2001). As oocysts are rarely found in dog feces (Schaes *et al.*, 2005), serological surveys provide the main clue for estimating the prevalence of canine neosporosis (Bjorkman *et al.*, 1994). The agent has a cosmopolitan distribution, with reports from various countries of Europe, USA, Canada, Australia, South Africa, Japan and Costa Rica (Dubey and Lindsay, 1996). There is also serological evidence of infection in equatorial Africa and South America (Barber *et al.*, 1997a). Seroprevalence (of infection, not disease) varies from 0.5-17% in Europe (Bjorkman *et al.*, 1994; Rasmussen and Jensen, 1996)

and has been reported at 2% in the USA. At present, only limited information is available on the occurrence and prevalence of *Neospora* infections in dogs in Iran. *N. caninum* infection was first reported by Malmasi *et al.* (2007) in household dogs and dogs living in dairy and beef cattle farms. The presence of antibodies against *N. caninum* in aborted and healthy dairy cattle was detected and neosporosis importance in cases of cattle abortion is being increasingly demonstrated in Iran (Razmi *et al.*, 2006; 2007; Sadrebazaz *et al.*, 2004; 2007; Habibi *et al.*, 2005; Bonfitto *et al.*, 2009; Saravanamurthy *et al.*, 2010; Solhjoo *et al.*, 2010). Nevertheless, no information on the prevalence of *N. caninum* infection in stray dogs is available in Iran. Because of this reason, the recent investigation was done to evaluate the seroprevalence of *Neospora* infection in stray dogs of northwest Iran.

### MATERIALS AND METHODS

**Animals:** A total of 100 sera from stray dogs of mixed breeds from districts of Tabriz (63 males, 37 females; <1-6 years of age) were randomly sampled. All of the dogs were examined after the drawing of blood samples. Information regarding access to bovines, clinical signs, gender and age of the animals was also obtained and were registered for retrospective correlation with the anti-*N. caninum* antibody titers in the positive animals.

**Preparation of the serum samples:** Blood samples were drawn from cephalic vein of each dog using venopuncture. The samples were centrifuged at 700×g for 15 min to obtain the serum, which was aliquoted and stored in microtubes and kept frozen at -20°C until analysis.

**Indirect fluorescent antibody test:** The sera were screened for IgG antibodies to *N. caninum* by commercial Indirect Fluorescent Antibody Test (IFAT; Mega Screen Fluoneospora, Horbranz, Austria). All sera were screened at dilutions of 1:50 in phosphate-buffered saline (pH7.2). Serum samples showing fluorescence at the dilution of 1:50 (cutoff value) were further tittered using twofold serial dilutions (Dubey *et al.*, 1988a, 1988b). The entire surface fluorescence of tachyzoite was considered positive (Silva *et al.*, 2007). On each slide, a negative and a positive control were included.

**Statistical analysis:** For the statistical analysis of the possible effects of the attributes of different gender and age groups range on the prevalence of anti-*N. caninum* antibodies, chi-squared test was used. Statistical significance was declared at  $p \leq 0.05$ . The analyses were performed by SPSS 12 statistics program for Windows.

**RESULTS**

Thirty-one out of 100 (31%) stray dogs were found seropositive. The IFAT antibody titers were as follows: 1:50 (n = 11), 1:100 (n = 10), 1:200 (n = 6), 1:400 (n = 1), 1:800 (n = 2) and 1:1,600 (n = 1; Table 1). Twenty-one of 63 (33%) male and 10 of 37 (27%) female stray dogs had Neospora infection (Table 2). The highest infection rate was observed in male dogs (29%). There were no significant differences in seropositivity between different genders ( $p > 0.05$ ). The infection in age group 2-3 years (20%) was higher than in older dogs (1%). There was a significant difference between the infection rate of Neospora among different age

groups ( $p < 0.05$ ). The seropositive results were being increased in the age group of 1-3 years (27%) compared to the dogs of age group 4-10 years (4%).

**DISCUSSION**

Assessing seroprevalence and hence the exposure of dog population of *N. caninum*, is an important part of investigating the possible transmission routes of the parasite as well as identifying populations in which neosporosis may occur. The IFAT is a well-established test for detecting anti-*N. caninum* antibodies in dogs (Dubey *et al.*, 1988a; 1988b). In Iran, the presence of antibodies against *N. caninum* in dairy cattle (Sadrebazzaz *et al.*, 2004; Razmi *et al.*, 2006), camels (Sadrebazzaz *et al.*, 2006) and farm and household dogs (Haddadzadeh *et al.*, 2007; Malmasi *et al.*, 2007) were reported. Based on the present study, this is the first report on the seroprevalence of in stray dogs of the Tabriz northwest of Iran. This finding confirms the presence of Neospora infection and the important role of dogs in the region. The prevalence of 31% was lesser than the prevalence of 33 and 28% reported by Malmasi *et al.* (2007) and Haddadzadeh *et al.* (2007), respectively. This prevalence is similar to the reported prevalence of 22% in urban areas in Argentina (Basso *et al.*, 2001). The higher seroprevalence have been reported in earlier studies on dogs conducted in several countries including USA and Canada (Cheadel *et al.*, 1999), Belgium (Barber *et al.*, 1997b), England (Trees *et al.*, 1993), southern Italy, the Netherlands (Wouda *et al.*, 1999) and Turkey (Coskun *et al.*, 2000). Serologic investigations showed that dogs coming from dairy properties have a greater seroprevalence than those from urban areas, suggesting that farm dogs have a higher risk of exposure to the parasite (Sawada *et al.*, 1998; Wouda *et al.*, 1999; Sanchez *et al.*, 2003).

Table 1: Seroprevalence of *Neospora* antibodies in different age groups of stray dogs

Age (years)	No. of infected dogs	Positives (IFA titers)						Total (%)
		≥50	100	200	400	800	1600	
<1-1	7	2	2	3	0	2	0	7(7)
3-Feb	20	7	7	3	0	1	1	20(20)
5-Apr	3	2	1	0	0	1	0	3(3)
10-Jun	1	0	0	0	1	0	0	1(1)
Total	31	11	10	6	1	2	1	31(31)

Table 2: Prevalence of anti-*N. caninum* based on gender of stray dogs

Antibody titer	1:50	1:100	1:200	1:400	1:800	1:1,600	Total (%)
Infected animal (%)	11(11)	10(10)	6(6)	1(1)	2(2)	1(1)	31(31)
Male	9(29)	7(22)	4(13)	1(3)	0	0	21(68)
Female	2(13)	3(10)	2(6)	0	2(6)	1(3)	10(32)

These higher seroprevalences can be due to the risk factors (consumption of placenta, materials of aborted fetuses, uterine discharge, hunting and close contact with potential intermediate hosts of the parasite (Dijkstra *et al.*, 2002; Fernandes *et al.*, 2004). The presence of anti-*N. caninum* antibodies in city dogs could be associated with subclinical transplacental transmission through several generations as well. To a lesser extent, dogs could become infected by the horizontal route if they consume sporulated oocysts of *N. caninum* (Yakhchali *et al.*, 2010; McAllister *et al.*, 1998; Dubey, 1999). For this reason, in this region, stray dogs should not be accessed to bovine placentas or uterine discharge around the farms or slaughterhouse. There was no significant difference in seropositivity between males (n = 21, 68%) and females (n = 10, 32%; p>0.05), in agreement with the results of other surveys (Trees *et al.*, 1993; Barber *et al.*, 1997a; Sawada *et al.*, 1998; Cheadel *et al.*, 1999; Haddadzadeh *et al.*, 2007; Malmasi *et al.*, 2007). In this study, there is a tendency for elevated risk of pathogen contact with increasing age, suggesting postnatal exposure to *N. caninum*. Similar reports from Brazil (Fernandes *et al.*, 2004) and Iran (Haddadzadeh *et al.*, 2007) confirm this finding. There was an association between seroprevalence and age. The youngest seropositive animal was 2-3 years of age and the oldest was 6-10 years. However, dogs of any age (18 months to 6 years old) may be affected by neosporosis (Knowler and Wheeler, 1995; Patitucci *et al.*, 1997). The role of age in seropositivity suggests that most of the dogs acquire the infection in the postnatal period by means of horizontal transmission. The role of age has been observed in similar studies (Barber and Trees, 1996; Sawada *et al.*, 1998; Wouda *et al.*, 1999; Haddadzadeh *et al.*, 2007; Malmasi *et al.*, 2007). The anti-*N. caninum* titers encountered varied from 50-1600. Thirty (97%) of 31 seropositive dogs had the titers of 1:800 and lower. None of these cases had previous clinical symptoms of the disease, suggesting that a large portion of infection occurs in a subclinical form. In our study, one of the examined dogs showed hind limb paraplegia and had an anti-*N. Caninum* IFAT titer of 1:1, 600. This may indicate an acute infection (Barber and Trees, 1996). Clinical and subclinical infections with *N. caninum* in dogs have been extensively reported worldwide with seroprevalence rates ranging from 0.2-54% (Moore, 2005). The presence of infected stray dog could be a risk factor for the occurrence of *N. caninum*-associated abortions in cattle. Further investigations could be useful to show the correlation between stray dog neosporosis and cattle abortion in this region.

## CONCLUSION

Based on the present study, this is the first report on the seroprevalence of in stray dogs of the Tabriz northwest of Iran. This finding confirms the presence of *Neospora* infection and the important role of dogs in the region.

## ACKNOWLEDGMENT

The researchers wish to thanks the Islamic Azad University, Tabriz Branch, Tabriz, Iran for the financial supports and all laboratory technicians for technical aids in this project.

## REFERENCES

- Anderson, M.L., P.C. Blanchard, B.C. Barr, J.P. Dubey and R.L. Hoffman, 1991. *Neospora*-like protozoan infection as a major cause of abortion in California dairy cattle. *J. Am. Vet. Med. Assoc.*, 198: 241-244. PMID: 2004983
- Barber, J.S. and A.J. Trees, 1996. Clinical aspects of 27 cases of neosporosis in dogs. *Vet. Rec.*, 139: 439-443. DOI: 10.1136/vr.139.18.439 PMID: 8931299
- Barber, J.S., R.B. Gasser, J. Ellis, M.P. Reichel and D. McMillan *et al.*, 1997a. Prevalence of antibodies to *Neospora caninum* in different canid populations. *J. Parasitol.*, 83: 1056-1058. PMID: 9406778
- Barber, J.S., L.V. Ham, I. Polis and A.J. Trees, 1997b. Seroprevalence of antibodies to *neospora caninum* in belgian dogs. *J. Small Anim. Pract.*, 38: 15-16. DOI: 10.1111/j.1748-5827.1997.tb02978.x PMID: 9121127
- Basso, W., L. Venturini, M.C. Venturini, P. Moore and M. Rambeau, 2001. Prevalence of *Neospora caninum* infection in dogs from beef-cattle farms, dairy farms and from urban areas of Argentina. *J. Parasitol.*, 87: 906-907. DOI: 10.1645/0022-3395(2001)087[0906:PONCII]2.0.CO;2 PMID: 11534656
- Bjorkman, C., A. Lunden and A. Uggla, 1994. Prevalence of antibodies to *neospora caninum* and *toxoplasma gondii* in swedish dogs. *Acta Vet. Scand.*, 35: 445-447. PMID: 7676930
- Bonfitto, N.L.B., A.C.F. Motta, R.B. Furini, M.C. Komesu and M.M.P.D. Nascimento *et al.*, 2009. Determination of the salivary anti-phenolic glycolipid-1 antibody in leprosy patients as a tool to monitoring multidrugtherapy. *Am. J. Infect. Dis.*, 5: 314-319. DOI: 10.3844/ajidsp.2009.314.319

- Cheadel, M.A., D.S. Lindsay, S. Rowe, C.C. Dykstra and M.A. Williams, 1999. Prevalence of antibodies to *Neosporacanium* dogs. Int. J. Parasitol., 29: 1537-1543. DOI: 10.1016/S0020-7519(99)00140-X
- Coskun, S.Z., L. Aydyn and C. Bauer, 2000. Seroprevalence of *Neospora caninum* infection in domestic dogs in Turkey. Vet. Rec., 146: 649-649. PMID: 10872789
- Dijkstra, T., M. Eysker, G. Schares, F.J. Conraths and W. Wouda *et al.*, 2001. Dogs shed *Neosporacanium* oocysts after ingestion of naturally infected bovine placenta but not after ingestion of colostrum spiked with *Neosporacanium* tachyzoites. Int. J. Parasitol., 31: 747-752. DOI: 10.1016/S0020-7519(01)00230-2 PMID: 11403764
- Dijkstra, T., H.W. Barkemaa, M. Eysker, J.W. Hesselink and W. Wouda, 2002. Natural transmission routes of *Neospora caninum* between farm dogs and cattle. Vet. Parasitol., 105: 99-104. DOI: 10.1016/S0304-4017(02)00010-9 PMID: 11900923
- Dubey, J.P. and D.S. Lindsay, 1996. A review of *neosporea caninum* and neosporosis. Vet. Parasitol., 67: 1-59. PMID: 9011014
- Dubey, J.P., 1999. Neosporosis in cattle: Biology and economic impact. J. Am. Vet. Med. Assoc., 214: 1160-1163. PMID: 10212674
- Dubey, J.P., 2003. Review of *neosporea caninum* and neosporosis in animals. Korean J. Parasitol., 41: 1-16. PMID: 12666725
- Dubey, J.P., J.L. Carpenter, C.A. Speer, M.J. Topper and A. Uggla, 1988a. Newly recognized fatal protozoan disease of dogs. J. Am. Vet. Med. Assoc., 192: 1269-1285. PMID: 3391851
- Dubey, J.P., A.L. Hattel, D.S. Lindsay and M.J. Topper, 1988b. Neonatal *neosporea caninum* infection in dogs: Isolation of the causative agent and experimental transmission. J. Am. Vet. Med. Assoc., 193: 1259-1263. PMID: 3144521
- Fernandes, B.C.T.M., S.M. Gennari, S.L.P. Souza, J.M. Carvalho and W.G. Oliveira *et al.*, 2004. Prevalence of anti-*neosporea caninum* antibodies in dogs from urban, periurban and rural areas of the city of Uberlândia, Minas Gerais-Brazil. Vet. Parasitol., 123: 33-40. DOI: 10.1016/j.vetpar.2004.05.016
- Habibi, G.R., R. Hashemi-Fesharaki, A. Sadrebazaz, S. Bozorgi and N. Bordbar, 2005. Seminested PCR for diagnosis of *Neospora caninum* infection in cattle. Arch. Razi. Instit., 59: 55-64.
- Haddadzadeh, H.R., A. Sadrebazaz, A. Malmasi, H.T. Ardakani and P.K. Nia *et al.*, 2007. Seroprevalence of *neosporea caninum* infection in dogs from rural and urban environments in Tehran, Iran. Parasitol. Res., 101: 1563-1565. DOI: 10.1007/s00436-007-0678-5 PMID: 17687566
- Knowler, C. and S.J. Wheeler, 1995. *Neospora caninum* infection in three dogs. J. Small Anim. Pract., 36: 172-177. DOI: 10.1111/j.1748-5827.1995.tb02875.x PMID: 7603059
- Malmasi, A., M. Hosseinijad, H. Haddadzadeh, A. Badii and A. Bahonar, 2007. Serologic study of anti-*Neospora caninum* antibodies in household dogs and dogs living in dairy and beef cattle farms in Tehran, Iran. Parasitol. Res., 100: 1143-1145. DOI: 10.1007/s00436-006-0385-7 PMID: 17120042
- McAllister, M.M., J.P. Dubey, D.S. Lindsay, W.R. Jolley and R.A. Wills *et al.*, 1998. Rapid communication: Dogs are definitive hosts of *Neospora caninum*. Int. J. Parasitol., 28: 1473-1479. DOI: 10.1016/S0020-7519(98)00138-6
- Moore, D.P., 2005. Neosporosis in South America. Vet. Parasitol., 127: 87-97. DOI: 10.1016/j.vetpar.2004.10.001 PMID: 15631900
- Patitucci, A.N., M.R. Alley, B.R. Jones and W.A. Charleston, 1997. Protozoal encephalomyelitis of dogs involving *Neospora caninum* and toxoplasma gondii in New Zealand. N.Z. Vet. J., 45: 231-235. PMID: 16031995
- Rasmussen, K. and A.L. Jensen, 1996. Some epidemiologic features of canine neosporosis in Denmark. Vet. Parasitol., 62: 345-349. DOI: 10.1016/0304-4017(95)00867-5 PMID: 8686180
- Razmi, G.R., G.R. Mohammadi, T. Garossi, N. Farzaneh and A.H. Fallah *et al.*, 2006. Seroepidemiology of *Neospora caninum* infection in dairy cattle herds in mashhad area, Iran. Vet. Parasitol., 135: 187-189. DOI: 10.1016/j.vetpar.2005.09.004
- Razmi, G.R., M. Maleki, N. Farzaneh, M.T. Garoussi and A.H. Fallah, 2007. First report of *Neospora caninum*-associated bovine abortion in Mashhad area, Iran. Parasitol. Res., 100: 755-757. DOI: 10.1007/s00436-006-0325-6
- Ruehlmann, D., M. Podell, M. Oglesbee and J.P. Dubey, 1995. Canine neosporosis: A case report and literature review. J. Am. Anim. Hosp. Assoc., 31: 174-183. PMID: 7773765
- Sadrebazaz, A., H. Haddadzadeh, K. Esmailnia, G. Habibi and M. Vojgani *et al.*, 2004. Serological prevalence of *Neospora caninum* in healthy and aborted dairy cattle in Mashhad, Iran. Vet. Parasitol., 124: 201-204. DOI: 10.1016/j.vetpar.2004.06.027

- Sadrebazzaz, A., H. Haddadzadeh and P. Shayan, 2006. Seroprevalence of *Neospora caninum* and *Toxoplasma gondii* in camels (*Camelus dromedarius*) in Mashhad, Iran. *Parasitol. Res.*, 98: 600-601. DOI: 10.1007/s00436-005-0118-3
- Sadrebazzaz, A., G. Habibi, H. Haddadzadeh and J. Ashrafi, 2007. Evaluation of bovine abortion associated with *neospora caninum* by different diagnostic techniques in Mashhad, Iran. *Parasitol. Res.*, 100: 1257-1260. DOI: 10.1007/s00436-006-0417-3
- Sanchez, G.F., S.E. Morales, M.J. Martinez and J.F. Trigo, 2003. Determination and correlation of anti-*Neospora caninum* antibodies in dogs and cattle from Mexico. *Can. J. Vet. Res.*, 67: 142-145. PMID: 12760481
- Saravanamurthy, P.S., P. Rajendran, P.M. Miranda, G. Ashok and S.S. Raghavan *et al.*, 2010. A cross-sectional study of sexual practices, sexually transmitted infections and human immunodeficiency virus among male-to-female transgender people. *Am. Med. J.*, 1: 113-119. DOI: 10.3844/amjpsp.2010.113.119
- Sawada, M., C.H. Park, H. Kondo, T. Morita and A. Shimada *et al.*, 1998. Serological survey of antibody to *Neospora caninum* in Japanese dogs. *J. Vet. Med. Sci.*, 60: 853-854. DOI: 10.1292/jvms.60.853 PMID: 9713815
- Schares, G., N. Pantchev, D. Barutzki, A.O. Heydorn and C. Bauer *et al.*, 2005. Oocysts of *Neospora caninum*, *Hammondia heydorni*, *Toxoplasma gondii* and *Hammondia hammondi* in faeces collected from dogs in Germany. *Int. J. Parasitol.*, 35: 1525-1537. DOI: 10.1016/j.ijpara.2005.08.008 PMID: 16197949
- Silva, D.A.O., J. Lobato, T.W.P. Mineo and J.R. Mineo, 2007. Evaluation of serological tests for the diagnosis of *Neospora caninum* infection in dogs: Optimization of cut off titers and inhibition studies of cross-reactivity with *Toxoplasma gondii*. *Vet. Parasitol.*, 143: 234-244. DOI: 10.1016/j.vetpar.2006.08.028
- Solhjo, K., A.S. Jahromi and A. Parnian-Rad, 2010. Anti-*Toxoplasma gondii* antibodies in haemodialysis patients. *Am. J. Infect. Dis.*, 6: 13-17. DOI: 10.3844/ajidsp.2010.13.17
- Trees, A.J., F. Guy, B.J. Tennant, A.H. Balfour and J.P. Dubey, 1993. Prevalence of antibodies to *Neospora caninum* in a population of urban dogs in England. *Vet. Rec.*, 132: 125-126. DOI: 10.1136/vr.132.6.125 PMID: 8447050
- Wouda, W., T. Dijkstra, A.M.H. Kramer, C.V. Maanen and J.M.A. Brinkhof, 1999. Seroepidemiological evidence for a relationship between *Neospora caninum* infections in dogs and cattle. *Int. J. Parasitol.*, 29: 1677-1682. DOI: 10.1016/S0020-7519(99)00105-8
- Yakhchali, M., S. Javadi and A. Morshedi, 2010. Prevalence of antibodies to *Neospora caninum* in stray dogs of urmia, Iran. *Parasitol. Res.*, 106: 1455-1458. DOI: 10.1007/s00436-010-1824-z