

Homogenization of Earthquake Catalogue in Terms of Magnitude in Iran and Adjoining Region

¹Noushin Naraghiaraghi, ¹Mohd Nawawi,
¹Syed Mustafizur Rahman, ²Ali Beitollahi, ¹Rosli Saad and ³Samieh Joneidi

¹School of Physics, University Science Malaysia, Penang, Malaysia

²Department of Earthquake, Building and Housing Research Center, Tehran, Iran

³Petroiran Development Company (PEDCO), Tehran, Iran

Article history

Received: 12-10-2014

Revised: 31-03-2016

Accepted: 01-04-2016

Corresponding Author:

Noushin Naraghiaraghi
School of Physics, University
Science Malaysia, Penang,
Malaysia

Email: noushin_na@yahoo.com

Abstract: In any seismic hazard assessment a uniform earthquake catalogue is an essential parameter. In this research, an earthquake catalogue of Iran and adjacent areas was studied, using national and international catalogues. The considered region covers a quadrangle limited by 23 to 42°N and 42 to 65°E including Iran and adjacent areas. Earthquake data from the third millennium BC until 2014 were considered in this study. The standardization of the catalogue in terms of magnitude was achieved and new relations were generated to convert all types of magnitude into moment magnitude by using the orthogonal regression technique. Based on the proposed relations, M_W can be estimated and considered as a unified magnitude scale.

Keywords: Orthogonal Regression, Earthquake Catalogue, Magnitude Conversion, Iran and Adjoining Region

Introduction

The earthquake magnitude scale is an important parameter for quantification of earthquakes. Station distribution, changes in instrumentation, the magnitude formula and the data reduction method cause different magnitude scale reports therefore use of a uniform scale is not always possible (Kanamori, 1983).

As a result, different magnitude scales such as M_L , M_S and m_b have been developed and are currently in use. This research aims to standardize the event catalogue in terms of magnitude and generate a uniform catalogue with moment magnitude because compilation of a homogeneous earthquake catalogue is an essential tool for the seismic hazard evaluation. All available international and national catalogues were used to compile the new catalogue. To achieve this goal orthogonal regression method between different magnitude types is used. The final catalogue includes the events from the third millennium BC to 2014.

Different magnitude scales behave variously for all magnitude ranges and for large earthquakes they show saturation effects at different levels. These limitations cause over-estimation or under-estimation of earthquake magnitudes (Scordilis, 2006). Other magnitude scale based on seismic moment, M_W , is considered as the most reliable magnitude because of having no saturation limits

(Kasahara, 1981). Moment magnitude can be calculated while moment-tensor solution is available and it has several benefits comparing with other magnitude scales.

Moment magnitude is a physical parameter of the earthquake and quantitatively links the earthquake process to tectonic deformation (Kagan 2002a; Bird and Kagan, 2004). It does not saturate for large earthquakes and the accuracy is two to three times higher than other magnitudes (Kagan, 2002b; 2003).

The main objective of this research is to develop valid empirical relations converting all kind of magnitude scales to moment magnitudes in order to make a uniform event catalogue based on moment magnitudes. Such relations are useful for compiling uniform earthquake catalogues.

Materials and Methods

Magnitude of an earthquake is the most commonly used descriptor for earthquake size. Body wave magnitude (m_b), Surface wave magnitude (M_S) and Local magnitude (M_L). Local magnitudes (M_L) are the most commonly reported magnitudes in seismic catalogues defined by Richter (1935). Body wave magnitude and the surface wave magnitude were proposed later.

Historical events as well as Instrumental data were used in this study. A comprehensive historical catalogue

for Iran earthquakes was compiled by (Berberian, 1994 and Ambraseys and Melville, 1982).

International databases and regional sources have been consulted for the catalogue construction as followed:

- International Institute of Earthquake Engineering and Seismology (IIEES)
- Iranian Seismological Center (IRSC)
- Building and Housing Research Center (BHRC)
- ISC, International Seismological Centre UK
- NEIC, National Earthquake Information Center
- HRVD, Harvard CMT Catalogue, Harvard Centroid Moment Tensor Catalogue
- MOS, Institution of the Russian Academy of Sciences

Iran plateau is situated between the interaction of Arabian plate and Eurasian plate. Iran is situated between two old continents Eurasia; in north and Africa-Arabia in south and is known for its tectonics (Jarahi *et al.*, 2016).

Figure 1 shows the epicenter of earthquakes in study area. The quantity and reliability of these contributions ensured that the catalogue is well defined. For most of the Iranian earthquake events after 1963 only m_b was reported. m_b saturates for magnitudes bigger than 6.2

(Singh *et al.*, 1983). M_S saturation level is around magnitude 8, so it would be reliable for earthquakes less than magnitude 8. The most suitable magnitude for earthquake and seismic studies is moment magnitude because of having no saturation limits (Kasahara, 1981) and moment magnitude is an input parameter for most of the predicted ground motion equations (Karimiparidari *et al.*, 2013).

Moment magnitude is not reported for historical events and this work has extended the relationships achieved from the instrumental part. These relations are used to convert the magnitudes of historical events into moment magnitude. All catalogues together have been used to develop the equations.

For studies related to earthquake catalogues, it is important to know how different magnitude scales compared with each other (Kagan, 2003). Many empirical relationships have been developed between various magnitude scales for mapping one magnitude type into the other for different regions in the world.

Use of least square linear regression may lead to incorrect results. In such situation, it is suitable to use orthogonal regression procedure (Stromeyer *et al.*, 2004; Joshi and Sharma, 2008; Thingbaijam *et al.*, 2008; Ristau, 2009; Das *et al.*, 2012).




Fig. 1. Epicenters of the earthquakes in study area

However, this regression procedure requires the knowledge of the error variance ratio for the two magnitudes, which is usually not known. An alternative to this problem is to take the error variance ratio equal to unity, assuming that error variance of different magnitudes are approximately equal (Stromeyer *et al.*, 2004; Gutdeutsch *et al.*, 2002; Kaverina *et al.*, 1996; Cavallini and Rebez, 1996; Panza *et al.*, 1993; Gusev, 1991; Ambraseys, 1990). Orthogonal standard regression is used to generate the relations. The details of the orthogonal regression procedure for estimating regression parameters are explained in the literature (Castellaro and Bormann, 2007; Carroll and Ruppert, 1996; Madansky, 1959; Kendall and Stuart, 1946).

This study reviews Orthogonal Regression (OR) (Castellaro and Bormann, 2007), General Orthogonal Regression (GOR), Inverted Standard Least Squares Regression (ISR) and Standard Least-Squares Regression (SR). The relationships were developed for M_s , m_b and M_L (local magnitude).

Results

Homogeneity of the data is considered as one of prime requirements for the catalogue. This study developed several relationships between moment magnitude and other magnitude types in order to convert them into M_W .

The Conversion of Body-Wave Magnitude Information

The conversion relation between body-wave magnitude and moment magnitude has been developed on the basis of 500 earthquake events which body-wave magnitude (m_b) and moment magnitude (M_W) were independently reported.

The proposed conversion relations are expressed in Fig. 2. The relationships are given as followed:

$$\begin{aligned} \text{SR: } M_W &= 1.038216(\pm 0.013) m_b - 0.100564(\pm 0.068) \\ \text{OR: } M_W &= 1.191018(\pm 0.027) m_b - 0.909681(\pm 0.148) \\ \text{ISR: } M_W &= 1.315352(\pm 0.0001) m_b - 1.568054(\pm 0.0001) \end{aligned}$$

The Conversion of Surface-Wave Magnitude Information

In order to develop a reliable relationship between magnitude M_s and M_W , regression analysis was applied for two magnitude ranges: (a) events with magnitude $M_s < 6.1$ and (b) earthquakes with magnitude $6.1 \leq M_s$. The conversion relationship of information expressed as surface-wave magnitude has been built on the basis of 423 events ($M_s < 6.1$) for which both measures of magnitude (M_s and M_W) were reported independently. The obtained conversion relationship is expressed in Fig. 3.




Fig. 2. The regression between M_W and m_b data




Fig. 3. Regression between M_W and M_s for $M_s < 6.1$




Fig. 4. Regressions between M_W and M_s for $6.1 \leq M_s$

About 42 earthquake events have been considered for conversion of M_s magnitudes in the magnitude range of $6.1 \leq M_s$ and the relationship is given in Fig. 3.

The established relationships between M_W and M_s based on 423 events for $M_s < 6.1$ are:

SR: $M_W = 0.545821(\pm 0.013) M_S + 2.559413(\pm 0.066)$
 OR: $M_W = 0.667339(\pm 0.029) M_S + 1.942052(\pm 0.158)$
 ISR: $M_W = 0.998898(\pm 0.0001) M_S + 0.257606(\pm 0.0001)$

For conversion of higher M_S magnitudes in the magnitude range $6.1 \leq M_S \leq 7.4$, we considered 42 events and the relationships are given as followed Fig. 4:

SR: $M_W = 0.834312(\pm 0.043) M_S + 1.002939(\pm 0.288)$
 OR: $M_W = 0.974457(\pm 0.086) M_S + 0.059964(\pm 0.568)$
 ISR: $M_W = 1.145546(\pm 0.046) M_S - 1.091219(\pm 0.307)$

The Conversion of Local Magnitude Information

About 362 earthquake events of the compiled catalogue with information in local magnitude scale (M_L) and moment magnitude scale (M_W) were available. However, the relationships for the conversion have been built based on them. The proposed conversion relationships between M_L - M_W are expressed in Fig. 5.

SR and ISR relationships are also plotted along with OR to illustrate the differences of using these methods. The established relationships between M_W and M_L based on 362 events are:

SR: $M_W = 0.625359(\pm 0.020) M_L + 2.229484(\pm 0.088)$
 OR: $M_W = 0.913323(\pm 0.054) M_L + 0.949395(\pm 0.271)$
 ISR: $M_W = 1.416868(\pm 0.0001) M_L - 1.289013(\pm 0.0001)$

Castellaro and Bormann (2007) suggested calculating OR, SR and ISR. Their study presented that if slope of OR lies in the angular midst between slope of SR and slope of ISR and slope of OR ≈ 1 , then slope of OR is the best regression because in this case, slope of OR is equal to slope of GOR so in this study for all magnitudes slope of OR were considered as the best regression. In order to create a uniform catalogue in terms of magnitude, if is available must use, otherwise, one of the calculated equations must be used to convert to moment magnitude (Karimiparidari *et al.*, 2013).




Fig. 5. Regression between M_L and M_W data

Discussion and Conclusion

The new seismic event catalogue contains data from the third millennium BC to 2014. Orthogonal Regression (OR) method was applied to develop several relationships between different types of magnitudes in order to standardize catalogue in terms of magnitude.

Scordilis (2006) investigated empirical global relations converting, m_b and M_S to moment magnitude and Karimiparidari *et al.* (2013) developed relations to convert m_b , M_S , M_N and M_L into moment magnitude in Iran.

Comparing the results of the present study with a similar investigation done by (Scordilis, 2006; Karimiparidari *et al.*, 2013) for m_b and M_S to M_W and only with (Karimiparidari *et al.*, 2013) for M_L to M_W have been shown in Fig. 6 to 8.

Generally, result of linear regression models in this study is properly similar with the other studies for M_W via M_S conversion but for M_W via m_b the results are significantly different and it can be clearly seen that new relations are more similar to Karimiparidari *et al.* (2013). The new relationship yields slightly lower M_W . The differences between the results of this study beside other studies might be because of using different methods or different magnitude ranges.

Figure 8 shows that the M_L - M_W curves are significantly different and the new relationship yields slightly higher.

As the procedures and priorities are established, updating this catalog should be a straightforward task. There was a lack of sufficient evidence about some historical events, so they were omitted. A future study on Iranian historical earthquakes to consider probable fake events is also recommended.




Fig. 6. Result comparison of this study with M_W versus m_b obtained by (Scordilis, 2006; Karimiparidari *et al.*, 2013)




Fig. 7. Comparison result of this study with M_W versus M_S obtained by (Scordilis, 2006; Karimipardari *et al.*, 2013) for $3 < M_S < 6.1$




Fig. 8. Comparison result of this study with M_W versus M_L and Karimipardari *et al.* (2013)

Acknowledgement

Researchers are grateful to University Science Malaysia for support this research.

Author's Contributions

Noushin Naraghiaraghi: Designed the study, collected data, made the interpretation and wrote the manuscript.

Mohd Nawawi: Collected data and made the interpretation and edited the manuscript.

Syed Mustafizur Rahman: Wrote the manuscript.

Ali Beitollahi: Collected data.

Rosli Saad: Wrote the manuscript.

Samieh Joneidi: Collected data.

Ethics

This article is original. All authors have read and approved the manuscript and no ethical issues involved.

References

- Ambraseys, N.N. and C.P. Melville, 1982. A History of Persian Earthquakes. 1st Edn., Cambridge University Press, Cambridge, ISBN-10: 52124112X, pp: 236.
- Ambraseys, N.N., 1990. Uniform magnitude re-evaluation of European earthquakes associated with strong-motion records. *Earthquake Eng. Structural Dynamics*, 19: 1-20. DOI: 10.1002/eqe.4290190103
- Berberian, M., 1994. Natural Hazards and the First Earthquake Catalogue of Iran. 1st Edn., International Institute of Earthquake Engineers and Seismology, pp: 603.
- Bird, P. and Y.Y. Kagan, 2004. Plate-tectonic analysis of shallow seismicity: Apparent boundary width, beta, corner magnitude, coupled lithosphere thickness and coupling in seven tectonic settings. *Bull. Seismol. Soc. Am.*, 94: 2380-2399. DOI: 10.1785/0120030107
- Carroll, R. and D. Ruppert, 1996. The use and misuse of orthogonal regression in linear errors-in-variables models. *Am. Statist.*, 50: 1-6. DOI: 10.1080/00031305.1996.10473533
- Castellaro, S. and P. Bormann, 2007. Performance of different regression procedures on the magnitude conversion problem. *Bull. Seismol. Society Am.*, 97: 1167-1175. DOI: 10.1785/0120060102
- Cavallini, F. and A. Rebez, 1996. Representing earthquake intensity-magnitude relationship with a nonlinear function. *Bull. Seismol. Society Am.*, 86: 73-78.
- Das, R., H.R. Wason and M.L. Sharma, 2012. Homogenization of earthquake catalog for northeast India and adjoining region. *Pure Applied Geophys.*, 169: 725-731. DOI: 10.1007/s00024-011-0339-6
- Gusev, A.A., 1991. Intermagnitude relationships and asperity statistics. *Pure Applied Geophys.*, 136: 515-527. DOI: 10.1007/BF00878585
- Gutdeutsch, R., D. Kaiser and G. Jentzsch, 2002. Estimation of earthquake magnitudes from epicentral intensities and other focal parameters in Central and Southern Europe. *Geophys. J. Int.*, 151: 824-834. DOI: 10.1046/j.1365-246X.2002.01804.x
- Jarahi, H., N. Naraghiaraghi and M. Nadalian, 2016. Short period spectral acceleration zonation of tehran a comparison between slip and activity rates data's. *Am. J. Geosci.* DOI: 10.3844/ofsp.10450
- Joshi, G.C. and M.L. Sharma, 2008. Uncertainties in the estimation of M_{max} . *J. Earth Syst. Sci.*, 117: 671-682. DOI: 10.1007/s12040-008-0063-5
- Kagan, Y.Y., 2002a. Seismic moment distribution revisited: II. Moment conservation principle. *Geophys. J. Int.*, 149: 731-754. DOI: 10.1046/j.1365-246X.2002.01671.x

- Kagan, Y.Y., 2002b. Modern California earthquake catalogs and their comparison. *Seism. Res. Lett.*, 73: 921-929. DOI: 10.1785/gssrl.73.6.921
- Kagan, Y.Y., 2003. Accuracy of modern global earthquake catalogs. *Phys. Earth Planetary Interiors*, 135: 173-209. DOI: 10.1016/S0031-9201(02)00214-5
- Kanamori, H., 1983. Magnitude scale and quantification of earthquakes. *Tectonophysics*, 93: 185-199. DOI: 10.1016/0040-1951(83)90273-1
- Karimiparidari, S., M. Zare, H. Memarian and A. Kijko, 2013. Iranian earthquakes, a uniform catalog with moment magnitudes. *J. Seismol.*, 17: 897-911. DOI: 10.1007/s10950-013-9360-9
- Kasahara, K., 1981. *Earthquake mechanics*. Cambridge University Press Cambridge.
- Kaverina, A.N., A.V. Lander and A.G. Prozorov, 1996. Global creepex distribution and its relation to earthquake-source geometry and tectonic origin. *Geophys. J. Int.*, 125: 249-265. DOI: 10.1111/j.1365-246X.1996.tb06549.x
- Kendall, M.G. and A. Stuart, 1946. *The advanced theory of statistics*. Charles Griffin and Co, London.
- Madansky, A., 1959. The fitting of straight lines when both variables are subject to error. *J. Am. Statist. Associat.*, 54: 173-205.
- Panza, G.F., A.G. Prozorov and G. Pazzi, 1993. Extension of global creepex definition (M_s - m_b) to local studies (M_d - M_l): The case of the Italian region. *Terra Nova*, 5: 150-156. DOI: 10.1111/j.1365-3121.1993.tb00240.x
- Richter, C.F., 1935. An instrumental earthquake magnitude scale. *Bull. Seismol. Soc. Am.*, 25: 1-32.
- Ristau, J., 2009. Comparison of magnitude estimates for New Zealand earthquakes: Moment magnitude, local magnitude and teleseismic body-wave magnitude. *Bull. Seismol. Society Am.*, 99: 1841-1852. DOI: 10.1785/0120080237
- Scordilis, E.M., 2006. Empirical global relations converting M_S and m_b to moment magnitude. *J. Seismol.*, 10: 225-236. DOI: 10.1007/s10950-006-9012-4
- Singh, S., M. Rodriguez and L. Esteva, 1983. Statistics of small earthquakes and frequency of occurrence of large earthquakes along the Mexican subduction zone. *Bull. Seismol. Society Am.*, 73: 1779-1796.
- Stromeyer, D., G. Grünthal and R. Wahlström, 2004. Chi-square regression for seismic strength parameter relations and their uncertainties, with applications to an M_w based earthquake catalogue for central, northern and northwestern Europe. *J. Seismol.*, 8: 143-153. DOI: 10.1023/B:JOSE.0000009503.80673.51
- Thingbaijam, K.K.S., S.K. Nath, A. Yadav, A. Raj and M.Y. Walling *et al.*, 2008. Recent seismicity in Northeast India and its adjoining region. *J. Seismol.*, 12: 107-123. DOI: 10.1007/s10950-007-9074-y