

Cost Analysis of a Two Dissimilar-Unit Cold Standby Redundant System Subject to Inspection and Random Change in Units

¹G.S. Mokaddis and ²C.H. Matta

¹Department of Mathematics, Faculty of Science,
Ain Shams University, Cairo, Egypt

²Department of Statistics, Faculty of Economics and Political Sciences,
Cairo University, Cairo, Egypt

Abstract: Problem statement: This study dealt with the cost analysis of a two dissimilar unit cold standby redundant system subject to inspection and random change in units. In this system each unit works in two different modes normal and total failure. Assuming that the failure, repair, post repair, interchange of units and inspection times are stochastically independent random variables each having an arbitrary distribution. **Approach:** The system was analyzed by semi Markov process technique. **Results:** The time-dependent availability, steady-state availability, busy period analysis, expected number of visits by the repairman were obtained numerically and cost analysis was obtained numerically and graphically. **Conclusion:** Expected cost per unit time decreased with respect to the increase of failure rate.

Key words: Availability, busy period, expected number of visits by the repairman, cost analysis

INTRODUCTION

Several authors have analyzed two-unit redundant system with two states of operation-operative and failed. Models have been formulated to treat many situations. The system analyses by the semi-Markov process technique. The transition probabilities, mean sojourn time and the mean time to system failure has been obtained by (Mokaddis *et al.*, 1997; 2010).

The purpose of the present study is to study the pointwise availability, steady state availability, busy period, expected number of visits and cost per unit time of the system. This study deals with a model of a two dissimilar-unit cold standby redundant system with two modes normal and failure, where a single repair facility is available with the system for inspection, repair and post repair of the failed unit. The operative and standby units are interchanged at random times to achieve high reliability of the system. After repair, the unit is sent for inspection to decide whether the repair is satisfactory or not. If the failed unit is found unsatisfactory on inspection then the unit sent to pose repair. After repair or post repair of the failed unit it becomes like a new one. The results obtained by (Goel *et al.*, 1992) are derived from this study as special cases.

The following assumptions and notations are used to analyze the system:

- The system consists of a two-dissimilar units, the first is operative and the second is kept as cold standby, which of course does not fail unless it goes into operation
- A unit has two possible modes-normal (the unit functions with full capacity) and total failure (the unit capacity is reduced below the specified level)
- After a random amount of time, the operative unit becomes standby and the standby unit becomes operative if the standby is available
- After failure of an operative unit, the cold standby unit becomes operative instantaneously
- Failure, repair, post repair, interchange of units and inspection times are stochastically independent random variables each having an arbitrary distribution
- There is only one repair facility available with the system to repair, inspection and post repair of the failed unit
- After repair, a unit goes for inspection to decide whether the repair is satisfactory or not, if the repaired unit is found to be unsatisfactory then it is sent for post repair
- The probability of having satisfactory repair is fixed
- On repair or post repair a unit acts like a new unit
- Service discipline is first come first served
- All random variables are mutually independent

Corresponding Author: G.S. Mokaddis, Department of Mathematics, Faculty of Science, Ain Shams University, Cairo, Egypt

Notations and states of the system:

E_0 State of the system at epoch $t = 0$
 E set of regenerative states; $\{S_0, S_1, S_2, S_3, S_4, S_5, S_6, S_7\}$
 \bar{E} set of non-regenerative states; $\{S_8, S_9, S_{10}, S_{11}, S_{12}, S_{13}\}$
 $f_i(t), F_i(t)$ Pdf and cdf of failure time of the i -th unit; $i = 1, 2$
 $\ell_i(t), L_i(t)$ Pdf and cdf of time after the i -th operative unit changes; $i = 1, 2$
 $g_i(t), G_i(t)$ Pdf and cdf of repair time of the i -th failed unit; $i = 1, 2$
 $h_i(t), H_i(t)$ Pdf and cdf of time to complete inspection of the i -th failed unit; $i = 1, 2$
 $k_i(t), K_i(t)$ Pdf and cdf of time to complete post repair of the i -th unsatisfactory failed unit; $i = 1, 2$
 $p_i = (1 - q_i)$ Probability that the repair of the i -th unit is satisfactory after the inspection; $i = 1, 2$
 $q_{ij}(t), Q_{ij}(t)$ Pdf and cdf of first passage time from regenerative state i to a regenerative state j or to a failed state j without visiting any other regenerative state in $(0, t]$; $i, j \in E$
 $q_{ij}^{(k)}(t), Q_{ij}^{(k)}(t)$ Pdf and cdf of first passage time from regenerative state i to a regenerative state j or to a failed state j visiting state k only once in $(0, t]$; $i, j \in E, k \in \bar{E}$
 p_{ij} One step transition probability from state i to state j ; $i, j \in E$
 $p_{ij}^{(k)}$ Probability that the system in state i goes to state j passing through state k ; $i, j \in E, k \in \bar{E}$
 $\pi_i(t)$ Cdf of first passage time from regenerative state i to a failed state
 $M_i(t)$ Probability that the system having started from state i is up at time t without making any transition into any other regenerative state
 $B_i(t)$ Probability that the server is busy at time t given that the system entered regenerative state i at time $t = 0$
 $V_i(t)$ Expected number of visits by the server given that the system started from regenerative state i at time $t = 0$
 μ_{ij} Contribution mean sojourn time in state i when transition is to state j is $-\tilde{Q}_{ij}(0) = q_{ij}^*(0)$
 μ_i Mean sojourn time in state i , $\mu_i = \sum_j [\mu_{ij} + \sum_k \mu_{ij}^{(k)}]$

\sim Symbol for Laplace-Stieltjes transform, e.g., $\tilde{F}(s) = \int e^{-st} dF(t)$
 $*$ Symbol for Laplace transform, e.g., $f^*(s) = \int e^{-st} f(t) dt$
 s Symbol for Stieltjes convolution, e.g., $A(t) s B(t) = \int_0^t B(t-u) dA(u)$
 \odot Symbol for ordinary convolution, e.g., $a(t) \odot b(t) = \int_0^t a(u) b(t-u) du$

For simplicity, whenever integration limits are $(0, \infty)$, they are not written.

Symbols used for the states:

N_{oi} The i -th unit is operative in normal mode; $i = 1, 2$
 N_{Si} The i -th unit is standby in normal mode; $i = 1, 2$
 F_{wri} The i -th unit is in total failure mode and waiting for repair; $i = 1, 2$
 F_{ri} The i -th unit is in total failure mode and under repair; $i = 1, 2$
 F_{Ri} The i -th unit is in total failure mode with repair continued from earlier state; $i = 1, 2$
 F_{Ii} The i -th unit is in total failure mode and under inspection; $i = 1, 2$
 F_{pri} The i -th unit is in total failure mode and under post repair after inspection; $i = 1, 2$

Considering these symbols, the system may be in one of the following states:

$S_0 \equiv (N_{o1}, N_{S2}), S_1 \equiv (N_{S1}, N_{o2}), S_2 \equiv (F_{r1}, N_{o2}),$
 $S_3 \equiv (N_{o1}, F_{r2}), S_4 \equiv (F_{r1}, N_{o2}), S_5 \equiv (N_{o1}, F_{r2}),$
 $S_6 \equiv (F_{pr1}, N_{o2}), S_7 \equiv (N_{o1}, F_{pr2}), S_8 \equiv (F_{R1}, F_{wr2}),$
 $S_9 \equiv (F_{wr1}, F_{R2}), S_{10} \equiv (F_{I1}, F_{wr2}), S_{11} \equiv (F_{wr1}, F_{I2}),$
 $S_{12} \equiv (F_{pr1}, F_{wr2}), S_{13} \equiv (F_{wr1}, F_{pr2})$

Up states: $S_0, S_1, S_2, S_3, S_4, S_5, S_6, S_7.$

Down states: $S_8, S_9, S_{10}, S_{11}, S_{12}, S_{13}.$

States and possible transitions between them are shown in Fig. 1.

Availability analysis: From arguments used in the theory of regenerative processes, the point wise availabilities $A_i(t)$ are seen to satisfy the following relations:

$$\begin{aligned}
 A_0(t) &= M_0(t) + q_{01}(t) \odot A_1(t) + q_{02}(t) \odot A_2(t), \\
 A_1(t) &= M_1(t) + q_{10}(t) \odot A_0(t) + q_{13}(t) \odot A_3(t), \\
 A_2(t) &= M_2(t) + q_{24}(t) \odot A_4(t) + q_{2,10}^{(8)}(t) \odot A_{10}(t), \\
 A_3(t) &= M_3(t) + q_{35}(t) \odot A_5(t) + q_{3,11}^{(9)}(t) \odot A_{11}(t), \\
 A_4(t) &= M_4(t) + q_{41}(t) \odot A_1(t) + q_{46}(t) \odot A_6(t) + \\
 &\quad q_{43}^{(10)}(t) \odot A_3(t) + q_{4,12}^{(10)}(t) \odot A_{12}(t), \\
 A_5(t) &= M_5(t) + q_{50}(t) \odot A_0(t) + q_{57}(t) \odot A_7(t) + \\
 &\quad q_{52}^{(11)}(t) \odot A_2(t) + q_{5,13}^{(11)}(t) \odot A_{13}(t), \\
 A_6(t) &= M_6(t) + q_{61}(t) \odot A_1(t) + q_{63}^{(12)}(t) \odot A_3(t), \\
 A_7(t) &= M_7(t) + q_{70}(t) \odot A_0(t) + q_{72}^{(13)}(t) \odot A_2(t), \\
 A_{10}(t) &= q_{10,3}(t) \odot A_3(t) + q_{10,12}(t) \odot A_{12}(t), \\
 A_{11}(t) &= q_{11,2}(t) \odot A_2(t) + q_{11,13}(t) \odot A_{13}(t), \\
 A_{12}(t) &= q_{12,3}(t) \odot A_3(t), \\
 A_{13}(t) &= q_{13,2}(t) \odot A_2(t)
 \end{aligned}
 \tag{1}$$

Where:

$$\begin{aligned}
 M_0(t) &= \bar{F}_1(t)\bar{L}_1(t), M_1(t) = \bar{F}_2(t)\bar{L}_2(t), M_2(t) = \\
 &\quad \bar{F}_2(t)\bar{G}_1(t)M_3(t) = \bar{F}_1(t)\bar{G}_2(t) \\
 M_4(t) &= \bar{F}_2(t)\bar{H}_1(t), M_5(t) = \bar{F}_1(t)\bar{H}_2(t), M_6 \\
 &\quad (t) = \bar{F}_2(t)\bar{K}_1(t) = \bar{F}_1(t)\bar{K}_2(t)
 \end{aligned}
 \tag{2}$$

Taking Laplace transforms of Eq. 1 and solving for $A_0^*(s)$, it gives:

$$A_0^*(s) = N_1(s) / D_1(s) \tag{3}$$

Where:

$$\begin{aligned}
 N_1(s) &= M_0^*(1 - b_{23}b_{32} - q_{23}^*b_{32}b_{21}) + M_1^*[q_{01}^*(1 - b_{23}b_{32}) + [q_{02}^*b_{21} \\
 &\quad + q_{01}^*q_{13}^*(b_{32}b_{26} + b_{37}) + q_{02}^*[b_{26} + b_{37}(b_{23} + b_{21}q_{13}^*)]]
 \end{aligned}$$




Fig. 1: States and possible transitions

and:

$$\begin{aligned}
 D_1(s) &= (1 - b_{23}b_{32})(1 - q_{01}^*q_{10}^*q_{13}^*(b_{30}q_{01}^* + b_{32}b_{21}) \\
 &\quad - q_{02}^*[b_{21}(q_{10}^* + q_{13}^*b_{30}) + b_{23}b_{30}]), \\
 b_{21} &= q_{24}^*(q_{41}^* + q_{46}^*q_{61}^*), \\
 b_{23} &= q_{24}^*(q_{46}^*q_{63}^{(12)*} + q_{43}^{(10)*} + q_{4,12}^{(10)*}q_{12,3}^*) + q_{2,10}^{(8)*}(q_{10,3}^* + q_{10,12}^*q_{12,3}^*), \\
 b_{26} &= M_2^* + q_{24}^*(M_4^* + q_{46}^*M_6^*), b_{30} = q_{35}^*(q_{50}^* + q_{57}^*q_{70}^*), \\
 b_{32} &= q_{35}^*(q_{57}^*q_{72}^{(13)*} + q_{52}^{(11)*} + q_{5,13}^{(11)*}q_{13,2}^*) + q_{3,11}^{(9)*}(q_{11,2}^* + q_{11,13}^*q_{13,2}^*), \\
 b_{37} &= M_3^* + q_{35}^*(M_5^* + q_{57}^*M_7^*)
 \end{aligned}$$

The steady state availability of the system is:

$$A_0(\infty) = N_1/D_1 \tag{4}$$

Where:

$$\begin{aligned}
 N_1 &= \mu_0(1 - \bar{b}_{23}\bar{b}_{32} - p_{13}\bar{b}_{32}\bar{b}_{21}) + \mu_1[p_{01}(1 - \bar{b}_{23}\bar{b}_{32}) + p_{02}\bar{b}_{21}] \\
 &\quad + p_{01}p_{13}(\bar{b}_{23}\bar{b}_{26} + \bar{b}_{37}) + p_{02}[\bar{b}_{26} + \bar{b}_{37}(\bar{b}_{23} + \bar{b}_{21}p_{13})]
 \end{aligned}$$

and:

$$\begin{aligned}
 D_1 &= (1 - \bar{b}_{23}\bar{b}_{32})(\mu_0p_{10} + p_{01}\mu_{10}) - (\bar{b}_{23}\bar{b}'_{32} + \bar{b}'_{23}\bar{b}_{32})(1 - p_{02}p_{10}) + \\
 &\quad \mu_{13}(\bar{b}_{30}p_{01} + \bar{b}_{32}\bar{b}_{21}) - p_{13}(\bar{b}'_{30}p_{01} + \bar{b}_{30}\mu_{01} + \bar{b}'_{32} + \bar{b}_{21} + \bar{b}_{32}\bar{b}'_{21}) \\
 &\quad + \mu_{02}[\bar{b}_{21}(p_{10} + p_{13}\bar{b}_{30}) + \bar{b}_{23}\bar{b}_{30}] - p_{02}[\bar{b}'_{21}(p_{10} + p_{13}\bar{b}_{30}) \\
 &\quad - \bar{b}_{21}(\mu_0 + \mu_{13}\bar{b}_{30} - p_{13}\bar{b}'_{30}) + \bar{b}'_{23}\bar{b}_{30} + \bar{b}_{23}\bar{b}'_{30}], \\
 \bar{b}_{21} &= p_{24}(p_{41} + p_{46}p_{61}), \\
 \bar{b}_{23} &= p_{24}(p_{46}p_{63}^{(12)} + p_{43}^{(10)} + p_{4,12}^{(10)}p_{12,3}) + p_{2,10}^{(8)}(p_{10,3} + p_{10,12}p_{12,3}), \\
 \bar{b}_{26} &= \mu_2 + p_{24}(\mu_4 + p_{46}\mu_6), \\
 \bar{b}_{30} &= p_{35}(p_{50} + p_{57}p_{70}), \\
 \bar{b}_{32} &= p_{35}(p_{57}p_{72}^{(13)} + p_{52}^{(11)} + p_{5,13}^{(11)}p_{13,2}) + p_{3,11}^{(9)}(p_{11,2} + p_{11,13}p_{13,2}), \\
 \bar{b}_{37} &= \mu_3 + p_{35}(\mu_5 + p_{57}\mu_7), \\
 \bar{b}'_{21} &= -\mu_{24}(p_{41} + p_{46}p_{61}) - p_{24}(\mu_{41} + \mu_{46}p_{61} + p_{46}\mu_{61}), \\
 \bar{b}'_{23} &= -\mu_{24}(p_{46}p_{63}^{(12)} + p_{43}^{(10)} + p_{4,12}^{(10)}p_{12,3}) - p_{24}(\mu_{46}p_{63}^{(12)} \\
 &\quad + p_{46}\mu_{63}^{(12)} + \mu_{43}^{(10)} + \mu_{4,12}^{(10)}p_{12,3} + p_{4,12}^{(10)}\mu_{12,3})\mu_{2,10}^{(8)} \\
 &\quad + (p_{10,3} + p_{10,12}p_{12,3}) - p_{2,10}^{(8)}(\mu_{10,3} + \mu_{10,12}p_{12,3} + p_{10,12}\mu_{12,3}), \\
 \bar{b}'_{30} &= -\mu_{35}(p_{50} + p_{57}p_{70})p_{35}(m_{50} + m_{57}p_{70} + p_{57}m_{70}), \\
 \bar{b}_{23} &= p_{24}(p_{46}p_{63}^{(12)} + p_{43}^{(10)} + p_{4,12}^{(10)}p_{12,3}) + \\
 &\quad p_{2,10}^{(8)}(p_{10,3} + p_{10,12}p_{12,3}), \\
 \bar{b}'_{32} &= -\mu_{35}(p_{57}p_{72}^{(13)} + p_{52}^{(11)} + p_{5,13}^{(11)}p_{13,2}) - p_{35}(\mu_{57}p_{72}^{(13)} \\
 &\quad + p_{57}\mu_{72}^{(13)} + \mu_{52}^{(11)} + \mu_{5,13}^{(11)}p_{13,2} + p_{5,13}^{(11)}\mu_{13,2}) \\
 &\quad - \mu_{3,11}^{(9)}(p_{11,2} + p_{11,13}p_{13,2}) - p_{3,11}^{(9)}(\mu_{11,2} + \mu_{11,13}p_{13,2} \\
 &\quad + p_{11,13}\mu_{13,2})
 \end{aligned}$$

Busy period analysis: We define $B_i(t)$ is the probability that the system is under repair at epoch t given that the system entered a regenerative state S_i at $t = 0$. By probabilistic arguments we have:

$$\begin{aligned}
 B_0(t) &= q_{01}(t) \odot B_1(t) + q_{02}(t) \odot B_2(t), \\
 B_1(t) &= q_{10}(t) \odot B_0(t) + q_{13}(t) \odot B_3(t), \\
 B_2(t) &= V_2(t) + q_{24}(t) \odot B_4(t) + q_{2,10}^{(8)}(t) \odot B_{10}(t), \\
 B_3(t) &= V_3(t) + q_{35}(t) \odot B_5(t) + q_{3,11}^{(9)}(t) \odot B_{11}(t), \\
 B_4(t) &= V_4(t) + q_{11}(t) \odot B_1(t) + q_{46}(t) \odot B_6(t) + \\
 &\quad q_{43}^{(10)}(t) \odot B_3(t) + q_{4,12}^{(10)}(t) \odot B_{12}(t), \\
 B_5(t) &= V_5(t) + q_{50}(t) \odot B_0(t) + q_{57}(t) \odot B_7(t) + \\
 &\quad q_{52}^{(11)}(t) \odot B_2(t) + q_{5,13}^{(11)}(t) \odot B_{13}(t), \\
 B_6(t) &= V_6(t) + q_{61}(t) \odot B_1(t) + q_{63}^{(12)}(t) \odot B_3(t), \\
 B_7(t) &= V_7(t) + q_{70}(t) \odot B_0(t) + q_{72}^{(13)}(t) \odot B_2(t), \\
 B_{10}(t) &= V_{10}(t) + q_{10,3}(t) \odot B_3(t) + q_{10,12}(t) \odot B_{12}(t), \\
 B_{11}(t) &= V_{11}(t) + q_{11,2}(t) \odot B_2(t) + q_{11,13}(t) \odot B_{13}(t), \\
 B_{12}(t) &= V_{12}(t) + q_{12,3}(t) \odot B_3(t), \\
 B_{13}(t) &= V_{13}(t) + q_{13,2}(t) \odot B_2(t)
 \end{aligned}
 \tag{5}$$

Where:

$$\begin{aligned}
 V_2(t) &= \bar{F}_2(t)\bar{G}_1(t), V_3(t) = \bar{F}_1(t)\bar{G}_2(t), V_4(t) \\
 &= \bar{F}_2(t)\bar{H}_1(t), V_5(t) = \bar{F}_1(t)\bar{H}_2(t) \\
 V_6(t) &= \bar{F}_2(t)\bar{K}_1(t), V_7(t) = \bar{F}_1(t)\bar{K}_2(t), \\
 V_{10}(t) &= \bar{H}_1(t), V_{11}(t) = \bar{H}_2(t) \\
 V_{12}(t) &= \bar{K}_1(t), V_{13}(t) = \bar{K}_2(t)
 \end{aligned}
 \tag{6}$$

Taking Laplace transforms of Eq. 5 and solving for $B_0^*(s)$, it follows:

$$B_0^*(s) = N_2(s)/D_1(s) \tag{7}$$

Where:

$$\begin{aligned}
 N_2(s) &= q_{01}^*q_{13}^*(b_{2,12}b_{32} + b_{3,12}) + q_{02}^*[b_{2,12} + b_{23}b_{3,13} \\
 &\quad + b_{21}q_{13}^*b_{3,13}]
 \end{aligned}$$

and:

$D_1(s)$, $b_{21}, b_{23}, b_{25}, b_{30}, b_{32}, b_{37}$ are given in (4):

$$\begin{aligned}
 b_{2,12} &= V_2^* + q_{2,10}^{(8)}(V_{10}^* + q_{10,12}^*V_{12}^*) + q_{24}^*(V_4^* + q_{46}^*V_6^* \\
 &\quad + q_{4,12}^{(10)}V_{12}^*) \\
 b_{3,13} &= V_3^* + q_{3,11}^{(9)}(V_{11}^* + q_{11,13}^*V_{13}^*) + q_{35}^*(V_5^* + q_{57}^*V_7^* \\
 &\quad + q_{5,13}^{(11)}V_{13}^*)
 \end{aligned}$$

In long run the fraction of time for which the server is busy is given by:

$$B_0(\infty) = N_2/D_1 \tag{8}$$

Where:

$$\begin{aligned}
 N_2 &= p_{01}p_{13}(\bar{b}_{2,12}\bar{b}_{32} + \bar{b}_{3,13}) + p_{02}(\bar{b}_{2,12} + \bar{b}_{23}\bar{b}_{3,13} \\
 &\quad + \bar{b}_{21}p_{13}\bar{b}_{3,13})
 \end{aligned}$$

and:

$D_1, \bar{b}_{21}, \bar{b}_{23}, \bar{b}_{26}, \bar{b}_{30}, \bar{b}_{32}, \bar{b}_{37}, \bar{b}'_{21}, \bar{b}'_{23}, \bar{b}'_{30}, \bar{b}'_{32}$ are given in (4):

$$\begin{aligned}
 \bar{b}_{2,12} &= \mu_2 + p_{2,10}^{(8)}(\mu_{10} + p_{10,12}\mu_{12}) + p_{24}(\mu_4 + p_{46}\mu_6 + p_{4,12}^{(10)}\mu_{12}), \\
 \bar{b}_{3,13} &= \mu_3 + p_{3,11}^{(9)}(\mu_{11} + p_{11,13}\mu_{13}) + p_{35}(\mu_5 + p_{57}\mu_7 + p_{5,13}^{(11)}\mu_{13})
 \end{aligned}
 \tag{9}$$

The expected busy period of server in $(0, 1]$ is:

$$\mu_b(t) = \int_0^t B_0(u)du$$

So that:

$$\mu_b^*(s) = B_0^*(s)/s$$

Thus one can evaluate $\mu_b(t)$ by taking inverse Laplace transform of $\mu_b^*(s)$.

Expected idle time of the repairman in $(0, t]$ is:

$$\mu_1(t) = 1 - \mu_b(t)$$

Expected number of visits by the repairman: According to the definition of $V_i(t)$ by elementary probability arguments, the following relations are obtained:

$$\begin{aligned}
 V_0(t) &= Q_{01}(t) s V_1(t) + Q_{02}(t) s [1 + V_2(t)], V_1(t) \\
 &= Q_{10}(t) s V_0(t) + Q_{13}(t) s [1 + V_3(t)], \\
 V_2(t) &= Q_{24}(t) s V_4(t) + Q_{2,10}^{(8)}(t) s V_{10}(t), V_3(t) = \\
 &\quad Q_{35}(t) s V_5(t) + Q_{3,11}^{(9)}(t) s V_{11}(t), \\
 V_4(t) &= Q_{41}(t) s V_1(t) + Q_{46}(t) s V_6(t) + Q_{43}^{(10)}(t) s V_3(t) \\
 &\quad + Q_{4,12}^{(10)}(t) s V_{12}(t), \\
 V_5(t) &= Q_{50}(t) s V_0(t) + Q_{57}(t) s V_7(t) + Q_{52}^{(11)}(t) s V_2 \\
 &\quad (t) + Q_{5,13}^{(11)}(t) s V_{13}(t), \\
 V_6(t) &= Q_{61}(t) s V_1(t) + Q_{63}^{(12)}(t) s V_3(t), V_7(t) \\
 &= Q_{70}(t) s V_0(t) + Q_{72}^{(13)}(t) s V_2(t), \\
 V_{10}(t) &= Q_{10,3}(t) s V_3(t) + Q_{10,12}(t) s V_{12}(t), \\
 V_{11}(t) &= Q_{11,2}(t) s V_2(t) + Q_{11,13}(t) s V_{13}(t), \\
 V_{12}(t) &= Q_{12,3}(t) s V_3(t), V_{13}(t) = Q_{13,2}(t) s V_2(t)
 \end{aligned}
 \tag{10}$$

Taking Laplace-Stieltjes transforms of Eq. 10 and solving for $\tilde{V}_0(s)$, dropping the argument “s” for brevity, it follows:

$$\tilde{V}_0(s) = N_3(s) / D_2(s) \tag{11}$$

Where:

$$N_3(s) = (\tilde{Q}_{01}\tilde{Q}_{13} + \tilde{Q}_{02})(1 - C_{23}C_{32}) + \tilde{Q}_{02}C_{21}\tilde{Q}_{13}(1 - C_{32})$$

and:

$$\begin{aligned} D_2(s) &= (1 - \tilde{Q}_{01}\tilde{Q}_{10})(1 - C_{23}C_{30}) - \tilde{Q}_{13}(C_{30}\tilde{Q}_{01} + C_{32}C_{21}) - \tilde{Q}_{02}[C_{21}(\tilde{Q}_{10} + \tilde{Q}_{13}C_{30}) + C_{23}C_{30}], \\ C_{21} &= \tilde{Q}_{24}(\tilde{Q}_{41} + \tilde{Q}_{46}\tilde{Q}_{61}), \\ C_{23} &= \tilde{Q}_{24}(\tilde{Q}_{46}\tilde{Q}_{63}^{(12)} + \tilde{Q}_{43}^{(10)} + \tilde{Q}_{4,12}^{(10)}\tilde{Q}_{12,3}) + \tilde{Q}_{2,10}^{(8)}(\tilde{Q}_{10,3} + \tilde{Q}_{10,12}\tilde{Q}_{12,3}), \\ C_{30} &= \tilde{Q}_{35}(\tilde{Q}_{50} + \tilde{Q}_{57}\tilde{Q}_{70}), \\ C_{32} &= \tilde{Q}_{35}(\tilde{Q}_{57}\tilde{Q}_{72}^{(13)} + \tilde{Q}_{52}^{(11)} + \tilde{Q}_{5,13}^{(11)}\tilde{Q}_{13,2}) + \tilde{Q}_{3,11}^{(9)}(\tilde{Q}_{11,2} + \tilde{Q}_{11,13}\tilde{Q}_{13,2}) \end{aligned} \tag{12}$$

In steady state, number of visits per unit is given by:

$$V_0(\infty) = N_3/D_2 \tag{13}$$

Where:

$$N_3 = (p_{01}p_{13} + p_{02})(1 - \bar{C}_{23}\bar{C}_{32}) + p_{02}\bar{C}_{21}p_{13}(1 - \bar{C}_{32})$$

and:

$$\begin{aligned} D_2 &= (m_{01}p_{10} + p_{01}m_{10})(1 - \bar{C}_{23}\bar{C}_{30})(1 - p_{01}p_{10}) (\bar{C}'_{23}\bar{C}_{30} + \bar{C}_{23}\bar{C}'_{30}) + \mu_{13}(\bar{C}_{30}p_{01} + \bar{C}_{32}\bar{C}_{21}) + p_{13}(-\bar{C}'_{30}p_{01} + \bar{C}_{30}\mu_{01} \\ &\quad - \bar{C}'_{32}\bar{C}_{21} - \bar{C}_{32}\bar{C}'_{21}) + \mu_{02}[\bar{C}_{21}(p_{10} + p_{13}\bar{C}_{30}) + \bar{C}_{23}\bar{C}_{30}] + p_{02}[-\bar{C}'_{21}(p_{10} + p_{13}\bar{C}_{30}) + \bar{C}_{21}(\mu_{10} + \mu_{13}\bar{C}_{30} - p_{13}\bar{C}'_{30}) \\ &\quad - \bar{C}'_{23}\bar{C}_{30} - \bar{C}_{23}\bar{C}'_{30}], \\ \bar{C}_{21} &= p_{24}(p_{41} + p_{46}p_{61}), \\ \bar{C}_{23} &= p_{24}(p_{46}p_{63}^{(12)} + p_{43}^{(10)} + p_{4,12}^{(10)}p_{12,3}) + p_{2,10}^{(8)}(p_{10,3} + p_{10,12}p_{12,3}), \\ \bar{C}_{30} &= p_{35}(p_{50} + p_{57}p_{70}), \\ \bar{C}_{32} &= p_{35}(p_{57}p_{72}^{(13)} + p_{52}^{(11)} + p_{5,13}^{(11)}p_{13,2}) + p_{3,11}^{(9)}(p_{11,2} + p_{11,13}p_{13,2}), \\ \bar{C}'_{21} &= -m_{02}(p_{41} + p_{46}p_{61}) - p_{24}(m_{41} + m_{46}p_{61} + p_{46}m_{61}), \\ \bar{C}'_{23} &= -\mu_{24}(p_{46}p_{63}^{(12)} + p_{43}^{(10)} + p_{4,12}^{(10)}p_{12,3}) - p_{24}(\mu_{46}p_{63}^{(12)} + p_{46}\mu_{63}^{(12)} + \mu_{43}^{(10)} + \mu_{4,12}^{(10)}p_{12,3} + p_{4,12}^{(10)}\mu_{12,3}) - \mu_{2,10}^{(8)}(p_{10,3} + p_{10,12}p_{12,3}) \\ &\quad - p_{2,10}^{(8)}(\mu_{10,3} + \mu_{10,12}p_{12,3} + p_{10,12}\mu_{12,3}), \\ \bar{C}'_{30} &= -m_{35}(p_{50} + p_{57}p_{70}) - p_{35}(m_{50} + m_{57}p_{70} + p_{57}m_{70}), \\ \bar{C}'_{32} &= -\mu_{35}(p_{57}p_{72}^{(13)} + p_{52}^{(11)} + p_{5,13}^{(11)}p_{13,2}) - p_{35}(\mu_{57}p_{72}^{(13)} + p_{57}\mu_{72}^{(13)} + \mu_{52}^{(11)} + \mu_{5,13}^{(11)}p_{13,2} + p_{5,13}^{(11)}\mu_{13,2}) - \mu_{3,11}^{(9)}(p_{11,2} + p_{11,13}p_{13,2}) \\ &\quad - p_{3,11}^{(9)}(\mu_{11,2} + \mu_{11,13}p_{13,2} + p_{11,13}\mu_{13,2}) \end{aligned} \tag{14}$$

Cost analysis: The cost function of the system obtained by considering the mean-up time of the system, expected busy period of the server and the expected number of visits by the server, therefore, the expected cost incurred in (0, t] is:

$$C(t) = \text{expected total revenue in } (0, t] - \text{expected total service cost in } (0, t] - \text{expected cost of visits by server in } (0, t] = K_1 m_{up}(t) - K_2 m_b(t) - K_3 V_0(t) \tag{15}$$

The expected profit per unit time in steady-state is:

$$C = K_1 A_0 - K_2 B_0 - K_3 V_0 \tag{16}$$

Where:

- K_1 = The revenue per unit up time
- K_2 = The cost per unit time for which system is under repair
- K_3 = The cost per visit by repair facility

Special cases: The two units are dissimilar with exponential distributions:

Let:

- α_i Failure rate of the i-th operative; $i = 1, 2$
- β_i Rate of changes of the i-th operative unit; $i = 1, 2$
- γ_i Rate of repair of the i-th failed unit; $i = 1, 2$
- θ_i Rate of complete inspection of the i-th failed unit; $i = 1, 2$
- λ_i Rate of complete post repair of the i-th unsatisfactory failed unit; $i = 1, 2$
- $p_i = -\langle q_i \rangle$ Probability that the repair of the i-th unit is satisfactory after the inspection; $i = 1, 2$

Transition probabilities are:

$$\begin{aligned} p_{01} &= \beta_1 / (\alpha_1 + \beta_1), & p_{02} &= \alpha_1 / (\alpha_1 + \beta_1), & p_{10} &= \beta_2 / (\alpha_2 + \beta_2), \\ p_{13} &= \alpha_2 / (\alpha_2 + \beta_2), & p_{24} &= \gamma_1 / (\alpha_2 + 1), & p_{28} &= p_{2,10}^{(8)} = \beta_2 / (\alpha_2 + \gamma_1), \\ p_{35} &= \gamma_2 / (\alpha_1 + \gamma_2), & p_{39} &= p_{3,11}^{(9)} = \alpha_1 / (\alpha_1 + \gamma_2), & p_{41} &= p_1 \theta_1 / (\alpha_2 + \theta_1), \\ p_{46} &= q_1 \theta_1 / (\alpha_2 + \theta_1), & p_{4,10} &= \alpha_2 / (\lambda_1 + \alpha_2), & p_{50} &= p_2 \theta_2 / (\alpha_1 + \theta_2), \\ p_{57} &= q_2 \theta_2 / (\alpha_1 + \theta_2), & p_{5,11} &= \alpha_1 / (\alpha_1 + \theta_2), & p_{61} &= \lambda_1 / (\alpha_2 + \lambda_1), \\ p_{6,12} &= p_{63}^{(12)} = \alpha_2 / (\alpha_2 + \lambda_1), & p_{70} &= \lambda_2 / (\alpha_1 + \lambda_2), & p_{7,13} &= p_{72}^{(13)} = \alpha_1 / (\alpha_1 + \lambda_2), \\ p_{10,3} &= p_1, & p_{10,12} &= q_1, & p_{11,2} &= p_2, \\ p_{11,13} &= q_2, & p_{43}^{(10)} &= p_1 \alpha_2 / (\alpha_2 + \theta_1), & p_{4,12}^{(10)} &= q_1 \alpha_2 / (\alpha_2 + \theta_1), \\ p_{52}^{(11)} &= p_2 \alpha_1 / (\alpha_1 + \theta_2), & p_{5,13}^{(11)} &= q_2 \alpha_1 / (\alpha_1 + \theta_2), & p_{8,10} &= p_{9,11} = p_{12,3} = p_{13,2} = 1 \end{aligned}$$

The mean sojourn times are:

$$\begin{aligned} \mu_0 &= 1 / (\alpha_1 + \beta_1), & \mu_1 &= 1 / (\alpha_2 + \beta_2), \mu_2 = 1 / (\alpha_2 + \beta_1), & \mu_3 &= 1 / (\alpha_1 + \gamma_2), \\ \mu_4 &= 1 / (\alpha_2 + \theta_1), & \mu_5 &= 1 / (\alpha_1 + \theta_2), \mu_6 = 1 / (\alpha_2 + \lambda_1), & \mu_7 &= 1 / (\alpha_1 + \lambda_2), \\ \mu_{10} &= 1 / \theta_1, & \mu_{11} &= 1 / \theta_2, \mu_{12} = 1 / \lambda_1, & \mu_{13} &= 1 / \lambda_2 \end{aligned}$$

The mean time to system failure with starting state S_0 is:

$$MTSF = \hat{N}_0 / \hat{D}_0$$

Where:

$$\begin{aligned} \hat{N}_0 &= \frac{1}{(\alpha_1 + \beta_1)(\alpha_2 + \beta_2)} \left\{ \left[\frac{(\beta_2 + \alpha_2 \hat{a}_{30})}{(\alpha_2 + \beta_2)} - \alpha_2 \hat{a}'_{30} \right] \left[(\beta_1 + \alpha_1 \hat{a}_{21}) + \left[\frac{(\beta_1 + \alpha_1 \hat{a}_{21})}{(\alpha_1 + \beta_1)} - \alpha_1 \hat{a}'_{21} \right] (\beta_2 + \alpha_2 \hat{a}_{30}) \right] + \frac{\beta_1 \alpha_2}{(\alpha_1 + \beta_1)(\alpha_2 + \beta_2)} \right. \\ &\quad \left\{ \left[\frac{1}{(\alpha_1 + \beta_1)} + \frac{1}{(\alpha_2 + \beta_2)} \right] \hat{a}_{39} - \hat{a}'_{39} \right\} + \frac{\alpha_1}{(\alpha_1 + \beta_1)} \left[\hat{a}_{28} + \frac{\alpha_2}{(\alpha_2 + \beta_2)} \hat{a}_{21} \hat{a}_{39} \right] - \frac{\alpha_1}{(\alpha_1 + \beta_1)} \left[\hat{a}'_{28} - \frac{\alpha_2}{(\alpha_2 + \beta_2)} \hat{a}'_{21} \hat{a}'_{39} \right] \\ &\quad \left. + \frac{\alpha_2}{(\alpha_2 + \beta_2)} \left(\hat{a}'_{28} \hat{a}_{39} + \hat{a}_{28} \hat{a}'_{39} \right) \right\} \end{aligned}$$

and:

$$\begin{aligned} \hat{D}_0 &= \frac{(\beta_1 + \alpha_1 \hat{a}_{21})(\beta_2 \alpha_2 \hat{a}_{30})}{(\alpha_1 + \beta_1)(\alpha_1 + \beta_2)} \hat{a}_{30} = \frac{\gamma_2 \theta_2}{(\alpha_1 + \gamma_2)(\alpha_1 + \theta_2)} \left[p_2 + \frac{q_2 \lambda_2}{(\alpha_1 + \lambda_2)} \right] \\ \hat{a}_{21} &= \frac{\gamma_1 \theta_1}{(\alpha_2 + \gamma_1)(\alpha_2 + \theta_1)} \left[p_1 + \frac{q_1 \lambda_1}{(\alpha_2 + \lambda_1)} \right] \hat{a}_{28} = \frac{\alpha_2}{(\alpha_2 + \gamma_1)} \left\{ 1 + \frac{\lambda_1}{(\alpha_2 + \lambda_1)} \left[1 + \frac{q_1 \theta_1}{(\alpha_2 + \theta_1)} \right] \right\} \\ \hat{a}_{39} &= \frac{\alpha_1}{(\alpha_1 + \gamma_2)} \left\{ 1 + \frac{\lambda_2}{(\alpha_1 + \lambda_2)} \left[1 + \frac{q_2 \theta_2}{(\alpha_1 + \theta_2)} \right] \right\} \\ \hat{a}'_{30} &= \frac{\gamma_2 \theta_2}{(\alpha_1 + \gamma_2)(\alpha_1 + \theta_2)} \left\{ \frac{1}{(\alpha_1 + \gamma_2)} \left[p_2 + \frac{q_2 \lambda_2}{(\alpha_1 + \gamma_2)} \right] + \frac{p_2}{(\alpha_1 + \theta_2)} + \frac{q_2 \lambda_2}{(\alpha_1 + \lambda_2)} \left[\frac{1}{(\alpha_1 + \theta_2)} + \frac{1}{(\alpha_1 + \lambda_2)} \right] \right\} \\ \hat{a}'_{21} &= \frac{\gamma_1 \theta_1}{(\alpha_2 + \gamma_1)(\alpha_2 + \theta_1)} \left\{ \frac{1}{(\alpha_2 + \gamma_1)} \left[p_1 + \frac{q_1 \lambda_1}{(\alpha_2 + \gamma_1)} \right] + \frac{p_1}{(\alpha_2 + \theta_1)} + \frac{q_1 \lambda_1}{(\alpha_2 + \lambda_1)} \left[\frac{1}{(\alpha_2 + \theta_1)} + \frac{1}{(\alpha_2 + \lambda_1)} \right] \right\} \\ \hat{a}'_{28} &= \frac{\alpha_2}{(\alpha_2 + \gamma_1)^2} - \frac{\gamma_1 \alpha_2}{(\alpha_2 + \gamma_1)(\alpha_2 + \lambda_1)} \left\{ \frac{1}{(\alpha_2 + \gamma_1)} \left[1 + \frac{q_1 \theta_1}{(\alpha_2 + \theta_1)} \right] + \frac{q_1 \theta_1}{(\alpha_2 + \theta_1)} \left[\frac{1}{(\alpha_2 + \theta_1)} + \frac{1}{(\alpha_2 + \lambda_1)} \right] - \frac{1}{(\alpha_2 + \lambda_1)} \right\} \\ \hat{a}'_{39} &= \frac{\alpha_1}{(\alpha_1 + \gamma_2)^2} - \frac{\gamma_2 \alpha_1}{(\alpha_1 + \gamma_2)(\alpha_1 + \lambda_2)} \left\{ \frac{1}{(\alpha_1 + \gamma_2)} \left[1 + \frac{q_2 \theta_2}{(\alpha_1 + \theta_2)} \right] + \frac{q_2 \theta_2}{(\alpha_1 + \theta_2)} \left[\frac{1}{(\alpha_1 + \theta_2)} + \frac{1}{(\alpha_1 + \lambda_2)} \right] - \frac{1}{(\alpha_1 + \lambda_2)} \right\} \end{aligned}$$

In this case, $\hat{M}_i(t)$ are:

$$\begin{aligned} \hat{M}_0(t) &= e^{-(\alpha_1 + \beta_1)t}, & \hat{M}_1(t) &= e^{-(\alpha_2 + \beta_2)t}, & \hat{M}_0(t) &= e^{-(\alpha_1 + \beta_1)t}, & \hat{M}_3(t) &= e^{-(\alpha_1 + \gamma_2)t} \\ \hat{M}_4(t) &= e^{-(\alpha_2 + \theta_1)t}, & \hat{M}_5(t) &= e^{-(\alpha_1 + \theta_2)t}, & \hat{M}_6(t) &= e^{-(\alpha_2 + \lambda_1)t}, & \hat{M}_7(t) &= e^{-(\alpha_1 + \lambda_2)t} \end{aligned}$$

The steady state availability of the system is:

$$\hat{A}_0(\infty) = \hat{N}_1 / \hat{D}_1$$

Where:

$$\begin{aligned} \hat{N}_1 &= \frac{1}{(\alpha_1 + \beta_1)} \left\{ \left[1 - \frac{\hat{b}_{23} \hat{b}_{32}}{\alpha_2 + \beta_2} - \frac{\alpha_2}{(\alpha_2 + \beta_2)} \frac{\hat{b}_{32} \hat{b}_{21}}{\alpha_2 + \beta_2} \right] + \frac{1}{(\alpha_2 + \beta_2)} \left[\beta_1 \left(1 - \frac{\hat{b}_{23} \hat{b}_{32}}{\alpha_2 + \beta_2} \right) + \alpha_1 \hat{b}_{21} \right] + \alpha_1 \left[\hat{b}_{26} + \hat{b}_{37} \left(\frac{\hat{b}_{23}}{\alpha_2 + \beta_2} + \frac{\alpha_2}{\alpha_2 + \beta_2} \hat{b}_{21} \right) \right] \right\} \\ \hat{D}_1 &= \frac{\beta_1 \beta_2 \left(1 - \frac{\hat{b}_{23} \hat{b}_{32}}{\alpha_2 + \beta_2} \right)}{(\alpha_1 + \beta_1)(\alpha_2 + \beta_2)} \left[\frac{1}{(\alpha_1 + \beta_1)} + \frac{1}{(\alpha_2 + \beta_2)} \right] - \left(\frac{\hat{b}_{23} \hat{b}'_{32} + \hat{b}'_{23} \hat{b}_{32}}{\alpha_2 + \beta_2} \right) \left[1 - \frac{\beta_1 \beta_2}{(\alpha_1 + \beta_1)(\alpha_2 + \beta_2)} \right] + \frac{\alpha_2}{(\alpha_2 + \beta_2)} \\ &\quad \left\{ \frac{\hat{b}_{30} \beta_1}{(\alpha_2 + \beta_2)(\alpha_1 + \beta_1)} + \hat{b}_{32} \hat{b}_{21} - \left[\frac{\beta_1}{(\alpha_1 + \beta_1)} \left(\hat{b}'_{30} + \frac{\hat{b}_{30}}{\alpha_1 + \beta_1} \right) + \hat{b}'_{32} \hat{b}_{21} + \hat{b}_{32} \hat{b}'_{21} \right] \right\} + \frac{\alpha_1}{(\alpha_1 + \beta_1)} \left\{ \frac{1}{(\alpha_1 + \beta_1)} \left[\frac{\hat{b}_{21}}{(\alpha_2 + \beta_2)} \left(\beta_2 + \alpha_2 \hat{b}_{30} \right) + \hat{b}_{23} \hat{b}_{30} \right] \right. \\ &\quad \left. - \frac{\hat{b}'_{21}}{(\alpha_2 + \beta_2)} \left(\beta_2 + \alpha_2 \hat{b}_{30} \right) + \frac{\hat{b}_{21}}{(\alpha_2 + \beta_2)} \left[\frac{\beta_2}{(\alpha_2 + \beta_2)} + \frac{\alpha_2 \hat{b}_{30}}{(\alpha_2 + \beta_2)} - \alpha_2 \hat{b}'_{30} \right] - \hat{b}'_{23} \hat{b}'_{30} - \hat{b}_{23} \hat{b}'_{30} \right\}, \\ \hat{b}_{21} &= \frac{\gamma_1 \theta_1}{(\alpha_2 + \gamma_1)(\alpha_2 + \theta_1)} \left[p_1 + \frac{q_1 \lambda_1}{(\alpha_2 + \lambda_1)} \right], \hat{b}_{23} = \frac{1}{(\alpha_2 + \gamma_1)} \left\{ \frac{\gamma_1 \alpha_2}{(\alpha_2 + \theta_1)} \left[\frac{q_1 \theta_1}{(\alpha_2 + \lambda_1)} + 1 \right] + \alpha_2 \right\}, \\ \hat{b}_{26} &= \frac{1}{(\alpha_2 + \gamma_1)} \left\{ 1 + \frac{\gamma_1}{(\alpha_2 + \theta_1)} \left[1 + \frac{q_1 \theta_1}{(\alpha_2 + \lambda_1)} \right] \right\}, \hat{b}_{30} = \frac{\gamma_2 \theta_2}{(\alpha_1 + \gamma_2)(\alpha_1 + \theta_2)} \left[p_2 + \frac{q_2 \lambda_2}{(\alpha_1 + \lambda_2)} \right], \\ \hat{b}_{32} &= \frac{1}{(\alpha_1 + \gamma_2)} \left\{ \frac{\gamma_2 \alpha_1}{(\alpha_1 + \theta_2)} \left[\frac{q_2 \theta_2}{(\alpha_1 + \lambda_2)} + 1 \right] + \alpha_1 \right\}, \hat{b}_{37} = \frac{1}{(\alpha_1 + \gamma_2)} \left\{ 1 + \frac{\gamma_2}{(\alpha_1 + \theta_2)} \left[1 + \frac{q_2 \theta_2}{(\alpha_1 + \lambda_2)} \right] \right\}, \end{aligned}$$

$$\begin{aligned} \hat{b}'_{21} &= \frac{-\gamma_1\theta_1}{(\alpha_2 + \gamma_1)(\alpha_2 + \theta_1)} \left\{ p_1 \left[\frac{1}{(\alpha_2 + \gamma_1)} + \frac{1}{(\alpha_2 + \theta_1)} \right] + \frac{q_1\lambda_1}{(\alpha_2 + \lambda_1)} \left[\frac{1}{(\alpha_2 + \gamma_1)} + \frac{1}{(\alpha_2 + \theta_1)} + \frac{1}{(\alpha_2 + \lambda_1)} \right] \right\} \\ \hat{b}'_{30} &= \frac{-\gamma_2\theta_2}{(\alpha_1 + \gamma_2)(\alpha_1 + \theta_2)} \left\{ p_2 \left[\frac{1}{(\alpha_1 + \gamma_2)} + \frac{1}{(\alpha_1 + \theta_2)} \right] + \frac{q_2\lambda_2}{(\alpha_1 + \lambda_2)} \left[\frac{1}{(\alpha_1 + \gamma_2)} + \frac{1}{(\alpha_1 + \theta_2)} + \frac{1}{(\alpha_1 + \lambda_2)} \right] \right\}, \\ \hat{b}'_{32} &= \frac{-\gamma_2\alpha_1}{(\alpha_1 + \gamma_2)(\alpha_1 + \theta_2)} \left\{ 1 + \frac{q_2\theta_2}{(\alpha_1 + \lambda_2)} \left[\frac{1}{(\alpha_1 + \theta_2)} + \frac{1}{(\alpha_1 + \lambda_2)} + 1 \right] + \frac{1}{(\alpha_1 + \theta_2)} + \frac{q_2}{\lambda_2} \right\} - \frac{\alpha_1}{(\alpha_1 + \gamma_2)} \left[\frac{1}{(\alpha_1 + \gamma_2)} + \frac{1}{\theta_2} + \frac{q_2}{\lambda_2} \right] \end{aligned}$$

In this case, $\hat{V}_i(t)$ are:

$$\begin{aligned} \hat{V}_2(t) &= e^{-(\alpha_2 + \gamma_1)t}, & \hat{V}_3(t) &= e^{-(\alpha_1 + \gamma_2)t}, & \hat{V}_4(t) &= e^{-(\alpha_2 + \theta_1)t}, \\ \hat{V}_5(t) &= e^{-(\alpha_1 + \theta_2)t}, & \hat{V}_6(t) &= e^{-(\alpha_2 + \lambda_1)t}, & \hat{V}_7(t) &= e^{-(\alpha_1 + \lambda_2)t} \end{aligned}$$

In long run, the function of time for which the server is busy is given by:

$$\hat{B}_0(\infty) = \hat{N}_2 / \hat{D}_1$$

Where:

$$\hat{N}_2 = \frac{1}{(\alpha + \gamma)} \left\{ \frac{\alpha}{(\theta + \gamma)^2} \left[1 - \frac{\beta\lambda}{(\beta + \delta)(\lambda + \delta)} \right] + \frac{\beta}{(\lambda + \delta)^2(\beta + \delta)} \left[\frac{\alpha\theta(1 - \theta) + \alpha\gamma}{\theta(\theta + \gamma)} \right] \right\}$$

In steady state, number of visits per unit is given by:

$$\hat{V}_0(\infty) = \hat{N}_3 / \hat{D}_2$$

Where:

$$\hat{N}_3 = \frac{1}{(\alpha_1 + \beta_1)} \left\{ \left[\frac{\beta_1\alpha_2}{(\alpha_2 + \beta_2)} + \alpha_1 \right] \left(1 - \hat{C}_{23}\hat{C}_{32} \right) + \frac{\alpha_1\alpha_2\hat{C}_{21}(1 - \hat{C}_{32})}{(\alpha_2 + \beta_2)} \right\}$$

and:

$$\begin{aligned} \hat{D}_2 &= \frac{\beta_1\beta_2(1 - \hat{C}_{23}\hat{C}_{32})}{(\alpha_1 + \beta_1)(\alpha_2 + \beta_2)} \left[\frac{1}{(\alpha_1 + \beta_1)} + \frac{1}{(\alpha_2 + \beta_2)} \right] - \left[1 - \frac{\beta_1\beta_2}{(\alpha_1 + \beta_1)(\alpha_2 + \beta_2)} \right] \left(\hat{C}'_{23}\hat{C}_{30} + \hat{C}_{23}\hat{C}'_{30} \right) + \frac{\alpha_2}{(\alpha_2 + \beta_2)} \\ &\left\{ \frac{1}{(\alpha_2 + \beta_2)} \left[\frac{\beta_1\hat{C}_{30}}{(\alpha_1 + \beta_1)} + \hat{C}_{32}\hat{C}_{21} \right] + \frac{\beta_1}{(\alpha_1 + \beta_1)} \left[\frac{\hat{C}_{30}}{(\alpha_1 + \beta_1)} - \hat{C}_{30} \right] - \hat{C}'_{32}\hat{C}_{21} - \hat{C}_{32}\hat{C}'_{21} \right\} + \frac{\alpha_1}{(\alpha_1 + \beta_1)} \\ &\left\{ \frac{1}{(\alpha_1 + \beta_1)} \left[\frac{\hat{b}_{21}}{(\alpha_2 + \beta_2)} (\beta_2 + \alpha_2\hat{b}_{30}) + \hat{b}_{23}\hat{b}_{30} \right] - \frac{\hat{b}'_{21}}{(\alpha_2 + \beta_2)} (\beta_2 + \alpha_2\hat{b}_{30}) \frac{\hat{C}_{21}}{(\alpha_2 + \beta_2)} \left[\frac{(\beta_2 + \alpha_2\hat{C}_{30})}{(\alpha_2 + \beta_2)} - \alpha_2\hat{C}'_{30} \right] - \hat{C}'_{23}\hat{C}_{30} - \hat{C}_{23}\hat{C}'_{30} \right\}, \end{aligned}$$

$$\hat{C}_{21} = \frac{\gamma_1\theta_1}{(\alpha_2 + \gamma_1)(\alpha_2 + \theta_1)} \left[p_1 + \frac{q_1\lambda_1}{(\alpha_2 + \lambda_1)} \right], \hat{C}_{23} = \frac{1}{(\alpha_2 + \gamma_1)} \left\{ \frac{\gamma_1\alpha_2}{(\alpha_2 + \theta_1)} \left[\frac{q_1\theta_1}{(\alpha_2 + \lambda_1)} + 1 \right] + \alpha_2 \right\},$$

$$\hat{C}'_{21} = \frac{\theta_1}{(\alpha_2 + \theta_1)} \left\{ \frac{\alpha_1}{(\alpha_1 + \beta_1)^2} \left[p_1 + \frac{q_1\lambda_1}{(\alpha_2 + \lambda_1)} \right] + \frac{\gamma_1}{(\alpha_2 + \gamma_1)} \left[\frac{p_1}{(\alpha_2 + \theta_1)} + \frac{q_1\lambda_1}{(\alpha_2 + \lambda_1)} \left(\frac{1}{(\alpha_2 + \theta_1)} + \frac{1}{(\alpha_2 + \lambda_1)} \right) \right] \right\},$$

$$\hat{C}_{30} = \frac{\gamma_2\theta_2}{(\alpha_1 + \gamma_2)(\alpha_1 + \theta_2)} \left[p_2 + \frac{q_2\lambda_2}{(\alpha_1 + \lambda_2)} \right], \hat{C}_{32} = \frac{1}{(\alpha_1 + \gamma_2)} \left\{ \frac{\gamma_2\alpha_1}{(\alpha_1 + \theta_2)} \left[\frac{q_2\theta_2}{(\alpha_1 + \lambda_2)} + 1 \right] + \alpha_1 \right\},$$

$$\hat{C}_{23}^1 = -\frac{\alpha_2}{(\alpha_2 + \gamma_1)} \left\{ \frac{\gamma_1}{(\alpha_2 + \theta_1)} \left[\frac{1}{(\alpha_2 + \gamma_1)} \left(\frac{q_1 \theta_1}{(\alpha_2 + \lambda_1)} + 1 \right) + \frac{q_1 \theta_1}{(\alpha_2 + \lambda_1)} \left(\frac{1}{(\alpha_2 + \theta_1)} + \frac{1}{(\alpha_2 + \lambda_1)} \right) + \frac{1}{(\alpha_2 + \theta_1)} + \frac{q_1}{\lambda_1} \right] + \frac{1}{(\alpha_2 + \gamma_1)} + \frac{1}{\theta_1} + \frac{q_1}{\lambda_1} \right\}$$

$$\hat{C}_{30}^1 = \frac{-\theta_2}{(\alpha_1 + \theta_2)} \left\{ \frac{\alpha_2}{(\alpha_2 + \beta_2)^2} \left[p_2 + \frac{q_2 \lambda_2}{(\alpha_1 + \lambda_2)} \right] + \frac{\gamma_2}{(\alpha_1 + \gamma_2)} \left[\frac{p_2}{(\alpha_1 + \theta_2)} + \frac{q_2 \lambda_2}{(\alpha_1 + \lambda_2)} \left(\frac{1}{(\alpha_1 + \theta_2)} + \frac{1}{(\alpha_1 + \lambda_2)} \right) \right] \right\}$$

$$\hat{C}_{32}^1 = \frac{\alpha_1}{(\alpha_1 + \gamma_2)} \left\{ \frac{\gamma_2}{(\alpha_1 + \theta_2)} \left[\frac{1}{(\alpha_1 + \gamma_2)} \left(\frac{q_2 \theta_2}{(\alpha_1 + \lambda_2)} + 1 \right) + \frac{q_2 \theta_2}{(\alpha_1 + \lambda_2)} \left(\frac{1}{(\alpha_1 + \theta_2)} + \frac{1}{(\alpha_1 + \lambda_2)} \right) + \frac{1}{(\alpha_1 + \theta_2)} + \frac{q_2}{\lambda_2} \right] + \frac{1}{(\alpha_1 + \gamma_2)} + \frac{1}{\theta_2} + \frac{q_2}{\lambda_2} \right\}$$

Table 1: Relation between the failure rate of the operative unit and the expected cost per unit time at different value of the rate of repair of the failed unit

α	c		
	$\gamma = 0.2$	$\gamma = 0.5$	$\gamma = 0.8$
0.05	1524.83	1150.98	944.38
0.10	1495.48	1100.01	920.75
0.15	1328.60	1000.21	874.95
0.20	1174.51	943.26	840.53
0.25	1049.94	916.48	827.76
0.30	949.44	883.24	820.59
0.35	886.69	845.96	807.00
0.40	835.08	806.64	787.90
0.45	780.45	766.81	764.55

Table 2: Relation between the failure rate of the operative unit and the expected cost per unit time at different value of the rate of complete post repair of unsatisfactory failed unit

α	c		
	$\lambda = 0.2$	$\lambda = 0.5$	$\lambda = 0.8$
0.05	833.95	874.47	890.28
0.10	813.45	870.22	887.06
0.15	795.34	865.46	883.91
0.20	765.26	858.92	880.37
0.25	724.85	852.46	875.99
0.30	678.64	839.02	870.84
0.35	630.55	819.29	864.67
0.40	583.28	794.63	861.03
0.45	538.40	766.49	846.88

Numerical example: Let:

$$\beta = 0.7, \quad \theta = 0.7, \quad q = 0.6, \quad p = 0.4,$$

$$\lambda = 0.5, \quad K_1 = 1000, \quad K_2 = 100, \quad K_3 = 50$$

Let:

$$\beta = 0.6, \quad \theta = 0.7, \quad q = 0.6, \quad p = 0.4,$$

$$\gamma = 0.5, \quad K_1 = 1000, \quad K_2 = 100, \quad K_3 = 50$$

The expected cost per unit time in steady state is:

$$\hat{C} = K_1 \hat{A}_0 - K_2 \hat{B}_0 - K_3 \hat{V}_0$$




Fig. 2: Relation between the failure rate of the operative unit and the expected cost per unit time




Fig. 3: Relation between the failure rate of the operative unit and the expected cost per unit time

The system was analyzed by semi Markov process technique.

The time-dependent availability, steady-state availability, busy period analysis, expected number of visits by the repairman were obtained numerically and cost analysis was obtained numerically as shown in Table 1 and 2 and graphically and as shown in Fig. 2 and 3.

CONCLUSION

Table 1 and 2 compute the expected cost per unit time of the system for different values of failure rate α . The system is decrease with increase of failure rate α as shown in Fig. 2 and 3.

REFERENCES

- Goel, L.R., R.K. Agnihotri and R. Gupta, 1992. Profit evaluation of a two unit cold standby system with random change in units. *Int. J. Syst. Sci.*, 24: 367-377. DOI: 10.1080/00207729208949212
- Mokaddis, G.S., M.L. Tawfek and S.A.M. Elhssia, 1997. Analysis of a two-dissimilar unit cold standby redundant system subject to inspection and random change in units. *Microelect. Reliab.* 37: 329-334. DOI: 10.1016/0026-2714(95)00207-3
- Mokaddis, G.S., M.S. El-Sherbiny and E. Al-Sayeh, 2010. Compare between two-unit cold standby and warm standby outdoor electric power systems in charging weather. *J. Math. Stat.*, 6: 17-22. <http://www.scipub.org/fulltext/jms2/jms26117-22.pdf>