

## Bioactive Compounds Screening from Zingiberaceae Family as Influenza A/Swine Flu Virus Neuraminidase Inhibitor through Docking Approach

Usman Sumo Friend Tambunan, Fadilah and Arli Aditya Parikesit  
Department of Chemistry, Faculty of Mathematics and Science,  
University of Indonesia, Depok 16424, Indonesia

---

**Abstract: Problem statement:** Influenza A/H1N1 is a disease caused by infection of influenza a virus subtype H1N1. It is a major health problem in tropical and subtropical countries. This virus constantly mutates and consequently will be developed into new drug-resistant strains. **Approach:** In this research, we have conducted docking study to screen bioactive compounds from Zingiberaceae family, which has a role as neuraminidase inhibitor of influenza a virus. **Results:** The docking result identified that 1, 2-di-O-  $\beta$ -D-glucopyranosyl-4-allylbenzene (BGA) compound has the affinity and ability to inhibit neuraminidase. There are fourteen residues contact of BGA compound to neuraminidase and eight residues contact of enzyme that formed hydrogen bond with catalytic site. **Conclusion/recommendation:** The docking result showed that BGA has better binding energy and affinity compared with other bioactive compounds and the standard compounds.

**Keywords:** Neuraminidase inhibitory, molecular docking, influenza A/H1N1, bioactive compounds, neuraminidase inhibitor, initial screening process, screen bioactive, subtropical countries, catalytic site, hydrogen bond, conducted docking study,  $\beta$ -D-glucopyranosyl-4-allylbenzene (BGA), surface glycoprotein

---

### INTRODUCTION

Influenza A/H1N1 virus is one of subtype influenza a virus of the Orthomyxoviridae family and one of the main causes of influenza in humans (Van Reeth, 2009). Influenza remains a major cause of morbidity and mortality (Morens *et al.*, 2010). The Lastest influenza pandemic infected 25-30% of the world's population and killing at least 20-50 million world-wide, including more than half a million people in the United States (Thompson *et al.*, 2009). The utilization of antiviral drugs has increased dramatically along with the spread of pandemic influenza A/H1N1 which began in April 2009. The virus is always mutated, forming new strain that are resistant to existing antiviral drug, which also has the potential to infect humans by the means of animal to human transmission or human to human transmission (Moss *et al.*, 2010).

Influenza viruses have a single-stranded nucleic acid, which consists of eight gene segments that encode 11 proteins. The envelope of influenza viruses is composed of complex carbohydrates and two glycoproteins, namely hemagglutinin and neuraminidase (Bantia *et al.*, 2011). Neuraminidase is a

surface glycoprotein, which has an important activity for the replication of influenza A and B virus. This enzyme is responsible for the catalytic cleavage of Sia( $\alpha$ 2-6)Gal or Sia( $\alpha$ 2-3)Gal protein ketosidic bond that exists between the terminal sialic acid and amino acid residues. Bond cleavage has some very important effects. First, it allows the release of virus from infected cells. Second, it prevents the formation of viral aggregates after its release from the host cells (Skehel, 2009; Lai *et al.*, 2010). The neuraminidase is important in viral replication, releases and pathogenicity. Those facts made neuraminidase as a potential antiviral target (Durrant and McCammon, 2010).

The extensive studies of herbs from diversity of medicinal plants present a great opportunity as potential source of good-quality compounds to be developed as anti-influenza drugs. Lead compounds from herbs such as those from Zingiberaceae family have the potential to inhibit neuraminidase and are expected to be further investigated as alternative drugs to inhibit the development of influenza virus A (Grienke *et al.*, 2009) They exist in plants that contain glycoside active compounds as well as polyphenols (Ryu *et al.*, 2010), flavonoids (Ryu *et al.*, 2009a; Jeong *et al.*, 2009; Arora *et al.*, 2010) and xanthon

---

**Corresponding Author:** Usman Sumo Friend Tambunan, Department of Chemistry, Faculty of Mathematics and Science, University of Indonesia, Depok 16424, Indonesia Tel: +62-21-7270027 Fax: +62-21-7863432

(Ryu *et al.*, 2009b), which can be developed as inhibitors of hemagglutinin and neuraminidase biological activity (Sung *et al.*, 2010).

In this research, we have screened bioactive compounds based on their interactions with neuraminidase, using computer software applications (*in silico* method) (Vidal *et al.*, 2011) to determine the best drug lead compounds (Wang *et al.*, 2009). The interaction profiles of the bioactive compounds with neuraminidase were determined by docking experiments (Sandhya *et al.*, 2010; Lee *et al.*, 2009) as an initial screening process for selecting lead compounds, which can bind to the enzyme active site. Analysis was based on Gibbs Free energy values, inhibition constant (Elmi *et al.*, 2009), conformation of the structure, affinity and hydrogen bonding of enzyme and ligands (Krüger and Gohlke, 2010).

## MATERIALS AND METHODS

Neuraminidase enzyme is used in this study. Neuraminidase sequences were obtained from complete sequences of influenza A/H1N1 virus and the data time range is selected from January 2009 until September 2009. Complete sequences were downloaded from NCBI (<http://www.ncbi.nlm.nih.gov/genomes/flu/>). Multiple sequence alignment method was using ClustalW-program ([www.ebi.ac.uk/Tools/clustalw2/index.html](http://www.ebi.ac.uk/Tools/clustalw2/index.html) website). Homology modeling was performed using the Swiss Model server, which can be accessed through <http://swissmodel.expasy.org/SWISS-MODEL.html>. The selected three-dimensional structure was the one with highest similarity toward neuraminidase sequences from 2HTY chain F (PDB Code).

Optimization and minimization of three-dimensional structure of the enzyme were employed using the software of MOE 2008.10, with addition of hydrogen atoms. Protonation was employed with protonate 3D programs. Furthermore, partial charges and force field was employed with MMFF94x. Solvation of enzymes was performed in the form of a gas phase with a fixed charge with a RMS gradient of 0.05 kcal / A<sup>0</sup>mol and other parameters using the standard default values in MOE 2008.10 software.

Preparation of bioactive compounds as a ligand. Bioactive compounds of Zingiberaceae (Lakshmipathy and Kannabiran, 2009; Erukainure *et al.*, 2011; Kiranmayee *et al.*, 2010; Duru and Onyedineke, 2010; Jinsong *et al.*, 2010; Nilesh *et al.*, 2011; Saeed *et al.*, 2010; Cheng *et al.*, 2010; Bhuiyan *et al.*, 2009; Aja *et al.*, 2010) were modeled into three-dimensional structure using ACD Labs software. Three-dimensional shape was obtained by saving it in the 3D viewer in ACDLabs. Furthermore, the output format was changed into Molfile MDL Mol format using software Vegazz to

conform for the docking process. Another adjustments were made with the ligand partial charge and partial charge optimization using MMFF94 forcefield. The conformation structure energy of ligands was minimized using the RMS gradient energy with 0.001 Kcal/A mol. Other parameters were in accordance with default values in the software.

**Docking simulation:** The docking process was begun with the docking preparation, that was employed using a docking program from MOE 2008.10 software. Docking simulations were performed by the Compute-Simulation dock program. Placement method was conducted using a triangle matcher with 1,000,000 repetition energy readings each position and other parameters were in accordance with existing default in the MOE software. Furthermore, scoring functions used London DG, refinement of the configuration repetition forcefield with 1,000 populations. The first repetition of 100 times and the second setting was shown only for one of the best results.

## RESULTS

Neuraminidase sequence > gi|284999325 |gb|ADC32390.1|neuraminidase [Influenza A virus (A/Ancona/15/2009 (H1N1))] was used as target sequence which has a 91.429% similarity with template of 2HTY chain F (PDB Code) and value of Superimpose 0.88 Å<sup>0</sup>.

The bioactive ligands were isolated from species *Alpinia galangal*, *Alpinia conchigera*, *Alpinia officinarum* (Lakshmipathy and Kannabiran, 2009; Erukainure *et al.*, 2011; Kiranmayee *et al.*, 2010; Duru and Onyedineke, 2010; Jinsong *et al.*, 2010), *Curcuma longa*, *Curcuma aromatica*, *Curcuma xanthorrhiza* and *Zingiber zerumbet* (Bhuiyan *et al.*, 2009; Aja *et al.*, 2010). There are a total 99 bioactive compounds, including 13 of flavonoids, 47 of the diarylheptanoids, 18 of the glycosides, 10 phenolics and 11 zerumbones (Kiing *et al.*, 2009; Islas-Osuna *et al.*, 2010; Suthindhiran and Kannabiran, 2009; Vani *et al.*, 2009; Gayosso-García *et al.*, 2010).

The docking results showed that there are 10 out from 99 bioactive ligands that have low Gibbs Free energy. Furthermore, the docking studies of the 10 compounds were repeated with ZNM (Zanamivir) ligand as standard 1 and ILG (Isoliquartigenin) ligand as standard 2. From the 10 of best ligands, 4 ligands have a lower Gibbs energy than ZNM ligand standard. The analysis of the docking results is shown in Table 1.

The BGA compound has of the lowest ΔG value with ΔG = -8.1176 kcal/mol and it is lower than the ΔG of ILG ligand (-7.7494 kcal/mol) isolated from *Glucyrrhiza uralensis* (Gong *et al.*, 2009) and the ΔG of Zanamivir (-5.4946 kcal/mol).

Table 1: Docking results and minimized energy of the complexes enzyme ligands

Ligan	Compound 2*	Compound 3*	Compound 9*	Compound 10*	Std 1*	Std 2*
□G (kcal/mol)	-5.9891	-8.1176	-6.2940	-6.5707	-5.4946	-7.7494
MR	442.461	474.4590	353.3500	354.3550	334.3290	550.5130
Log P	-1.554	-3.2280	3.4220	-0.1880	-6.8550	-1.3630
pKi (µM)	11.896	14.2300	11.2090	10.4820	15.7490	13.7390
H Don	5.000	5.0000	1.0000	2.0000	7.0000	5.0000
H Acc	4.000	8.0000	3.0000	4.0000	6.0000	7.0000

\*) Name of compounds/legends; 1.Compound 2: 1-O-(6-O- $\alpha$ -L-rhamnopyranosyl-  $\beta$ -D-glucopyranosyl)-4-allylbenzene (RGA)2. Compound 3: 1,2-di-O-  $\beta$ -D-glucopyranosyl-4-allylbenzene (BGA) 3: Compound 9: 1E,4Z,6E)-1,7-bis (3,4-dimethoxyphenyl)-5-hydroxyhepta-1,4,6-trien-3-one (DPH) 4. Compound 10: 5Z,7 (E)-6-methoxy-8-(4-hydroxy-3-methoxyphenyl)-5-[(2E)- (4-hydroxy-3- methoxyphenyl)octa-5,7-dienoic acid (MPP) 5. Std 1 : Zanamivir (ZNM) 6. Std 2: Isoliqurtingenin (ILG)

Table 2: Hydrogen bonds of neuraminidase residue with ligands

Ligand	Amino acid residue	Total of H bond
Compound 2 (RGA)	Glu119, Trp179, Trp179, Glu28, Glu278, Arg118, Glu119, Arg152, Arg293	9
Compound 3 (BGA)	Asp151, Ser247, Glu277, Glu277, Glu278, Asn344, Arg118, Ser247, Arg293, Arg293, Arg293, Asn295, Arg368, Arg368	14
Compound 9 (DPH)	Asp151, Arg118, Gln136, Arg368, Lys432	5
Compound 10 (MPP)	Glu119, Glu278, Glu119, Arg152, Arg152, Arg156	6
Std 1 (ZNM)	Glu119, Asp151, Asp151, Ser247, Glu277, Glu277, Glu278, Arg152, Ser247, Arg293, Arg293, Arg368, Arg368	13
Std 2 (ILG)	Val116, Gly147, Asp151, Glu277, Glu278, Gln136, Arg156, Arg156, Arg293, Arg368, Arg430, Thr438	12

\*Bold: catalytic site

Table 3: Score residue contact of neuraminidase with ligands

Ligand	Catalytic site (score %)				
	Arg118	Asp151	Glu278	Arg293	Arg368
Compound 2 (RGA)	40,5	-	23,7	33,9	-
Compound 3 (BGA)	64,0	35,4	42,2	45,3; 24,3; 11,4	76,7; 23,2
Compound 9 (DPH)	26,5	12,7	-	-	21,5
Compound 10 (MPP)	-	-	74,0	-	-
Std 1 (ZNM)	-	31,7; 45,9	27,1	39,7; 13,6	81,3; 83,2
Std 2 (ILG)	-	63,7	27,0	75,1	34,3

The docking result indicates that the BGA ligand has pKi value of 14,430 µM, suggesting that BGA ligand has stronger affinity for neuraminidase than standard 2 (pKi = 13,739 µM). However, when compared with the standard ZNM, pKi value of BGA ligand is lower than ZNM ligand. The results suggest that BGA ligand interacts strongly and effectively with neuraminidase (Table 1).

The hydrogen bonding network between neuraminidase residues and BGA or standard ligands is shown in Tab. 2. The BGA, ZNM and ILG ligands form 14, 13 and 12 hydrogen bonds to neuraminidase, respectively. BGA ligand binds to the catalytic site at the Arg118, Asp151, Glu278, Arg293 and Arg 368 residues and forms as many as 8 hydrogen bonds with the active site. BGA has a higher number of hydrogen bonds to the active site than ZNM standard (7 hydrogen bonds) and ILG standard (4 hydrogen bonds) (Table 2).

Thus, based on the hydrogen bonds, it suggests that the BGA ligand has strong interaction with neuraminidase than those of standards. All three ligands interact with Asp151, Glu78, Arg293 and Arg368 of the catalytic site; however, BGA ligand has additional interaction with the Arg118 residue. The active site binding scores of BGA ligand for neuraminidase are 64% for Arg118; 42.2% for

Glu278; 45.3%, 24.3%, 11.4% for Arg 293; and 76.7%, 23.2% for Arg368 (Table 3).

## DISCUSSION

The conformation of BGA ligand bound to the neuraminidase is shown in Fig. 1. Because the BGA ligand has more residue contacts on the active site than ZNM and ILG, it suggests that BGA interacts more tightly to neuraminidase than ZNM or ILG ligand.

The conformation of BGA is similar to that of ZNM when entering and filling the cavity of neuraminidase binding site. The structure of BGA was superimposed the structure of ZNM as shown in Fig. 2A and BGA filled more surface area of the cavity of the ligand binding site more than ILG (Fig. 2B).

BGA ligand has many interactions with residue contacts at the catalytic site that are similar to those of ZNM ligand (Fig. 1A and 2A). In addition, BGA ligand has better interactions to the catalytic site than ILG ligand (Fig. 1A and 2B). The chemical structure of BGA ligand is shown below:

The difference of the charge within the binding site of BGA, which occurs between the NH<sub>3</sub> clusters binding site of arginine 293 and arginine 368 residues, allowed the interaction with functional group of the ligand.




Fig. 1: The neuraminidase residue contact with ligands, (A) shows neuraminidase contact residues with BGA in 3D; (B) shows neuraminidase contact residues with BGA in 2D; (C) shows neuraminidase contact residues with ZNM in 3D; (D) shows neuraminidase contact residues with ZNM in 2D; (E) shows neuraminidase contact residues with ILG in 3D; (F) shows neuraminidase contact residues with ILG in 2D




Fig. 2: A. The conformation of BGA (blue) with ZNM (green), B. Conformation of BGA (blue) with ILG (red)

It will eventually increase the affinity and stability of the enzyme-ligand complex. Stability and affinity of the ligand complex with the active site was influenced by the distance of the hydrogen bond and complex score. The value of complex score is shown in Table 3.




Fig. 3: The chemical structure of 1, 2-di-O-β-D-glucopyranosyl-4-allylbenzene (BGA). Taken from (Duru and Onyedineke, 2010; Jinsong *et al.*, 2010)

It was indicated that the BGA could change the conformation of neuraminidase cavity upon entering the binding site of neuraminidase (Fig. 3).

Our results suggest that O-glycoside group such as BGA can interact with neuraminidase. This further confirms the results of Sung *et al.* (2010) that O-glycoside group is hydrophilic with a polar functional group, which can inhibit neuraminidase. It is in line with the polar hydrophobic characteristics of neuraminidase.

## CONCLUSION

In conclusion, BGA ligand was found to be the most potent inhibitor of neuraminidase out of 99

compounds evaluated from Zingiberaceae family. The complex between BGA ligand and neuraminidase produces 14 hydrogen bonds with 8 hydrogen bonds at the neuraminidase catalytic site. BGA has the lowest binding energy (-8.1176 kcal/mol) and highest pKi (14,430  $\mu$ M) compared to the two standards (ZNM and ILG). Thus, BGA can be considered to be lead compound for inhibition neuraminidase activity. In the future, BGA will be investigated further as a potential antiviral drug for treating influenza caused by A/H1N1 virus.

#### ACKNOWLEDGEMENT

This research is supported by The Ministry of Education (Grant Hibah Pasca). The authors are grateful to Prof. Teruna J. Siahaan, Ph.D., Department of Pharmaceutical Chemistry, The University of Kansas, Lawrence, USA for his critical comments and excellent assistance in proof-reading the manuscript and to Dr. Ridla Bakri, Chairman of Department of Chemistry, Faculty of Science, University of Indonesia for his support of this research.

#### REFERENCES

- Aja, P.M., A.N.C. Okaka, P.N. Onu, U. Ibiham and A.J. Urako, 2010. Phytochemical composition of *Talinum triangulare* (water leaf) leaves. Pak. J. Nutr., 9: 527-530. DOI: 10.3923/pjn.2010.527.530
- Arora, R., R. Chawla and R. Marwah, 2010. Potential of complementary and alternative medicine in preventive management of novel H1N1 Flu (Swine Flu) Pandemic: Thwarting potential disasters in the bud. Evid. Based Complement Alternat. Med., 1-16. DOI: 10.1155/2011/586506
- Bantia, S., D. Kellogg, C. Parker, R. Upshaw and N.A. Ilyushina *et al.*, 2011. A Single Intramuscular Injection of Neuraminidase inhibitor Peramivir Demonstrates Antiviral Activity against Novel Pandemic A/California/04/2009 (H1N1) Influenza Virus Infection in Mice. *Antiviral Res.* ISSN 0166-3542, DOI: 10.1016/j.antiviral.2011.02.001
- Bhuiyan, M.N.I., C. Jasim Uddin and B. Jaripa, 2009. Chemical investigation of the leaf and rhizome essential oils of *Zingiber zerumbet* (L.) Smith from Bangladesh. Bangladesh J. Pharmacol, 4: 9-12. DOI: 10.3329/bjp.v4i1.845
- Cheng, T.C., C.C. Lu, H.H. Chung, C.C. Hsu and N. Kakizawa *et al.*, 2010. Activation of muscarinic M-1 cholinceptors by curcumin to increase contractility in urinary bladder isolated from Wistar rats. *Neurosci. Lett.*, 473: 107-109. ISSN: 0304-3940, DOI: 10.1016/j.neulet.2010.02.029
- Durrant, J.D. and A. McCammon, 2010. Research article: Potential drug-like inhibitors of Group 1 influenza neuraminidase identified through computer-aided drug design. *Comput. Biol. Chem.*, 34: 97-105. DOI: 10.1016/j.compbiolchem.2010.03.005
- Duru, C.M. and N.E. Onyedineke, 2010. *In vitro* study on the antimicrobial activity and phytochemical analysis of ethanolic extracts of the mesocarp of *Voacanga Africana*. *Am. J. Plant Physiol.*, 5: 163-169. DOI: 10.3923/ajpp.2010.163.169
- Elmi, A., B. Mehdi, C. Madramootoo, R. Dam and D. Smith, 2009. Long-term effect of conventional and no-tillage production systems on nitrous oxide fluxes from corn (*Zea mays* L.) field in southwestern Quebec. *Am. J. Environ. Sci.*, 5: 238-246. DOI: 10.3844/ajessp.2009.238.246
- Erukainure, O.L., O.V. Oke, F.O. Owolabi and O.S. Adenekan, 2011. Chemical composition and antioxidant activities of *Aframomum sceptrum*. *Trends Appl Sci. Res.*, 6: 190-197. DOI: 10.3923/tasr.2011.190.197
- Gayosso-García, S., L.E. Yahia, M. Elhadi, M.A. Martínez-Téllez and G.A. González-Aguilar, 2010. Effect of maturity stage of papaya maradol on physiological and biochemical parameters. *Am. J. Agric. Biol. Sci.*, 5: 194-203. DOI: 10.3844/ajabssp.2010.194.203
- Gong, J., H. Fang, M. Li, Y. Liu and K. Yang *et al.*, 2009. Potential targets and their relevant inhibitors in anti-influenza fields. *Curr. Med. Chem.*, 16: 3716-3739. DOI: 10.2174/092986709789104984
- Grienke, U., M. Schmidtke, J. Kirchmair, K. Pfarr and P. Wutzler *et al.*, 2009. Antiviral potential and molecular insight into neuraminidase inhibiting diarylheptanoids from *alpinia katsumadai*. *J. Med. Chem.*, 53: 778-786. DOI: 10.1021/jm901440f
- Islas-Osuna, M.A., N.A. Stephens-Camacho, C.A. Contreras-Vergara, M. Rivera-Dominguez and E. Sanchez-Sanchez *et al.*, 2010. Novel postharvest treatment reduces ascorbic acid losses in mango (*Mangifera indica* L.) Var. Kent. *Am. J. Agric. Biol. Sci.*, 5: 342-349. DOI: 10.3844/ajabssp.2010.342.349
- Jeong, H.J., Y.B. Ryu, S. Park, J.H. Kim and H. Kwon *et al.*, 2009. Neuraminidase inhibitory activities of flavonols isolated from *Rhodiola rosea* roots and there *in vitro* anti-influenza viral activities. *Bioorganic Med. Chem. Lett.*, 17: 6816-6823. DOI: 10.1016/J.BMC.2009.08.036
- Jinsong, Z., J. Dou, S. Zhang, Q. Liang and Q. Meng, 2010. Chemical composition and antioxidant properties of the essential oil and methanol extracts of rhizoma *Alpinia officinarum* from China *in vitro*. *Afr. J. Biotechnol.*, 9: 4264-4271.

- Kiing, I.C., P.H. Yiu, A. Rajan and S.C. Wong, 2009. Effect of germination on  $\gamma$ -oryzanol content of selected sarawak rice cultivars. *Am. J. Applied Sci.*, 6: 1658-1661. DOI: 10.3844/ajassp.2009.1658.1661
- Kiranmayee, R., C. Bhuvanewari, M.N. Lakshmi and G. Archana, 2010. Antibacterial activity of *Alpinia galanga* (L) wild crude extracts. *Applied Biochem. Biotechnol.*, 162: 871-884. DOI: 10.1007/s12010-009-8900-9
- Krüger, D. and H. Gohlke, 2010. DrugScorePPI for scoring protein-protein interactions: improving A knowledge-based scoring function by atomtype-based QSAR. *J. Cheminform.*, 2: 1-20. DOI: 10.1186/1758-2946-2-S1-P20
- Lai, J.C.C., W.W.L. Chan, F. Kien, J.M. Nicholls and J.S.M. Peiris *et al.*, 2010. Michel formation of virus-like particles from human cell lines exclusively expressing influenza neuraminidase. *J. Gen. Virol.*, 91: 2322-2330. DOI: 10.1099/vir.0.019935-0
- Lakshminpathy, D.T. and K. Kannabiran, 2009. A morphological, biochemical and biological studies of *Halophilic Streptomyces* sp. Isolated from saltpan environment. *Am. J. Infect. Dis.*, 5: 200-206. DOI: 10.3844/ajidsp.2009.200.206
- Lee, H.S., C.S. Lee, S.K. Kim, D.H. Jeong and H. Choe, 2009. Improving virtual screening performance against conformational variations of receptors by shape matching with ligand binding pocket. *J. Chem. Inform. Model.*, 49: 2419-2428.
- Morens, D.M. and J.K. Taubenberger, H.A. Harvey and M.J. Memoli, 2010. The 1918 influenza pandemic: Lessons for 2009 and the future. 38: e10-e20. DOI: 10.1097/CCM.0b013e3181ceb25b
- Moss, R.B., R.T. Davey and F. Fang, 2010. Targeting pandemic influenza: A primer on influenza antivirals and drug resistance. *J. Antimicrob. Chemother.*, 65: 1086-1093. DOI: 10.1093/jac/dkq100
- Nilesh, N.K., B.B. Sandip, I.B. Bajaj, M.V. Bule and R.S. Singhal. 2011. Enzyme-assisted three phase partitioning: A novel approach for extraction of turmeric oleoresin. *Process Biochem.*, 46: 423-426. ISSN: 1359-5113, DOI: 10.1016/j.procbio.2010.09.010
- Ryu, Y.B., J.H. Kim, S.J. Park, J.S. Chang and M.C. Rho *et al.*, 2010. Inhibition of neuraminidase activity polyphenol compounds isolated from the roots of *Glycyrrhiza uralensis*. *Bioorg. Med. Chem Lett.*, 20: 971-974. DOI: 10.1016/j.bmcl.2009.12.106
- Ryu, Y.B., M.J. Curtis-Long, J.W. Lee, H.W. Ryu and J.Y. Kim *et al.*, 2009a. Structural characteristics of flavanones and flavones from *Cudrania tricuspidata* for neuraminidase. *Bioorg. Med. Chem Lett.*, 19: 4912- 915. DOI: 10.1016/j.bmcl.2009.07.098
- Ryu, Y.B., M.J. Curtis-Long, J.W. Lee, J.H. Kim and J.Y. Kim. 2009b. Characteristic of neuraminidase inhibitory xanthenes from *Cudrania tricuspidata*. *Bioorg. Med. Chem Lett.*, 17: 2744-2750. DOI: 10.1016/j.bmcl.2009.02.042
- Saeed, B.A., K.Y. Saour, R.S. Elias and N.A. Al-Masoudi, 2010. Antiviral and quantitative structure activity relationship study for dihydropyridones derived from curcumin. *Am J. Immunol.*, 6: 25-28. DOI: 10.3844/ajisp.2010.25.28
- Sandhya, R.N., R. Subhashini and B. Thiagarajan, 2010. Comparative docking analysis on natural compounds versus a synthetic drug as a therapeutic for acquired immuno deficiency syndrome. *Am. Med. J.*, 1: 148-150. DOI: 10.3844/amjisp.2009.148.150
- Skehel, J., 2009. An overview of influenza haemagglutinin and neuraminidase, *Biologicals*, Volume 37, Issue 3, New Cell for New Vaccines III, New Cell for New Vaccines Conference, pp: 177-178. ISSN: 1045-1056, DOI: 10.1016/j.biologicals.2009.02.012
- Sung, J.C. and A.W.V. Wynsberghe, R.E. Amaro, W.W. Li *et al.*, 2010. Role of secondary sialic acid binding sites in influenza N1 neuraminidase. *J. ACS.*, 132: 2883-2885. DOI: 10.1021/ja9073672
- Suthindhiran, K. and K. Kannabiran, 2009. Cytotoxic and antimicrobial potential of actinomycete species *Saccharopolyspora salina* VITSDK4 isolated from the bay of bengal coast of India. *Am. J. Infect. Dis.*, 5: 90-98. DOI: 10.3844/ajidsp.2009.90.98
- Thompson, W.W., M.R. Moore, E.C. Weintraub, C.B. Bridges and J.S. Bresee *et al.*, 2009. Estimating influenza-associated deaths in the united states. *Am. J. Public Health*, 99: S225-230. DOI: 10.2105/AJPH.2008.15194
- Van Reeth, K.N. 2009. A human case of swine influenza virus infection in Europe-implications for human health and research. *Eur. Surveill*, 14: 19124.
- Vani, S.R., S.F. Cheng and C.H. Chuah, 2009. Comparative study of volatile compounds from genus *ocimum*. *Am. J. Applied Sci.*, 6: 523-528. DOI: 10.3844/ajassp.2009.523.528
- Vidal, D., R. Garcia-Serna and J. Mestres, 2011. Chemoinformatics and computational chemical biology. *Meth. Molecular Biol.*, 672: 489-502. DOI: 10.1007/978-1-60761-839-3\_19
- Wang, Y., J. Xiao, T.O. Suzek, J. Zhang and J. Wang *et al.*, 2009. A public information system for analyzing bioactivities of small molecules. *Nucleic Acids Res.*, 37: 1-11. DOI: 10.1093/nar/gkp456