

## Genetic Capability of Young Layers for Survival after *Salmonella enteritidis* Challenge

<sup>1</sup>E.S. Soliman, <sup>2</sup>H. Busby, <sup>2</sup>C. Kilpatrick, <sup>1</sup>Y. Nagamalleswari, <sup>3</sup>P. McDuff and <sup>4</sup>M.A.A. Sobieh

<sup>1</sup>Department of Pathobiology, College of Veterinary Medicine, Nursing and Allied Health,  
Tuskegee University, Tuskegee,

<sup>2</sup>Alabama State Veterinary Diagnostic Laboratory, Auburn,

<sup>3</sup>Department of Biomedical Science, School of Veterinary Medicine, Auburn University, Auburn,  
Alabama, United States

<sup>4</sup>Department of Animal Hygiene, Zoonoses and Animal Behavior, College of Veterinary Medicine,  
Suez Canal University, Egypt

---

**Abstract: Problem statement:** Genetic line differences in resistance of layer hens and young chicks to *Salmonella enteritidis* have been identified through a lot of studies. That is why the agricultural industry was prepared for the potential phasing out of antibiotics for use in controlling bacterial pathogens. Early infection may result in long term colonization of layers with *Salmonella enteritidis*, resulting in shedding into table or hatching eggs. **Approach:** This study was carried to evaluate the genetic factors underlying early response to *Salmonella enteritidis*, genetic line differences in mortality and pathogen load at two sites (cecal lumen and spleen) were investigated. At day of hatch, chicks of four genetic lines were intra-esophageally inoculated with one of three doses of *Salmonella enteritidis* phage type 13 A. **Results:** There was a significant effect ( $p \leq 0.001$ ) of genetic line on chick 6 days survival. The effect of genetic line was significant ( $p \leq 0.05$ ) on survivors' *Salmonella enteritidis* burden in cecal content but not on *Salmonella enteritidis* load per gram of spleen. *Salmonella enteritidis* pathogen load of the spleen and the cecal content were not significantly correlated, indicating that independent host mechanisms are partly responsible for these two traits. **Conclusion:** Future *Salmonella enteritidis* control mechanism in poultry may be the same as it is used these days but it has to be for longer term sustainability, genetic resistance should be pursued. Sufficient genetic line variation should exist to suggest that it is feasible to effectively choose among pure breeder lines for those exhibits reduced *Salmonella enteritidis* induced mortality and cecal content *Salmonella enteritidis* pathogen load in young layer chicks.

**Key words:** Susceptibility, layers, *Salmonella enteritidis*, genetics

---

### INTRODUCTION

In the United States, an estimated 500,000 cases of human illness are annually attributed to *Salmonella enteritidis* contaminated food products<sup>[17]</sup>. The primary route for *Salmonellae* transmission in poultry is vertical transmission via contaminated eggs, but lateral transmission also occurs via contaminated feed, water and facilities or via host reservoirs such as wild birds, rodents, pet and humans<sup>[13]</sup>. The *Salmonella enteritidis* serotypes colonized at an early age are maintained throughout adulthood in layer chickens<sup>[8]</sup>. *Salmonella enteritidis* contamination in layer hens can result in decreased egg production and furthermore, the bacteria can contaminate the laid eggs<sup>[6]</sup>. Gast<sup>[9]</sup> has suggested that reduction of

*Salmonella enteritidis* pathogen load in chicks reduces the *Salmonella enteritidis* contaminated eggs produced by the hens. Management and pharmaceutical approaches such as vaccination, competitive exclusion and antibiotic treatments can help reduce the *Salmonella* burden in poultry<sup>[9]</sup>. Additional reduction in pathogen load in poultry may be obtainable through genetic selection for disease resistance; Lamont<sup>[14]</sup>. Previous studies have demonstrated the polygenic nature of disease resistance to *Salmonella enteritidis* in poultry; Bumstead and Barrow<sup>[3,4,12,15,16]</sup>. Others Gorham *et al.*<sup>[1,11]</sup> have measured mortality and frequency of colonization in young chicks, but little is known regarding the quantifiable pathogen load of *Salmonella enteritidis* in the internal organs. Genetic factors involved in early resistance and immune

---

**Corresponding Author:** E.S. Soliman, Department of Pathobiology, College of Veterinary Medicine,  
Nursing and Allied Health, Tuskegee University, Tuskegee, Alabama, United States

response in layer chicks are not well understood. The objective of this study was to evaluate genetic line effect on survival and pathogen load after *Salmonella enteritidis* exposure in young layer chicks.

## MATERIALS AND METHODS

**Birds:** Two hundred chicks from four layer lines equally represented for sex and genetic line. Three pure lines and one experimental cross were used. The experimental cross was produced from the three pure line strains. To ensure that the maternal immune status of all hens producing the chicks was equivalent and would not interfere with testing their chicks for *salmonella* response, all hens were kept under the same biosecurity measures and managemental conditions and were from breeding flocks that were tested weekly for freedom from *Salmonella* sp. The chicks were equally divided, with regard to genetic line, sex and inoculation dose, into three biosafety level two animal rooms. The chicks were given access to water and food *ad libitum*. The feed contained standard feed additive amounts of Amprolium (0.0125%).

***Salmonella enteritidis* challenge:** The *Salmonella enteritidis* phage type 13A was brought into the exponential growth phase by incubating in Luria-Bertani (LB) broth (Fisher Sci) for approximately 4 h at 37°C. Inoculation dose dilutions of *Salmonella enteritidis* were made based on a concentration estimated from the optical density at 600 nm. Actual *Salmonella enteritidis* inoculation doses were then confirmed by serial plate dilutions of the inoculum. Chicks were intraesophageally inoculated with *Salmonella enteritidis* via a syringe equipped with an infusion teat. Each chick received one of three inoculation doses ( $1 \times 10^3$ ,  $1 \times 10^5$ , or  $1 \times 10^7$  CFU/chick in 0.25 mL of LB broth). Chicks were monitored twice daily for clinical expression of disease. Morbid chicks were sacrificed. All 99 birds surviving to day 6 were euthanized to determine *Salmonella Enteritidis* burden in spleen tissue and cecal content. The spleen and one cecum were aseptically removed and rinsed with sterile PBS (Ambion, Fisher Int.). The spleen was aseptically weighed and minced with a sterile scalpel.

**Bacteriological analysis:** The spleen (1 g/10 mL) and cecal content (sterile swab collection of content from 3 cm section/10 mL) were enriched by adding the samples into tryptic soy broth (Remel, Fisher Int.) and incubated at 37°C for 18-20 h. One loopful was taken and inoculated into Tetrathionate-iodine broth (BD Diagnostic System, Fisher Int.) and then incubated at 27°C for 20-24 h. After incubation a loopful is

streaked on XLT4 medium plates (BD Diagnostic System, Fisher Int.) containing  $100 \mu\text{g mL}^{-1}$  nalidixic acid and incubated at 37°C for 20-24 h. If colony morphological identification of *Salmonella enteritidis* was questionable, the colony identity was confirmed by *Salmonella* antiserum group D agglutination (BD Diagnostic System, Fisher Int.). The enrichment cultures were held at 4°C during the plus-minus screening.

**Statistical analysis:** All analyses were performed by using SAS 9.2.0. Effects of interactions on survival were tested by the Wald method. The numbers of *Salmonella enteritidis* colonies  $\times 10^{-6}$  were used for analysis. Minor room effects on *Salmonella enteritidis* colony counts were adjusted<sup>[18]</sup>. ANOVA was performed with variables of genetic line, sex and inoculation dose and with all two and three way interactions. Pearson's correlation test was used to determine the correlated relationship between spleen and cecal content *Salmonella Enteritidis* pathogen load. If not otherwise indicated,  $p \leq 0.05$  was considered significant.

## RESULTS

Environmental scans of the birds, laboratory rooms, bedding, water and feed confirmed the absence of environmental *Salmonella enteritidis* prior to the experimental exposure. Four chicks (one from each genetic line) were sacrificed prior to inoculation with *Salmonella enteritidis* and bacterial cultures of spleen and cecal contents confirmed them to be *Salmonella enteritidis* free.

**Layers survival post-challenge:** All 200 chicks were included in the survival analysis. Chicks were categorized as alive or dead (euthanized and spontaneous mortality) at 6 days. The highest inoculation dose of *Salmonella Enteritidis* ( $1 \times 10^7$  CFU/bird) had a significantly ( $p \leq 0.03$ ) lower chick survival than the other two inoculation doses.

Survival for *Salmonella enteritidis* inoculation doses of  $1 \times 10^3$ ,  $1 \times 10^5$  and  $1 \times 10^7$  CFU/bird were 79.2, 72.9 and 54.2%, respectively. There was no significant effect of sex, interaction of sex with either inoculation dose or genetic line, or interaction of inoculation dose with genetic line on chick survival; therefore, inoculation doses and sexes were pooled for analysis of genetic line effect. There was a significant effect ( $p \leq 0.001$ ) of genetic line on chick 6 days survival post *Salmonella enteritidis* challenge. Survival rates were 47.2, 72.2 and 55.6% in the three pure lines (Line 1-3, respectively) and 100% in the experimental three way cross line (Line 4).




Fig. 1: *Salmonella enteritidis* pathogen load of cecal content A. liver (a) and B. spleen (b) by genetic lines

**Bacterial count in ceca and spleen:** The *Salmonella enteritidis* pathogen load was quantified in the 135 chicks that survived to Day 6, at which time the chick numbers were no longer equal for genetic line, sex, inoculation dose, or room. There was no significant effect of inoculation dose or sex on *Salmonella enteritidis* numbers in the cecal content or spleen. Data of all three inoculation doses and both sexes were, therefore, pooled for analysis of genetic line effect and adjusted for room effect. Genetic line had a significant effect ( $p \leq 0.05$ ) on *Salmonella enteritidis* burden in the cecal content but not the spleen tissue, Fig. 1. The *Salmonella enteritidis* pathogen loads in the cecal content and the spleen tissue were not correlated ( $r \leq 0.077$ ,  $p \leq 0.457$ ).

## DISCUSSION

The role of genetics in reducing the *Salmonella enteritidis* pathogen load in young chicks is not clearly understood. In this study, we demonstrated a significant effect ( $p < 0.001$ ) of genetic line on young

layer chick 6 days survival after *Salmonella enteritidis* challenge. Guillot *et al.*<sup>[12]</sup> have also reported varied degrees of genetic line effects on resistance to SE-induced mortality rate of young chicks. They ranked the chicks from broiler, layer and experimental lines according to the level of *Salmonella enteritidis* resistance to IM *Salmonella enteritidis* challenge, as evaluated by lethal dose 50 results at 15 days post-challenge. The three commercial layer lines and one experimental White Leghorn line in the study were all ranked as susceptible (lethal dose  $50 \leq 10^2$  CFU/bird) to *Salmonella enteritidis* relative to the other lines in the study.

The current study demonstrates a wide range of genetic line responses in survival of young layer chicks to oral *Salmonella enteritidis* challenge. The experimental three-way cross had survival significantly superior to each of the pure lines, consistent with expression of heterosis for this trait. Inoculation dose may affect pathogenicity of *Salmonella enteritidis* and the resulting resistance or susceptibility categorization of young chicks. In the current study, inoculation dose had a significant effect on chick 6 days survival but not on *Salmonella enteritidis* burden of the cecal content or spleen tissue of survivors. This result suggests that a wide range of exposure doses might all result in sustained colonization of survivors.

There are conflicting reports regarding the effect of genetic line on resistance to *Salmonella enteritidis* colonization and on which internal organs are differentially colonized<sup>[10,12,15,16]</sup>.

Despite variation in experimental protocols, chicken genetic line is clearly a contributing factor in the current study as well as in previous studies on *Salmonella enteritidis* resistance. In the current study, there were significant genetic line differences in the surviving chicks for *Salmonella enteritidis* burden in the cecal content but not the spleen tissue. Cecal content and spleen *Salmonella enteritidis* burden were not correlated; indicating independent host genetic mechanisms are partly responsible for pathogen load at the two sites. Duchet-Suchaux *et al.*<sup>[5]</sup> reported that high mortality rate in *Salmonella enteritidis* challenged, 1 day old chicks was incompatible with a persistent carrier state, thus suggesting that susceptibility to mortality is directly related to carrier state potential. In contrast, Gast and Beard<sup>[7]</sup> were unable to establish a strong correlation between chick mortality and frequency of production of SE-contaminated eggs. Future *Salmonella enteritidis* contamination control mechanisms in poultry may be basically the same as are currently being utilized, but for long-term sustainability, genetic resistance should be pursued;

Hafez<sup>[13]</sup>. A heritability estimate of 0.20 for *Salmonella enteritidis* burden of enriched cecum culture was reported for 1 week old chicks that were orally inoculated with *Salmonella enteritidis* phage type 4<sup>[2]</sup>.

### CONCLUSION

The current study was able to quantify *Salmonella enteritidis* load from enrichment cultures of the gastrointestinal tract (cecal content) and internal organ (spleen tissue) of young layer chicks.

The current study demonstrated significant differences among genetic lines in resistance to SE-induced mortality and in the *Salmonella enteritidis* burden in cecal contents of young layer chicks.

The genetic line variation exists to effectively choose pure breeder lines that characterized with exhibition of reduced *Salmonella enteritidis* load.

### REFERENCES

1. Beaumont, C., J. Protais, J.F. Guillot, P. Colin, K. Proux, N. Millet and P. Pardon, 1999. Genetic resistance to mortality of day-old chicks and carrier-state of hens after inoculation with *Salmonella enteritidis*. Avian Pathol., 28: 131-135. DOI: 10.1080/03079459994858.
2. Berthelot, F., C. Beaumont, F. Mompert, O. Girard-Santosuosso, P. Pardon and M. Duchet-Suchaux, 1998. Estimated heritability of the resistance to cecal carrier state of *Salmonella enteritidis* in chickens. Poult. Sci., 77: 797-801. PMID: 9628525.
3. Bumstead, N. and P. Barrow, 1993. Resistance to *Salmonella gallinarum*, *S. Pullorum* and *S. Enteritidis* in inbred lines of chickens. Avian Dis., 37: 189-193. PMID: 8452495.
4. Cotter, P.F., R.L. Taylor, Jr. and H. Abplanalp, 1998. B-complex associated immunity to *Salmonella enteritidis* challenge in congenic chickens. Poult. Sci., 77: 1846-1851. PMID: 9872588.
5. Duchet-Suchaux, M., P. Lechopier, J. Marly, P. Bernardet, R. Delaunay and P. Pardon, 1995. Quantification of experimental *Salmonella enteritidis* carrier state in B13 leghorn chicks. Avian Dis., 39: 796-803. PMID: 8719213.
6. Gast, R.K. and C.W. Beard, 1990. Production of *Salmonella enteritidis*-contaminated eggs by experimentally infected hens. Avian Dis., 34: 438-446. PMID: 2196046.
7. Gast, R.K. and C.W. Beard, 1992. Evaluation of chick mortality model for predicting the consequences of *Salmonella enteritidis* infection in layer hens. Poult. Sci., 71: 281-287. PMID: 1546038.
8. Gast, R.K. and P.S. Holt, 1998. Persistence of *Salmonella enteritidis* from one day of age until maturity in experimentally infected layer chickens. Poult. Sci., 77: 1759-1762. PMID: 9872575.
9. Gast, R.K., 1999. Applying Experimental Infection Models to Understand the Pathogenesis, Detection and Control of *Salmonella enterica* Serovar *enteritidis* in Poultry. In: *Salmonella enterica* Serovar *enteritidis* in Humans and Animals, Part III-Chapter 22, Saeed, A.M., R.K. Gast, M.E. Potter and P.G. Wall (Eds.). Iowa State University Press, Ames, IA., pp: 233-243. <http://www.lavoisier.fr/notice/fr270110.html>
10. Girard-Santosuosso, O., P. Menanteau, M. Duchet-Suchaux, F. Berthelot and F. Mompert, *et al.*, 1998. Variability in the resistance of four chicken lines to experimental intravenous infection with *Salmonella enteritidis* phage type 4. Avian Dis., 42: 462-496. PMID: 9777146
11. Gorham, S.L., K. Kadavil, H. Lambert, E. Vaughan, B. Pert and J. Abel, 1991. Persistence of *Salmonella enteritidis* in young chickens. Avian Pathol., 20: 433-437. PMID: 18680039
12. Guillot, J. F., C. Beaumont, F. Bellatif, C. Mouline, F. Lantier, P. Colin and J. Protais, 1995. Comparison of resistance of various poultry lines to infection by *Salmonella enteritidis*. Vet. Res., 26: 81-86. PMID: 7735307.
13. Oliver, S.P., Boor, K.J., Murphy, S.C., and Murinda, S.E., 2009. Poultry meat and food safety: Food safety hazards associated with consumption of raw milk. Foodborne Pathog. Dis., 6: 793-806. PMID: 19737059.
14. Lamont, S.J., 1998. Impact of genetics on disease resistance. Poult. Sci., 77: 1111-1118. PMID: 9706074.
15. Lindell, K.A., A.M. Saeed and G.P. McCabe, 1994. Evaluations of resistance of 4 strains of commercial layer hens to experimental infection with *Salmonella enteritidis* phage type eight. Poult. Sci., 73: 757-762.
16. Protais, J., P. Colin, C. Beaumont, J.F. Guillot, F. Lantier, P. Pardon and G. Bennejean, 1996. Line differences in resistance to *Salmonella enteritidis* PT4 infection. Br. Poult. Sci., 37: 329-339. PMID: 8773842.

17. Schlosser, W.D. and E.D. Ebel, 1998. A risk assessment of *Salmonella enteritidis* from eggs in the United States. USAHA. <http://www.usaha.org/speeches/speech98/s98schol.html>
18. Yonash, N., M.G. Kaiser, E.D. Heller, A. Cahaner and S.J. Lamont, 1999. Major Histocompatibility Complex (MHC) related cDNA probes associated with antibody response in meat-type chickens. *Anim. Genet.*, 30: 92-101. PMID: 10376299.