

## Mathematical Analysis of How the Electrical Training Center Site is Affected by Air Pollution

Bassam A. Al-Helou

Department of Engineering and Technology,  
Faculty of Engineering, Technology-Zarqa University, P.O. Box 132222 Zarqa, 13132, Jordan

Received 2012-07-18, Revised 2012-08-31; Accepted 2012-09-03

### ABSTRACT

The study has come to the result that the monitoring site at the Electrical Training Centre (ETC) at Al-Hashimiyeh town is exposed to grave pollution that goes beyond Jordanian standards of the ambient air. It has, indeed, shown that Sulfur dioxide (SO<sub>2</sub>) exceeds the limits and its seasonal concentration matches forward with the average wind speed. SO<sub>2</sub> concentration reaches its lowest level in the early morning of the day and its highest level at the end of the day. The study has also pointed out that the levels of carbon oxides do not exceed the Jordanian standards of the ambient air and that the mathematical analysis indicators have pointed to the continuing increase of nitrogen oxides due to the increase of calm winds year after year. In fact, these winds do not help the dispersal and decrease of emissions; on the contrary, they increase the emissions concentration, especially in the areas approximate to the pollution sources.

**Keywords:** Air Pollution, Concentration, Sulfur Dioxide, Carbon Oxides, Nitrogen Oxides, Jordanian Standard, Wind Speed, Mathematical Modeling, Wind Direction

### 1. INTRODUCTION

The aim of this study is to identify the degree of air pollution at the ETC resulting from some major pollution sources, such as the Jordanian Petroleum Refinery (JPR) and Al Hussein Thermal Power Plant (HTPP), in addition to the pollution emanating from the emissions of the transport routes and the numerous factories available at Al-Hashimiyeh in Zarqa Governorate.

This exactly means that the pollution affecting the ETC monitoring site is the outcome of the activities of oil refinery and electrical power generation in the region, in addition to the pollution caused by the vehicles movement on the roads (JPRC, 2011; HTPP, 2001; ME, 2010; MiE, 2010; Hamdi *et al.*, 2008). The urban population of Jordan has increased from 64% of the total in 1980 to 74% in 2000 (Reay, 2004). Al-Hashimiyeh area is located north of Zarqa city, 35 km northeast of Amman. It is bounded by Longitude 36° 04' to 39° 09' east and Latitude 32° 04' to 32° 10' north. This town is the most polluted city in Jordan (Odat, 2009; DEAZ, 2005).

It is well-known that exposure for long to pollutants or dust causes health damages. However, health disadvantages would be much greater if man is simultaneously exposed to higher levels of more than


one pollutant, namely SO<sub>2</sub>, NO, NO<sub>2</sub> and dust. Health damages especially affect children, elders and people stricken by allergy, chest diseases, asthma, (Peters *et al.*, 1997; WHO, 2005; Mage and Zali, 1992; Seaton *et al.*, 1995; APPA, 1998).

The following pollutants are monitored: sulfur dioxide, carbon dioxide, carbon monoxide and nitrogen oxides.

Sulfur dioxide (SO<sub>2</sub>) is a poisonous gas. It agitates the tissues of eyes, noses, pharynxes and lungs. So once absorbed by a human body, SO<sub>2</sub> operates as an internal toxin that would possibly impair the nervous and respiratory systems. SO<sub>2</sub> is a colorless gas and so cannot be visible when it infiltrates. Neither can it be seen in a stormy weather. It moves as an invisible cloud, usually in the wind direction. It is a heavier gas than air and tends to fall on low lands, such as ditches, drainages and holes, where it goes well as a fatal gas. The degree of SO<sub>2</sub> effect depends on its concentration in air, as well as on the victim's allergy and exposure duration. So if SO<sub>2</sub> condensation in air amounts to 100 ppm, it then develops eye and throat inflammation, headache, nausea, cough and the olfaction becomes useless within 3-15 min (Burki *et al.*, 1994; Jol and Kielland, 1997; Lefohn and Shadwick, 1998; Kalabokas *et al.*, 1999; WHO, 1979). The degree of SO<sub>2</sub> condensation in air is based on the quantity of fuel consumption and the ratio of sulfur in there.

**Table 1.** Some Pollutants Concentration Limits in Jordan, Standard 1140/2006 Jordanian Standards, 2006

Pollutant	Average exposure Time	Concentration Limit (ppm)	Number of allowed exceeds
SO <sub>2</sub>	1 h	0.30	Three times within any 12 months
SO <sub>2</sub>	24 h	0.14	One time within a year
SO <sub>2</sub>	Annual	0.04	-
CO	1 h	26.00	Three times within any 12 months
CO	8 h	9.00	Three times within any 12 months
NO <sub>2</sub>	1 h	0.21	Three times within any 12 months
NO <sub>2</sub>	24 h	0.08	Three times within any 12 months
NO <sub>2</sub>	yearly	0.05	-



**Fig. 1.** The geographical locations of the monitoring site and the pollution sources

Carbon Monoxide (CO) is a gas with no color or smell. It emanates from the incomplete combustion interaction of hydrocarbon fuel. However, the main source of this gas is the traffic movement on road, especially the benzene-operating vehicles. In fact, the CO affecting danger on man lies in its potential of creating carboxyl hemoglobin in man's blood, which really causes man's blood inability to carry oxygen to his organs. As such, it exposes to further danger those people who suffer from the inefficiency of extending oxygen to heart and brain (e.g., patients of coronary artery) (Al-Helou and Sinokrot, 2004; WHO, 2005; Brunekreef and Holgate, 2002; DH, 2005).

Nitrogen Oxides (NO<sub>x</sub>), the term used to describe the sum of NO and NO<sub>2</sub>, play a major role in the formation

of tropospheric ozone can also contribute to the formation of acid rain (Jol and Kielland, 1997; Mbuyi, 2003). Long-term exposures (e.g., one year) to nitrogen oxide concentrations may damage forest and vegetation (Fally *et al.*, 1995). Nitrogen oxides are formed in high-temperature combustion processes, mainly in petrol and diesel vehicles (Monroe and Li, 2002). The pollutants have been checked in this study within the limits determined by the Jordanian standards of the ambient air (Table 1).

### 1.1. Air Monitoring Site

Al-Hashimiyeh ETC Monitoring site lies only 2 kms to the south-east of JPR, 0.25 km south-east HTFP and about 5.5 kms south-west Samra Wastewater Treatment Station (SWTS) (Fig. 1).

## 1.2. Monitoring Procedures

The air samples are analyzed for Carbon Oxides (CO and CO<sub>2</sub>) concentrations using analyzer Infrared Fluorescence, Nitrogen Oxides (NO, NO<sub>x</sub> and NO<sub>2</sub>) concentrations using analyzer Chemiluminescence. The SO<sub>2</sub> analyze done using Ultra-Violet Fluorescence. The Wind speed and direction are measured using a Metro-Unit. Additionally, meteorological data are collected at the Zarqa weather station. Wind speed, wind direction, ambient temperature, barometric pressure and solar radiation are measured hourly.

## 1.3. Sulfur Dioxide (SO<sub>2</sub>)

ETC is affected by the SO<sub>2</sub> emission from the JPR and HTPP stations. Both stations constitute to be the consistent sources of sulfur dioxide that emanates from fuel combustion and oil refinery in Al-Hashimyeh area. In addition, the fact that sulfur dioxide contains high degree of sulfur which is carried by west and North West winds, at a range of about 950 ppm, although the European Standard set by 50 ppm (OPEC, 2005).

**Figure 2** shows that on January 6, 2010, the monitoring results of SO<sub>2</sub> in that year had registered SO<sub>2</sub>'s highest level of 1.766 ppm and those 783 cases of exceeding the Jordanian standard limits 1140/2006 were, in average, 0.3ppm, i.e. by 8.93%. Moreover, these monitoring results had shown that the max daily average was 0.623ppm and that the exceeding daily averages of those limits by 0.14 ppm were 68, i.e., 18.634%.

**Figure 3** shows the direction of wind distribution at the time of the daily exceeding of the Jordanian standards. This exactly clarifies the great pollution danger by SO<sub>2</sub> which the monitoring site is exposed to as west has winds increased to 20.1%, North West winds to 21.7% and calm winds to 30.7%. All together, those winds amounted to 72.5% a very high degree of wind accumulation that permanently exposes the monitoring site to SO<sub>2</sub> pollution.

**Table 2** shows the max recorded average in each of the monitored months. The average exceeds were recorded throughout those monitored months. The table also shows that calm winds had recorded high proportions of pollution during those months as a whole, which actually means that these winds do not disperse or reduce the emissions produced by HTPP. Rather they increase their concentration. This table clearly shows that the west and North West winds are relatively abundant and are capable to carry the emissions from JPR and HTPP to the monitoring site and subsequently increased SO<sub>2</sub> concentration there.

**Figure 4** shows the monthly average of SO<sub>2</sub> concentrations; its lowest level was in March and the max in June at a rate of 0.010 and 0.159 ppm respectively. It also clearly proves an exceeding to the 0.14 ppm limit of the Jordanian standards.

**Figure 5** shows the annual and seasonal average of SO<sub>2</sub> concentrations and the wind speed in each hour of the day. It can be noted that the SO<sub>2</sub> concentration fits forward with the wind speed average. It actually reaches its lowest level in the early morning and to its highest level at the end of the day. Air tranquility plays an important role in the distance which the wind may reach and in its concentration in the surrounding air as well.

Comparing the 2010 monitoring results with the previous ones, i.e. the results measured in: 2004-2005, 2005-2006, 2006-2007, 2007-2008 and in 2009 it becomes clearly obvious that 2010 results are the highest since 2006-2007 and the numbers of excessive averages were between 246 and 951. The highest excess was, indeed, in 2004-2005, while the highest average values ranged between 0.813 and 6.545ppm, (the latter being registered in 2004-2005). It is also noted that the percentage of the excess averages began their progressive increase since 2007-2008 up to 2010. In 2006, they recorded an increase of 3.13 and 8.93% in 2010 (**Fig. 6**).


**Figure 7** shows the daily averages of SO<sub>2</sub> concentration during the period of monitoring from 2004 to 2010. The maximum daily average was recorded in 2009 at 1.192ppm and that the number of excesses ranged between 24-115, the highest of which was in 2004-2005 and the lowest was in the monitoring period of 2006-2007.

The above- mentioned analysis points clearly to the excesses, in the monitoring site, of the Jordanian standard limits of SO<sub>2</sub> and that those excesses were emphatically noted throughout the years of monitoring. The percentages of those excesses underline the need for taking necessary measures of reducing, as much as possible, the sulfur quantity in fuel before it is burnt and for installing systems to treat the discharged air from chimneys before it darts into the ambient air.


## 1.4. Carbon Oxides (CO, CO<sub>2</sub>)

HTPP and JPR constitute the main and constant sources of CO and CO<sub>2</sub> emission that results from the fuel combustion and refining. CO<sub>2</sub> is generated from the full process of combustion, whereas CO is generated from the incomplete process of burning, especially when starting operation and halting the units of production.


The results of 2010 monitoring show that no excess to the CO average limits as determined by the Jordanian standards at 26 ppm, knowing that Jordanian standards contain no item regarding CO<sub>2</sub>. **Table 3** shows the maximum hourly averages of CO and CO<sub>2</sub> concentrations. It also shows the maximum 8-hours averages of CO concentrations. No excess to limits as determined by the Jordanian standards at 9 ppm was noted.




**Fig. 2.** Daily averages and max hourly averages concentrations of SO<sub>2</sub> at ETC




**Fig. 3.** Wind direction distribution during days at which SO<sub>2</sub> daily average concentration exceeded JS limit at ETC




**Fig. 4.** Monthly averages concentrations of SO<sub>2</sub> at ETC




**Fig. 5.** Averages diurnal concentrations of SO<sub>2</sub> and wind speed diurnal average (knot)




**Fig. 6.** Maximum hourly averages concentrations of SO<sub>2</sub> and No. of Exceeds at ETC (2004-2010)




**Fig. 7.** Maximum Daily averages concentrations of SO<sub>2</sub> and No. of Exceeds at ETC (2004-2010)



**Fig. 8.** Maximum hourly averages concentrations of NO, NO<sub>2</sub> and NO<sub>x</sub> at ETC



**Fig. 9.** Daily averages concentrations of NO, NO<sub>2</sub> and NO<sub>x</sub> at ETC



**Fig. 10.** Averages diurnal concentrations of NO<sub>x</sub> and wind speed diurnal average (knot) at ETC (ME, 2010)




Fig. 11. Wind direction distribution during days at which NO<sub>2</sub> daily average concentration exceeded JS limit of 0.08 ppm at ETC




Fig. 12. Maximum hourly averages concentrations of NO, NO<sub>2</sub> and NO<sub>x</sub> at ETC (2007-2010)




Fig. 13. Mathematical model of distribution wind speed (2004-2010)

**Table 2.** Maximum Hourly averages of SO<sub>2</sub> concentration (ppm) and percentage of wind direction

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug.	Sep.	Oct.	Nov.	Dec.
1.766		0.779	0.651	0.615	0.726	1.012	0.930	0.957	0.734	0.704	0.831	0.766
Calm %												
43.2		66.900	67.000	-	44.100	28.500	18.900	18.000	25.900	38.900	66.200	59.000
NW %												
5.5		6.300	5.500	-	18.400	25.500	45.600	11.300	0.000	23.200	3.000	4.800
W %												
18.4		20.900	16.500	-	35.000	33.900	11.900	0.000	0.000	6.500	0.900	2.400

**Table 3.** Maximum Hourly averages of CO, CO<sub>2</sub> and maximum 8-hours averages of CO concentrations (ppm)

Pollutant	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug.	Sep.	Oct.	Nov.	Dec.
CO	0.397	1.487	0.546	--	2.963	2.938	3.404	3.463	0.807	0.487	0.674	2.839
CO <sub>2</sub>	553.000	543.000	470.000	--	802.000	865.000	941.000	994.000	473.000	372.000	410.000	445.000
Maximum 8-hours averages of CO concentration (ppm)												
CO	0.386	1.286	0.325	--	1.149	1.08	1.708	1.370	0.474	0.160	0.351	1.065

### 1.5. Oxides (NO, NO<sub>2</sub>, NO<sub>x</sub>)

The results of 2010 monitoring show that 15 excesses to the maximum hourly average of NO<sub>2</sub>, as defined by the Jordanian standards at 0.21 ppm. **Figure 8** shows that the maximum averages of NO<sub>x</sub>, NO and NO<sub>2</sub> were 0.465, 0.879 and 0.895 ppm respectively. The first and second readings of the recording were carried in early September, while the third reading was done in February. **Figure 9** shows the daily averages of Nitrogen oxides, where the maximum daily average of NO<sub>x</sub> was 0.129 ppm, NO 0.059 ppm and NO<sub>2</sub> 0.104 ppm. Three excesses of NO<sub>2</sub> to Jordanian standards were recorded.

**Figure 10** explains the annual and seasonal averages of NO<sub>x</sub> concentration, as well as the wind speed per knot (knot = 1m/s). It can be, accordingly, said that there is progressive harmony with the wind speed. Both factors attain their lowest level in the early morning and highest levels at the end of the day. Such a phenomenon can be identified by analyzing a number of atmospheric factors that play a role in carrying emission to the monitoring sites. In addition, the speed and direction of the wind actually play an outstanding role in determining the distance that could be reached by emissions, as well as in the concentration of these emissions in the air.

**Figure 11** shows the distribution of wind direction wherever excess to the daily average, as defined by Jordanian standards, is recorded at the monitoring site. The above emphasized conclusions confirm that Al-Hashimiyeh area, lying to the east of JPR and HTPP, are gravely exposed to NO<sub>2</sub> pollution when west and North West winds intensify. All together, winds accumulate to 63.2%. **Figure 12** identifies the discrepancy between the results of 2010 monitoring and those of the previous years. It shows the values of the average nitrogen oxides between 2007 and 2010. The results prove the increase in the concentrations of these gases in 2010 as compared to the previous years. To understand this remarkable

increase, the wind speed at Al-Hashimiyeh area was analyzed for the years 2004-2010, after being divided in five speeds (0-2 knots). Eventually, those winds were labeled as calm winds, while the other winds were sorted out as 2-5, 5-10, 10-15 and >20 knots.

Analysis showed that calm wind was continuously increasing since 2004 up to 2010 and that the increase in 2010 was the highest compared to previous years. Indeed, this wind helps neither the dispersal nor the reduction of gas emissions; on the contrary, it increases its concentration, especially in approximate areas to pollution sources. Math predictions warn of new increases during the coming years, owing to the increase in calm wind speed a year after year (**Fig. 13**).

Notice that math analysis proves that the mat model of calm winds increase is **Equation 1**:

$$y = 3.01857x - 6361.5 \quad (1)$$

x: stands for the year when calm wind is counted for, whereas the correlation value  $R^2 = 0.8328$ , which is not low.

The math analyses also show that the (2-5 knots) and (5-10 knots) correspondently face decrease, as those analyses prove in the following two equations respectively **Equation 2** and 3:

$$y = -2.8986x + 5847.2 \quad (2)$$

$$y = -0.3579x + 747.2 \quad (3)$$

The correlation value of  $R^2 = 0.8469$  and  $R^2 = 0.2945$ . The latter value is not much. The high wind speeds 15-20 knots and >20 knots have not been analyzed mathematically, due to their low proportion during the years of monitoring.

## 2. CONCLUSION

This study has clearly shown that Al-Hashimiyeh area is overwhelmed by calm winds and subsequently exposed to increasing gas pollution in the ETC. It has also come out with the fact that SO<sub>2</sub> and nitrogen oxides have exceeded the limits of Jordanian standards, whereas carbon oxides maintained their low levels, with no record of violation of the standard limits. The study recommends the replacement of heavy fuel, which contains high degree of sulfur, by natural gas for power generation at HTPP.

Math analysis has shown the predicted the NO<sub>x</sub> increased pollution in the coming years, owing to the increased level of calm winds year after year. Such a fact helps the increase of these gas concentrations, especially in the approximate areas to pollution sources.

## 3. ACKNOWLEDGEMENT

The researchers would like to thank Jordanian Ministry of Environment for providing the air quality data necessary for this study.

## 4. REFERENCES

- Al-Helou, B. and B. Sinokrot, 2004. Air quality comparison between industrial and residential communities. *Environ. Ecol.*, 22: 586-591.
- APPA, 1998. La pollution de l'air: sources, effets, prevention, Association pour la Prevention de la Pollution Atmospherique. Ville de Besancon-Service Hygiene Sante, France.
- Brunekreef, B. and S.T. Holgate, 2002. Air pollution and health. *Lancet*, 360: 1233-1242. DOI: 10.1016/S0140-6736(02)11274-8
- Burki, D., J. Keller and B.E. Ga, 1994. Air pollution. Part 2: Concentrations of major air pollutants in Switzerland and trans boundary contributions. Paul Scherrer Institute, Switzerland.
- DEAZ, 2005. Reports about al-hashimeya area. Directorate of Environment in Al-Zarqa.
- DH, 2005. Hydrogen Sulfide Chemical Information Sheet. Department of Health.
- Fally, S., D. Joannes, D. Leduc and M. Scharll, 1995. La pollution, ses effets et ses Costs Special en region de Bruxelles-capitale. University Libre de Bruxelles, Belgique.
- Hamdi, M.R., A. Bdour and Z. Tarawneh, 2008. Diesel quality in Jordan: Impacts of vehicular and industrial emissions on urban air quality. *Environ. Eng. Sci.*, 25: 1333-1344. DOI: 10.1089/ees.2007.0234
- HTPP, 2001. Report for air quality monitoring for the selected site of assamra power project at Al-Hashimeya.
- Jol, A. and G. Kielland, 1997. Air pollution in Europe. European Environ. Agency.
- JPRC, 2011. Jordan Petroleum Refinery Company (JOPT)-Financial and strategic SWOT analysis review. MarketResearch.
- Kalabokas, P.D., L.G. Viras and C.C. Repapis, 1999. Analysis of the 11-year record (1987-1997) of air pollution measurements in Athens, Greece. part I: primary air pollutants. *Global Nest: Int. J.*, 1: 157-167.
- Lefohn, A.S. and D.S. Shadwick, 1998. Ozone, sulfur dioxide and nitrogen dioxide trends at rural sites located in the united states. *Atmospheric Environ.*, 25: 491-501. DOI: 10.1016/0960-1686(91)90321-W
- Mage, D. and O. Zali, 1992. Motor Vehicle Air Pollution: Public Health Impact and Control Measures. World Health Organization Division of Environmental Health. Switzerland.
- Mbuyi, M.H.T., 2003. Air pollution at the bily kriz mountains in the czech republic. *Int. J. Environ. Pollut.*, 19: 1-10.
- ME, 2010. Study the levels of pollutants in the area of Al-Hashimeya. Unpublished Annual Report, Ministry of Environment.
- Monroe, D. and W. Li, 2002. Desulfation dynamics of NO<sub>x</sub> storage catalysts. Proceedings of the SAE Power Train and Fluid Systems Conference and Exhibition, (PTFSCE' 02), SAE International, San Diego, CA, USA. DOI: 10.4271/2002-01-2886
- Odat, S., 2009. Diurnal and seasonal variation of air pollution at Al-Hashimeya Town, Jordan *J. Earth Environ. Sci.*, 2: 1-6.
- OPEC, 2005. Guide for petroleum product specifications. OPEC Organization.
- Peters, A., H.E. Wichmann, T. Tuch J. Heinrich and J. Heyder *et al.*, 1997. Respiratory effects are associated with the number of ultrafine particles. *Am. J. Respiratory Critical Care Med.*, 155: 1376-1383.
- Reay, M.E., 2004. Air quality and atmospheric pollution in the Arab Region. University of Alexandria.
- Seaton, A., D. Godden, W. MacNee and K. Donaldson, 1995. Particulate air pollution and acute health effects. *Lancet*, 345: 176-178. DOI: 10.1016/S0140-6736(95)90173-6
- WHO, 1979. Sulphur oxides and suspended particulate matter, environmental health criteria 8. World Health Organization, Geneva.
- WHO, 2005. WHO Air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide. World Health Organization.