

Trace Elements Distribution in the Brain of Stressed Rats

¹Zerrouk Kamal, ²Mohamed Najimi, ²Mohammed Chigr,
¹Mariama El Ouahli, ¹Ghizlane Er-Raoui and ¹Fatiha Chigr

¹Genie Biologique Laboratory, Faculty of Sciences and Techniques,
²Département of Chemistry and Environnement, Faculty of Sciences and Techniques,
Sultan Moulay Slimane University, Béni-Mellal, Morocco

Received 2012-03-12, Revised 2012-09-02; Accepted 2012-09-14

ABSTRACT

Because of their ability to modulate the Gamma Aminobutyric Acid (GABA) a receptor complex, the principal aim of the current work is to assess two essential heavy metals: Iron (Fe) and zinc (Zn) by atomic absorption in different brain areas of stressed rats. To do so, we investigated the effect of acute immobilization stress (single 1-hour session) on the distribution and the densities of GABA_A receptors as well as the concentrations of Zn and Fe in several rat brain regions of the stressed rats. Animals were randomly assigned to either control or stress conditions and changes in specific binding of the GABA_A receptor as labelled with T-Butylbicyclophosphonothionate (TBPS) (ligand useful for GABA_A receptor) were assessed by in vitro quantitative autoradiography with the aid of a computer-assisted image analysis system whereas the assessment of Fe and Zn concentrations was done by atomic absorption spectrophotometry. Exposure to 1h immobilization stress led to a significant increase in [³⁵S]-TBPS binding site density in stressed rats compared to controls (30-40% increase in cortex, hypothalamus, hippocampus and substantia nigra). In the other analyzed brain structures, specific binding of [³⁵S]-TBPS remained unchanged in stressed rats. The spectrophotometer analysis showed significant decrease in Zn levels in the whole forebrain structures as well as the mesencephalon of stressed rats. The striking differences are noticed in hippocampus and mesencephalon. Furthermore, Fe endogenous concentrations display similar pattern following stress. The present study demonstrates that immobilization stress induces an increase in GABA_A receptors concomitant to a reduction of Zn and Fe content in the stress sensitive rat brain structures. Besides supporting the alteration of the modulatory function occurring at the GABA_A receptor level after stress, our data also reveal that the measured brain concentrations of the investigated heavy metals remain not sufficient to efficiently modulate the activity of with efficacy such complex receptor. This could explain the higher densities of GABA_A receptors observed after acute stress.

Keywords: Zinc, Iron, GABA_A, Stress, Brain

1. INTRODUCTION

Gamma Aminobutyric Acid (GABA) A receptors are ligand-gated ion channels, mediating fast synaptic inhibition in the Central Nervous System (CNS) (Mody, 2012; Farrant and Nusser, 2005). These receptors are the targets of a variety of pharmacologically and clinically important drugs such as benzodiazepines, depressant

barbiturates and neuroactive steroids (Smith *et al.*, 2007). Beside these major modulators, some metal cations are able to inhibit the GABA response of neurons in a variety of organisms (Chen and Yung, 2006). Among them zinc and at a lesser extent iron, have been reported to be the most potent (Rahman, 2007). Zn²⁺ ions inhibit GABA_A receptor function by an allosteric mechanism that is critically dependent on the receptor subunit

Corresponding Author: Fatiha Chigr, Laboratoire de Génie Biologique, Faculté des Sciences et Techniques, PO. Box: 523,
Sultan Moulay Slimane University, Béni-Mellal 23000, Morocco Tel: 212-523485112 Fax: 212-523485201

composition: $\alpha\beta$ -subunit combinations show the highest sensitivity to Zn^{2+} (Krishek *et al.*, 1998; Saxena and Macdonald, 1996). Zn^{2+} antagonism is also affected by α - subunit (Draguhn *et al.*, 1990; Smart *et al.*, 1991; White and Gurley, 1995). The inhibitory effect of Zn^{2+} is reduced by exchanging $\alpha 1$ -subunit for $\alpha 2$ - or $\alpha 3$ -subunits. Finally, Zn^{2+} as picrotoxin and pregnenolone sulfate, two endogenous modulators of GABA_A receptor, exert their effect directly binding in the ion channel (Hosie *et al.*, 2003; Krishek *et al.*, 1996). The mechanism involving the Gamma-Aminobutyric Acid (GABA) in stress is well known (Foddi *et al.*, 1997). Indeed, many reports suggest that GABA_A receptors may mediate responses to stressors and the physiological control of stress (Drugan *et al.*, 1993; Gunn *et al.*, 2011). Stress has also been shown to be associated with alterations in the capacity of ligand binding to the GABA_A receptor complex (Concas *et al.*, 1993; Barbaccia *et al.*, 1996; Serra *et al.*, 2000). GABA_A receptors density and modulation have been documented to significantly increase in density following the application of acute stress (Chigr *et al.*, 2001a; 2001b; 2002) probably as a consequence of an alteration of endogenous modulator levels. recently, several stress paradigms have been shown to induce significant reduction in serum zinc concentration (Cieslic *et al.*, 2011). Fe^{2+} was also reported to interact with Zn^{2+} especially in stressful situations. Psychological stress may cause decreased Fe^{2+} absorption and iron redistribution in body resulting in low iron concentrations in the bone marrow, an effect counterbalanced by zinc (Li *et al.*, 2012). Therefore, the aim of the present study was to evaluate the changes that may occur for the concentrations of the two heavy metals Zn^{2+} and Fe^{2+} in the brain of stressed rats and to correlate such alterations to the [³⁵S]-TBPS binding on the benzodiazepine/GABA chloride ionophore receptor complex.

2. MATERIALS AND METHODS

2.1. *In vitro* Autoradiography Procedures

The experiments were performed on male Wistar rats (n = 16, weighting initially 180-200 g). The animals were housed (3-4 per standard cage) under controlled laboratory conditions (12-h light/dark cycle, constant temperature: 22±2°C and humidity: 60±2%) with food and water freely available. These conditions are rigorously respected to avoid any other stress situations. All experimental procedures were conducted between 8:00 a.m. and 10:00 p.m. Animals were handled for 5 days (20 min day⁻¹) to habituate them to the stress of handling and assigned to either a control (no stress) (n =

8) or immobilization stress (one 1-h stress session) (n = 8). At the time of experiments, the rats of the experimental group were picked up from their cages and were immediately subjected to stress immobilization. Immobilization stress conducted under this protocol was done as previously described by Kvetnansky and Mikulaj (1970) with slight modifications. Stress was applied by placing the rats on a piece of wood and the front and hind legs of the rats were immobilized with plastic collars and adhesive tape on the boards of the wood. This allowed normal breathing and only restricted movements were possible (free movement of the head and tail). The animals were kept immobilized for 1 h. These experimental measures minimized the pain of the rats. Ten minutes after the single immobilization stress session animals were killed by decapitation. Control rats were killed after removal from the home cages without prior manipulations. In all experiments, the immobilization procedure and sacrifice were performed between 10 and 12 h, to avoid possible circadian variations in the receptor function. After sacrifice, the brains were removed, snap-frozen in dry ice and kept at -80°C until sectioning. Serial coronal 20 μm thick sections from each brain were cut at -20°C in a cryostat (Figocut 2880, Reichert Jung), collected onto 2% gelatin-coated slides and kept at -20°C until used. [³⁵S] T-Butylbicyclophosphonothionate (TBPS) autoradiography was conducted as we previously described (Chigr *et al.*, 2002). Briefly, after a preincubation step of 30 min in 50 mM Tris-HCl, pH 7.4, brain slices were incubated with 3 nM [³⁵S]TBPS (100-140 Ci/mmol; NEN, Boston, MA) in 50 mM Tris-HCl buffer, pH 7.4, containing 500 mM NaCl and 10⁻⁴ M ascorbic acid, for 3 h at room temperature, in the absence (total binding) or in the presence of the unlabelled compound (picrotoxin, Sigma Chemical Co., St Louis, MO). The slices were then rinsed, dried out in ambient air, transferred to cardboard film cassettes along with slides containing labelled plastic standard (Amersham, Les Ulis, France) and apposed onto [³H]Hyperilm (Amersham). After an exposure period of 36 h at 4°C, the films were developed and fixed. The obtained autoradiographic labelling was quantified in many brainstem structures by computerized densitometry (Biocom, Les Ulis, France). Data were measured in relative Optical Density (OD) units for displacement studies. Drug-inhibited [³⁵S]-TBPS binding was then plotted as a percentage of the specific binding obtained in the absence of the drugs. For non-displacement studies, the optical densities of each structure, were measured from the exposed film and converted into femtomoles/mg (fmol/mg), based on calibration curves of standards with known radioactivity (Miller, 1991). Specific binding was defined as total binding minus non-specific binding (binding in the

presence of 10^{-5} M picrotoxin). For all experiments, each point was the mean of measurements in five to eight sections for each individual animal, repeated with at least five animals for either control or stressed rats. The data were compared by Analysis Of Variance (ANOVA) followed by Scheffe's test. A P value of <0.05 was considered statistically significant.

2.2. Brain Zinc and Iron Determination

For zinc and iron determination, two other groups of rats ($n = 6$, for each group) assigned to stress or not, were euthanized by decapitation 24 h after the session of stress. Rat brains were then collected and dissected to obtain: frontal cortex, hippocampus, thalamus, hypothalamus and mesencephalon the brain regions dissected were frozen and stored at -20°C until analysis. Each sample was wet-digested with nitric acid and hydrogen peroxide (microwave digestion). The zinc and iron concentrations were determined using flame atomic absorption spectrometry. The equipment used was a Thermo scientific, ICE 3000 Spectrophotometer with deuterium background correction (air flow- 4.2 L min^{-1} , acetylene flow- 1.2 L min^{-1} , analytical wavelength- 213.9 nm). Relative Standard Deviation (RSD) of the method (the whole analytical procedure: digestion + metal determination) did not exceed 2.4%. Mean recovery of zinc and for iron was 99%. The aqueous standard solutions were obtained from Zn (Zinc Standard Solution, Merck, Germany) and FeNO_3 (Sigma Aldrich). The data obtained were analysed by Analysis Of Variance (ANOVA) followed by Scheffe's test. A P value of <0.05 was considered statistically significant.

3. RESULTS

The distribution of GABA_A receptors, using $[^{35}\text{S}]\text{-TBPS}$ as a ligand, showed a heterogeneous distribution in both stressed and unstressed rat brain. However, the quantification of the radioautographic labeling revealed the presence of significant differences in the density of $[^{35}\text{S}]\text{-TBPS}$ binding sites. As shown in **Fig. 1**, high significant densities were principally found in the forebrain of stressed rats (30-40% increase). This includes hypothalamus, hippocampus and cerebral cortex. Detailed analysis, demonstrates that significant differences were confirmed in areas known to be sensitive to stress, such as layer IV of the cerebral cortex, the dentate gyrus in hippocampus and the paraventricular nucleus in the hypothalamus. In all these structures, the $[^{35}\text{S}]\text{-TBPS}$ binding (2 nM) was increased significantly ($p < 0.05$, $n = 8$ for each group and 6-8 measurements for each rat) after 1 h session of immobilization stress. The other structures belonging to cerebral cortex (as layer VI), thalamus (for example paraventricular nucleus) and hypothalamus did not display any significant differences in TBPS binding. In the brainstem, the only striking difference was observed in the substantia nigra at the mesencephalic level in accordance with our previous results ($p < 0.05$, $n = 8$ for each group and 6-8 measurements for each rat) following 1 h of immobilization stress session (Chigr *et al.*, 2001a; 2001b). The binding observed in other structures as collucili and substantia gelatinosa was also not affected after stress (**Fig. 1**).




Fig. 1. Effect of 1h-immobilization on TBPS binding sites in different brain structures as determined by *in vitro* quantitative autoradiography




Fig. 2. Effect of 1h-immobilization on endogenous zinc concentrations in different brain structures as assessed by atomic absorption spectrophotometry




Fig. 3. Effect of 1h-immobilization on endogenous iron concentrations in different brain structures as assessed by atomic absorption spectrophotometry

In other stressed and unstressed rat groups (n = 6, for each group), we in parallel measured the endogenous brain levels of zinc and iron. The spectrophotometric analysis showed that the mean concentrations of Zn²⁺ are significantly decreased in all the sensitive structures reported to be sensitive to stress (displaying high TBPS binding). This effect on endogenous zinc concentrations is well displayed in forebrain structures, principally hippocampus; where the lowest concentrations were observed (25-65% decrease in stressed rats compared to controls, p<0.05, n = 6 for each experimental group; **Fig. 2**). A significant decrease in Zn concentration was also measured in the mesencephalon (around 55% of decrease

in stressed rats compared to controls, p<0.05, **Fig. 2**). Thalamus, cerebellum and pontine which did not display increase in TBPS binding, also did not show any significant fluctuation in zinc concentration. Fe levels analysis showed significant decrease in the forebrain region and mesencephalon of the stressed rats (20-35% decrease in stressed rats compared to controls, compared to controls, p<0.05, **Fig. 3**). Similarly to endogenous zinc concentrations, stress does not affect significantly the endogenous iron concentrations in cerebellum and pontine (**Fig. 3**). Globally, the brain areas sensitive to stress, show both alterations in GABA_A receptors and zinc and iron concentrations.

4. DISCUSSION

The present study examined the effects of immobilization stress on the brain concentrations of two essential heavy metals, i.e., zinc and iron, i.e., zinc and iron proposed as endogenous modulators of the GABA_A receptor complex (for review Rahman, 2007). We firstly assessed the influence of the stress paradigm we used on the *in vitro* autoradiography [³⁵S]-TBPS binding, a ligand that binds at or near the chloride channel of the GABA_A receptor complex (Atack *et al.*, 2007). The experiments presented herein revealed that the GABA_A receptor complex as labeled by [³⁵S]-TBPS was affected by a single exposure to stress (1-h immobilization) in the rat brain. These findings also show that stress effects were specifically noticed at specific regions of the rat brain. Indeed, some regions of the cerebral cortex, hippocampus, hypothalamus and substantia nigra were the structures sensitive to stress immobilization. No significant differences were observed in the other structures. Thereafter, we investigated the effect of acute immobilization stress on the endogenous metals concentration and demonstrated that a significant decrease in their brain levels. Unfortunately, while the autoradiography technique allows the fine anatomical determination of structures concerned by stress, this is not possible with spectrophotometry absorption technique that necessitates important brain punches. Thus, it was impossible to determine the Zn values for the layer IV of cerebral cortex or for the paraventricular nucleus of the hypothalamus. The values of metal concentrations concern the whole structure. This could not allow us to determine if other subdivisions in these structures account for the decrease observed or not.

The findings concerning stress effect on GABA_A receptors density reinforce the idea that the GABAergic system seems to be differentially affected by stress according to the brain area studied (Losada, 1989). The enhancement of [³⁵S]-TBPS binding could be explained by the reduced concentrations of endogenous GABA, occurring specifically after stress. Such responses may, in turn enhanced the GABA_A receptor sensitivity to other endogenous ligands. In addition, the present findings are different from previous data showing decreases in the binding of [³⁵S]-TBPS in many brain structures after stress (Drugan *et al.*, 1993). The discrepancy may be due to several procedural factors or to the stress type applied. The mechanism whereby stress alters TBPS binding sites also allows other hypotheses. Thus, it is possible that the individual brain region differences observed could be due to the subunit heterogenous distribution in these brain structures (Fritschy and Mohler, 1995). As a

consequence, the potency and magnitude of the stress effect could be influenced by the subunit composition of the GABA_A receptor. In this sense, the selectivity of the stress effect for certain sub-unit assemblies could not be excluded. The observed alterations might also result from changes in brain neuroactive steroid levels known to exert anxiolytic (Bitran *et al.*, 1991) and anti-stress (Barbaccia *et al.*, 1996) effects in brain. It is likely that by reducing the synthesis and the release of neurosteroids, acute stress will enhance the [³⁵S]-TBPS binding. Finally, it is not excluded that the observed changes could be due to the fact that in the stress paradigm used, the concentration range over which several modulators were able to counteract the impairment in GABAergic transmission (Barbaccia *et al.*, 1996; Concas *et al.*, 1998) has not yet been reached in the brain structures sensitive to stress. This could explain why the low concentrations of zinc obtained after stress, are not able to efficiently inhibit GABA_A receptor function. Indeed, the transition metal ion Zn²⁺ is concentrated into zinc-containing neurons in the CNS and is thought to be involved in modulating the sensitivity of GABA and glutamate receptors to their respective neurotransmitters (Frederickson, 1989; Frederickson and Bush, 2001). Zinc is known to inhibit and to reduce the spontaneous and the miniature inhibitory postsynaptic currents (Chen and Yung, 2006). The decrease in zinc concentrations leads probably to the alteration of the capacity of the metal to modulate of the tonic GABAergic inhibition (Smart *et al.*, 2004). Among possible other explanations for this negative result is that synaptically released Zn²⁺ is confined to excitatory synapses and that Zn²⁺ spillover to GABAergic synapses is not sufficient to modulate the GABA_A receptors present (Ruiz *et al.*, 2003). Zinc as other active endogenous modulators of the GABA_A receptor, is negatively affected by stress which reduces its endogenous concentrations and could results probably in uncoupling of zinc and TBPS binding sites in the GABA_A receptor complex. As a consequence, the affinity of zinc towards its binding site in the GABA_A receptor complex will decrease. Finally it is not excluded that some of these changes may be explained by altered expression of subunits that affect the sensitivity of receptors to Zn²⁺. Zinc is known to exhibit antidepressant like activity (Hosie *et al.*, 2003) and any reduction in its endogenous levels leads to anxiety states (Huang *et al.*, 2012). This is an agreement with the anxiety behavior observed in stressed rats (Charrier *et al.*, 2006). Hence, supplementation of zinc was suggested to abolish or to reduce the anxiety-like state observed

following stressful conditions (Cope *et al.*, 2011). The supplemented zinc will reach the containing zinc neurons to probably reinforce the role of the endogenous zinc. Zinc supplementation was also reported to prevent stress effects and stress-induced decrease in iron levels (Li *et al.*, 2012). The data of the present investigations confirm low levels of endogenous cerebral iron after acute immobilization stress. This could be due to a changed systemic and brain iron homeostasis following the application of psychological stress (Ma *et al.*, 2008; Wang *et al.*, 2008) occurring via regulation of iron transport (Moss and Morgan, 2004). The role of transferrin seemed to be of high importance since stress induced reduction in ferritin and the subsequent reduction in iron storage and utilization (Huang *et al.*, 2012).

Our findings and the mentioned reports support the fact that a reduction in endogenous level zinc disturbs iron homeostasis in brain, which seems to be highly sensitive to zinc levels. The reduction in serum zinc concentration has also been reported following stress which is concomitant with a decrease in BDNF expression in hippocampus (Cieslic *et al.*, 2011). Indeed, zinc treatment has been shown to induce cortical BDNF gene expression (Nowak *et al.*, 2004). The mechanism, by which the reduction of Zn influences that of BDNF expression, passes probably via the GABAergic system. GABA is reported to diminish the glutamate-stimulated expression of BDNF transcripts and inhibited the stimulatory effect of glutamate on BDNF peptide content (Marmigere *et al.*, 2003). The alteration of both GABA_A receptors and the two heavy metals in substantia nigra is of high interest, as it is a component of the basal ganglia, known to be involved principally in motor control. Furthermore, substantia nigra receives GABA projections from globus pallidus (Erlj *et al.*, 2012). Any alteration in this circuit impaired locomotor behavior and locomotor activity. Interestingly, the impairment of these functions occurs also following stressful situations (Tamburella *et al.*, 2012). Thus, the alterations observed in substantia nigra in this study, could explain these stress-induced motor dysfunctions and suggest that modulatory function of GABA_A receptor by heavy metals could be important for basal ganglia functions. The effect of stress seemed to be specific, as the other parts of basal ganglia did not show any difference in TBPS binding and Zn and Fe concentrations (data not shown).

Taken together, these results support the alteration of the modulatory function occurring at the GABA_A receptor level after a single session of immobilization stress. The concentration of the heavy metals investigated is too low to efficiently modulate the

GABA_A receptor function. This could explain the higher densities of GABA_A receptors observed after acute stress. Finally, these investigations pointed the potential role played by essential heavy metals in plasticity processes related to the stress response and in the regulation of the homeostasis.

5. CONCLUSION

Immobilization stress induces up regulation of GABA_A receptors which is concomitant to down regulation of endogenous zinc, a potent modulator of this receptor as well as endogenous concentrations of iron in brain structures sensitive to stress. Whether the alteration of the two heavy metals concentration could be considered as a consequence or the cause of GABA_A receptor dysfunction, other investigations are needed to elucidate these findings.

6. REFERENCES

- Atack, J.R., Y. Ohashi and R.M. McKernan, 2007. Characterization of [³⁵S]t-butylbicyclophosphorothionate ([³⁵S]TBPS) binding to GABA_A receptors in postmortem human brain. *Br. J. Pharmacol.*, 150: 1066-1074. DOI: 10.1038/sj.bjp.0707186
- Barbaccia, M. L., G. Roscetti, M. Trabucchi, M.C. Mostallino and A. Concas, 1996. Time-dependent changes in rat brain neuroactive steroid concentrations and GABA_A receptor function after acute stress. *Neuroendocrinology*, 63: 166-172. DOI: 10.1159/000126953
- Bitran, D., R.J. Hilvers and C.K. Kellog, 1991. Anxiolytic effects of 3 α -hydroxy-5 α [β]-pregnan-20-one: endogenous metabolites of progesterone that are active at the GABA_A receptor. *Brain Res.*, 561: 157-161. DOI: 10.1016/0006-8993(91)90761-J
- Charrier, C., F. Chigr, C. Tardivel, S. Mahaut and A. Jean *et al.*, 2006. BDNF regulation in the rat dorsal vagal complex during stress-induced anorexia. *Brain Res.*, 1107: 52-57. DOI: 10.1016/j.brainres.2006.05.099
- Chen, L. and W.H. Yung, 2006. Zinc modulates GABAergic neurotransmission in rat globus pallidus. *Neurosci. Lett.*, 409: 163-167. DOI: 10.1016/j.neulet.2006.09.033
- Chigr, F., S. Ba M'hamed and M. Najimi, 2001a. Allosteric modulation of GABA_A receptor by somatostatin is altered under stress in rat brainstem. *Folia Biologica.*, 47: 196-199. PMID: 11768776

- Chigr, F., S. Ba M'Hamed and M. Najimi, 2002. Modulation of [³⁵S]-tert-butylbicyclophosphorothionate binding by somatostatin in rat hypothalamus. *Clin. Exp. Pharmacol. Physiol.*, 29: 291-298. DOI: 10.1046/j.1440-1681.2002.03645.x
- Chigr, F., S. BaM'Hamed and M. Najimi, 2001b. Allosteric modulation of the gabaa receptor in rat hypothalamus by somatostatin is altered by stress. *Clin. Exp. Pharmacol. Physiol.*, 28: 329-330. DOI: 10.1046/j.1440-1681.2001.03447.x
- Cieslic, K., M. Sowa-Kucma, G. Ossowska, B. Legutko and M. Wolak *et al.*, 2011. Chronic unpredictable stress-induced reduction in the hippocampal brain-derived neurotrophic factor (BDNF) gene expression is antagonized by zinc treatment. *Pharmacol. Rep.*, 63: 537-543. PMID: 21602609
- Concas, A., E. Sanna, T. Cuccheddu and M. Paola, 1993. Carbon dioxide inhalation, stress and anxiogenic drugs reduce the function of GABA_A receptor complex in the rat brain. *Prog. Neuropsychopharmacol. Biol. Psychiatry*, 17: 651-661. DOI: 10.1016/0278-5846(93)90012-H
- Concas, A., M.C. Mostallino, P. Porcu, P. Follesa and M.L. Barbaccia *et al.*, 1998. Role of brain allopregnanolone in the plasticity of γ -aminobutyric acid type A receptor in rat brain during pregnancy and after delivery. *Proc. Natl. Acad. Sci.*, 95: 13284-13289. DOI: 10.1073/pnas.95.22.13284
- Cope, E.C., D.R. Morris, A.G. Scrimgeour, J.W. VanLandingham and C.W. Levenson, 2011. Zinc supplementation provides behavioral resiliency in a rat model of traumatic brain injury. *Physiol. Behav.*, 104: 942-947. DOI: 10.1016/j.physbeh.2011.06.007
- Draguhn, A., T.A. Verdorn, M. Ewert, P.H. Seeburg and B. Sakmann, 1990. Functional and molecular distinction between recombinant rat GABA_A receptor subtypes by Zn²⁺. *Neuron*, 5: 781-788. DOI: 10.1016/0896-6273(90)90337-F
- Drugan, R.C., S.M. Paul and J.N. Crawley, 1993. Decreased forebrain [³⁵S]TBPS binding and increased [³H]muscimol binding in rats that do not develop stress-induced behavioral depression. *Brain Res.*, 631: 270-276. DOI: 10.1016/0006-8993(93)91545-4
- Erlj, D., J. Acosta-Garcia, M. Rojas-Marquez, B. Gonzalez-Hernandez and E. Escartin-Perez, 2012. Dopamine D4 receptor stimulation in GABAergic projections of the globus pallidus to the reticular thalamic nucleus and the substantia nigra reticulata of the rat decreases locomotor activity. *Neuropharmacol.*, 62: 1111-1118. DOI: 10.1016/j.neuropharm.2011.11.001
- Farrant, M. and Z. Nusser, 2005. Variations on an inhibitory theme: Phasic and tonic activation of GABA_A receptors. *Nat. Rev. Neurosci.*, 6: 215-29. DOI: 10.1038/nrn1625
- Foddi, M.C., M. Cinquanta and T. Mennini, 1997. Effect of psychological stress on [³⁵S]TBPS binding in rat brain. *Pharmacol. Biochem. Behav.*, 58: 373-377. DOI: 10.1016/S0091-3057(97)00253-0
- Frederickson, C.J. and A.I. Bush, 2001. Synaptically released zinc: physiological functions and pathological effects. *Biometals*, 14: 353-366. DOI: 10.1023/A:1012934207456
- Frederickson, C.J., 1989. Neurobiology of zinc and zinc-containing neurons. *Int. Rev. Neurobiol.*, 31: 145-238. DOI: 10.1016/S0074-7742(08)60279-2
- Fritschy, J.M., and H. Mohler, 1995. GABA_A-receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits. *J. Comp. Neurol.*, 359: 154-194. DOI: 10.1002/cne.903590111
- Gunn, B.G., A.R. Brown, J.J. Lambert and D. Belelli, 2011. Neurosteroids and GABA_A receptor interactions: A focus on stress. *Front Neurosci.*, 5: 131-131. DOI: 10.3389/fnins.2011.00131
- Hosie, A.M., E.L. Dunne, R.J. Harvey and T. Smart, 2003. Zinc-mediated inhibition of GABA_A receptors: discrete binding sites underlie subtype specificity. *Nat. Neurosci.*, 6: 362-369. DOI: 10.1038/nn1030
- Huang, K., H. Li, H. Shen and M. Li, 2012. Psychological stress expands low molecular weight iron pool in cerebral cortex, hippocampus, and striatum of rats. *Biol. Trace. Elem. Res.*, 146: 79-85. PMID: 21993966
- Krishek, B.J., S.J. Moss and T.G. Smart, 1998. Interaction of H⁺ and Zn²⁺ on recombinant and native rat neuronal GABA_A receptors. *J. Physiol.*, 507: 639-652. DOI: 10.1111/j.1469-7793.1998.639bs.x
- Kvetnansky, R. and L. Mikulaj, 1970. Adrenal and urinary catecholamines in rats during adaptation to repeated immobilization stress. *Endocrinology*, 87: 738-743. PMID: 5453288
- Li, Y., Y. Zheng, J. Qian, X. Chen and Z. Shen *et al.*, 2012. Preventive effects of zinc against psychological stress-induced iron dyshomeostasis, erythropoiesis inhibition, and oxidative stress status in rats. *Biol. Trace. Elem. Res.*, 147: 285-291. PMID: 22274754
- Losada, M.E.O. 1989. Acute stress and GABAergic function in the rat brain. *Br. J. Pharmacol.*, 96: 507-512.

- Ma, L., W. Wang, M. Zhao and M. Li, 2008. Foot-shock stress-induced regional iron accumulation and altered iron homeostatic mechanisms in rat brain. *Biol. Trace. Elem. Res.*, 126: 204-213. PMID: 18709494
- Marmigere, F., F. Rage and L. Tapia-Arancibia, 2003. GABA–glutamate interaction in the control of BDNF expression in hypothalamic neurons. *Neurochem Int.*, 42: 353-358. DOI: 10.1016/S0197-0186(02)00100-6
- Miller, J.A., 1991. The calibration of ³⁵S or ³²P with ¹⁴C-labeled brain paste or ¹⁴C-plastic standards for quantitative autoradiography using LKB Ultrofilm or Amersham Hyperfilm. *Neurosci. Lett.*, 121: 211-214. DOI: 10.1016/0304-3940(91)90687-O
- Mody, I., 2012. Plasticity of GABA_A Receptors Relevant to Neurosteroid Action. In: Jasper's Basic Mechanisms of the Epilepsies. Noebels, J.L. M. Avol, M.A. Rogawski, R.W. Olsen and A.V. Delgado-Escuta, (Eds.). Bethesda (MD): National Center for Biotechnology Information (US), pp: 1-9.
- Moss, T. and E.H. Morgan, 2004. The metabolism of neuronal iron and its pathogenic role in neurological disease: Review. *Ann. N.Y. Acad. Sci.*, 1012: 14-26. DOI: 10.1196/annals.1306.002
- Nowak, G., B. Legutko, B. Szewczyk, M. Papp and M. Sanak *et al.*, 2004. Zinc treatment induces cortical brain-derived neurotrophic factor gene expression. *Eur. J. Pharmacol.*, 492: 57-59. DOI: 10.1016/j.ejphar.2004.03.038
- Rahman, M., 2007. Effects of Neuroactive Steroids on the Recombinant GABAA Receptor in *Xenopus* Oocyte. 1st Edn., Umea University, Umea, ISBN-10: 9789172642737, pp: 72.
- Ruiz, A., M.C. Walker, R. Fabian-Fine and D.M. Kullmann, 2004. Endogenous zinc inhibits GABA_A receptors in a hippocampal pathway. *J. Neurophysiol.*, 91: 1091-1096. DOI: 10.1152/jn.00755.2003
- Saxena, N.C., and R.L. Macdonald, 1996. Properties of putative cerebellar gamma-aminobutyric acid A receptor isoforms. *Mol. Pharmacol.*, 49: 567-579. PMID: 8643098
- Serra, M., M.G. Pisu, M. Littera, G. Papi and E. Sanna, 2000. Social isolation-induced decreases in both the abundance of neuroactive steroids and GABAA receptor function in rat brain. *J. Neurochem.*, 75: 732-740. DOI: 10.1046/j.1471-4159.2000.0750732.x
- Smart, T.G., A.M. Hosie and P.S. Miller, 2004. Zn²⁺ ions: modulators of excitatory and inhibitory synaptic activity. *Neuroscientist*, 10: 432-42. DOI: 10.1177/1073858404263463
- Smart, T.G., S.J. Moss, X. Xie and R.L. Haganir, 1991. GABA_A receptors are differentially sensitive to zinc: dependence on subunit composition. *Br. J. Pharmacol.*, 103: 1837-1839. DOI: 10.1111/j.1476-5381.1991.tb12337.x
- Smith, S. S., H. Shen, Q.H. Gong and X. Zhou, 2007. Neurosteroid regulation of GABAA receptors: Focus on the $\alpha 4$ and δ subunits. *Pharmacol. Ther.*, 116: 58-76. DOI: 10.1016/j.pharmthera.2007.03.008
- Tamburella, A., G.M. Leggio, V. Micale, A. Navarria and C. Bucolo, 2012. Behavioural and neurochemical changes induced by stress-related conditions are counteracted by the neurokinin-2 receptor antagonist saredutant. *Int. J. Neuropsychopharmacol.*, 14: 1-11. PMID: 22695046
- Wang, L., W. Wang, M. Zhao M and L. Ma, 2008. Psychological stress induces dysregulation of iron metabolism in rat brain. *Neuroscience*, 155: 24-30. DOI: 10.1016/j.neuroscience.2008.03.091
- White, G. and D. Gurley, 1995. Benzodiazepine site inverse agonists can selectively inhibit subtypes of the GABAA receptor. *Neuroreport*, 6: 1313-1316. PMID: 7669993