

Original Research Paper

Efficiency of *Carica papaya* L. Seeds in Anticoccidial Control for Cockerel of Lohmann Brown Strain

¹Komlan Mawouli Wampah, ²Kosi Mawuéna Novidzro, ¹Namponi Dianam and ³Abalo Essosimna Kulo

¹Centre d'Excellence Régional sur les Sciences Aviaires, École Supérieure d'Agronomie, Université de Lomé, Lomé, Togo

²Département de Chimie, Faculté Des Sciences, Université de Lomé, Lomé, Togo

³Département des Sciences Animale et Vétérinaire, École Supérieure d'Agronomie, Université de Lomé, Lomé, Togo

Article history

Received: 13-09-2023

Revised: 02-11-2023

Accepted: 06-11-2023

Corresponding Author:
Kosi Mawuéna Novidzro
Département de Chimie,
Faculté Des Sciences,
Université de Lomé,
Lomé, Togo
Email: donnenovi@yahoo.fr

Abstract: To combat avian coccidiosis, chemical products are widely used in poultry farming. However, chemotherapy has inconveniences due to the bioresistance of *Coccidia* and the presence of undesirable residues in consumer products. The aim of this study is to investigate the toxicity of papaya seeds as a bio-insecticide against *Coccidia* in Lohmann brown chickens in order to improve their zootechnical parameters. Experimentally, 260 cockerels were divided into five groups among which three experimental groups, marked EX1-3, were treated per month with 5% of the papaya seed powder incorporated in the cockerel feed for two, three, and four consecutive days, "respectively". The 4th group T- which received no treatment, served as a negative control group; while the 5th group T+ was treated once with Amprolium 20%, introduced in the drinking water of the cockerels, served as a positive control group. The one-way ANOVA test was applied to discriminate the mean values calculated statistically, at the 5% probability threshold. The results of laparoscopic analyses, after 15-23 weeks of age, revealed a reduction in the average number of eggs per gram with rates of 66.77; 58.15, and 43.43%, respectively in groups EX1-3. Phytochemical tests carried out on the hydroethanolic extracts (50-50%: v/v) confirmed the presence of biomolecules such as alkaloids, tannins, flavonoids, terpenoids, coumarins, reducing compounds, and proteins in the seeds. The anticoccidial activity revealed in this study would therefore be justified by the presence of the suspected biomolecules in the seeds by the phytochemical tests. Monthly treatment of two consecutive days with 5% of papaya seed powder incorporated in the feed is effective in the fight against cockerel *Coccidia*. This study shows that papaya seed is an interesting natural alternative to synthetic chemical products in the fight against coccidiosis.

Keywords: Poltry, Coccidiosis, Chemotherapy, Papaya Seeds, Avian Safety

Introduction

The consumption of poultry meat and eggs in Sub-Saharan Africa is forecast to increase by up to 200% between 2010 and 2020 (Heise *et al.*, 2015). The relentless need for animal protein in connection with the exacerbated population explosion has led to an intensification of the breeding of short-cycle species; hence the growth of poultry farms around large cities, especially in Africa. Thus, each year, more than 50 billion broilers are raised, representing more than a third of the protein feed for human consumption (Quiroz-Castañeda and Dantán-González, 2015).

Unfortunately, poultry farming is confronted with several zootechnical, economic, health, and pathological constraints (Fall, 2010).

Avian coccidiosis is one of the most frequent and costly diseases for poultry farmers (Dakpogan and Salifou, 2013), causing annual losses that can exceed US\$2.4 billion (Quiroz-Castañeda and Dantán-González, 2015). This parasitosis, caused by intestinal protozoa of the genus *Eimeria* (Gilbert *et al.*, 2011) called *Coccidia* is the source of the decline in the zootechnical performance of parasitized birds (Kouakou *et al.*, 2010). The parasites damage the intestines of the birds, causing clinical symptoms such as intestinal lesions, hemorrhage, and

bloody stools resulting in the death of the birds (Muthamilselvan *et al.*, 2016). Mortality rates can be as high as 80-100% of total bird number (Buldgen *et al.*, 1996). In Africa, the conditions for the development of Coccidial oocysts are most favourable south of the Sahara. The most commonly used conventional approach to coccidiosis prevention for chickens is the incorporation of synthetic coccidiostats into their diet during the first weeks of life (Barbour *et al.*, 2015). However, chemoprevention continues to show its shortcomings, due to the high cost of chemicals, stock-outs, and bioresistance developed by the *Coccidia* themselves (Bichet *et al.*, 2003). As a result, undesirable residues are found in consumer products of avian origin (Chapman *et al.*, 2010), thus constituting a significant public health problem. As an alternative to chemotherapy, vaccines have been developed for most live species, allowing chickens to gradually build up robust immunity (Williams, 1994). About nine species of *Coccidia* are distinguished in chickens (Joyner and Long, 2008). However, simultaneous infection in the environment with several species of *Eimeria* is very common. Unfortunately, the same vaccine is not effective against all species of *Coccidia*. Therefore, there is a risk of introducing into the poultry farm environment the *Eimeria* species present in living vaccines. Consequently, the search for new, more natural, less expensive methods able to reduce infection and strengthen the immune system of the host is necessary (Kouakou *et al.*, 2010).

In this line of thought, medicinal plants that possess biomolecules with high antiparasitic potential (Bauri *et al.*, 2015) offer very good prospects. Indeed, extracts of some plants such as *Aloe vera*, *Azadirachta indica*, *Artemisia annua*, and *Pinus radiata* have been shown to reduce oocyst secretion and or intestinal lesions (Yim *et al.*, 2011). Traditional pharmacopoeia is now emerging as a record model for *Coccidia* control (Muthamilselvan *et al.*, 2016; Dakpogan *et al.*, 2018).

The papaya plant "*Carica papaya*" is a perennial medicinal plant that is easy to grow and adapts to various soils (Agboola *et al.*, 2018). Many studies have shown that the plant exhibits various antimicrobial properties (Abd Elgadir *et al.*, 2014) including anthelmintic (Soedji *et al.*, 2017), anticoccidial (Hema *et al.*, 2015; Dakpogan *et al.*, 2018) and immune system enhancing (Muazu and Aliyu-Paiko, 2020). Most studies on the use of *Carica papaya* seeds in poultry farming have focused mainly on their anthelmintic effects. Therefore, the originality of our study lies in the use of papaya seed as an alternative to synthetic chemical insecticides against avian coccidiosis.

In this context, the current study was undertaken to evaluate the anticoccidial properties of papaya seeds in cockerels, with a view to improving zootechnical

parameters and then to mention the phytoconstituents that would be responsible for the seeds' anticoccidial activities.

Materials and Methods

Study Frameworks and Period

All the works related to the current study were carried out in Togo, from September 2018 up to August 2019, precisely on a breeding site close to Badja village, called "Ayedole" farm. This site is characterized by a humid tropical climate with annual ambient temperatures ranging from 25°-32°C and an average humidity level of 75%. The poultry house is an open building with variable temperature and humidity.

The coproscopic analyses were carried out at the Centre d'Excellence Regional sur les Sciences Aviaires (CERSA) of the Université de Lomé.

Phytochemical tests were carried out at the "Laboratoire de Génie des Procédés et des Ressources Naturelles (LAGEPREN) of the Université de Lomé".

Plant and Animal Materials

Papaya seeds, variety 'solo' (Fig. 1), were used as plant material to combat the *Coccidial* effects of cockerels.

The seeds, purchased from papaya sellers in Lomé-Togo, were dried under air conditioning at 20°C in the CERSA laboratory, and then crushed in a mill. The powder obtained was stored, protected from sunlight, for later use.

To ensure that the total number of birds studied was representative and in the interest of obtaining statistically significant results, the animal material consisted of two hundred and sixty (260) Lohmann brown *Cockerels*, which had been reared together for eight weeks. During this period, the birds were vaccinated against Newcastle disease, infectious bronchitis, Gumboro disease, and smallpox and treated with Amprolium 20%.




Fig. 1: Photos of *Carica papaya* L. "Solo" (A = Fruiting plant and B = Fruit and seeds contained in fruit)

Experimental Methodologies

In accordance with current CERSA regulations, all ethical rules concerning the use of animals in experiments have been respected.

Disinfection of the Breeding Site and Groups' Constitution

Before installing the 260 cockerels on the site, the building reserved for the bioassay was washed, disinfected, and subjected to a sanitary vacuum for a fortnight, and then disinfection was repeated two days before the cockerels were installed. In order to protect the welfare of the chickens, regulations limit the number of chickens per m² to 17 cockerels in the standard poultry house.

After three weeks of acclimatization, the 260 cockerels were initially weighed before being randomly divided into five groups, each containing 52 cockerels. The five groups were numbered as follows:

- T-: Group that received no anticoccidial treatment throughout the experiment, served as a negative control group
- T+: The group that received a monthly anticoccidial treatment with Amprolium 20% for four consecutive days in the drinking water, served as a positive control group
- EX1: The experimental group that received a monthly treatment of 5% papaya seeds on two consecutive days
- EX2: The experimental group that received a monthly treatment of 5% papaya seed for three consecutive days
- EX3: The experimental group that received a monthly treatment of 5% papaya seed on four consecutive days

The cockerels were grown on floor bedding at a density of 17 cockerels/m², from start-up through to the growth phase. The experiments started on 23rd September 2018 and ended on 25th February 2019, for a duration of 20 weeks.

Throughout the trials, water and feed were provided *Ad libitum* to the cockerels.

Coproscopic Examination

This analysis is based on the Mc master method of *Eimeria* egg counting. According to the experimental protocol, 2 g of feces from each replication was taken with a spatula and crushed in a mortar with a pestle in 60 mL of a saturated solution of NaCl (50%). The mixture was filtered through a sieve to remove large particles. Using a Pasteur pipette, both cells of the Mc master slide were filled with the filtrate while avoiding the formation of air bubbles. After 5 min of waiting, the Number of Eggs Per Gram (NEPG: OPG) was determined

by observing the oocysts under a light microscope. The Reduction Rate (RR) of the NEPG was calculated using the following formula:

$$RR = \frac{(NEPG \text{ before treatment} - NEPG \text{ after treatment})}{t \times 100\%} \times NEPG \text{ before treatment}$$

Oocyst Extraction and Coccidia Identification

Hemorrhagic areas were checked for along the digestive tract by means of autopsy. Extraction of the intestinal contents allowed us to identify the species of *Coccidia* that affected the subjects. For this purpose, two subjects per replication were sacrificed and their digestive tracts were isolated. For each digestive tract, the intestine was then isolated and sectioned into its four different constituent portions, namely: Duodenum, jejunum, ileum, and caecum. Each portion was incised and a smear was taken and observed under the microscope. The remaining intestinal contents were blended in tap water and filtered through a sieve to remove large particles. The filtrate was left to settle for 2 h and the supernatant was removed. The pellet was centrifuged at 3000 rpm for 7 min. The supernatant was again removed and the pellet was suspended in salt water (50%) and centrifuged at 2000 rpm. The supernatant finally recovered was diluted in tap water and the mixture was centrifuged at 3000 rpm. A few drops of the pellet containing the oocysts were mounted between the slide and the coverslip and observed under a microscope at a magnification of 60. The remaining pellet was allowed to sporulate at room temperature for three days and observed again under the microscope.

Assessment of Zootechnical Parameters

Birds and feed were weighed with a CAMRY/EK 3250 scale to determine the following zootechnical parameters.

Daily Feed Consumption (DFC): This was calculated using the following formula:

$$DFC = \frac{\text{Amount of feed distributed (g)} - \text{Amount of feed remaining (g)}}{\text{Length of time} \times \text{number of subjects}}$$

Consumption Index (CI): This was determined using the following formula:

$$CI = \frac{\text{Amount of food consumed over time (g)}}{\text{Gain in weight during the time (g)}}$$

Cockerel weight progression: The weekly weight measurements of the birds were used to determine the Live Weight (LW) and to calculate the Average Daily Gain (ADG) using the following formulae:

$$LW = \frac{\text{Total weight of subjects (g)}}{\text{Number of subjects}}$$

$$ADG = \frac{\text{weight (g) over time}}{\text{length of time (day)}}$$

Phytochemical Screening of Papaya Seeds

Carica papaya L. seed powder was extracted with a hydroethanolic mixture (50-50%: v/v) using the soxhlet extraction method. The total extracts collected were used for qualitative and quantitative phytochemical analysis for the identification of secondary metabolites, mainly present in the seeds.

Qualitative Phytochemical Analyzes

Dragendorff, Wagner, and Mayer reagents were used for the search for alkaloids (Eke *et al.*, 2014; Yadav *et al.*, 2014). The foam test was used in the search for saponosides (Eke *et al.*, 2014). Tannins were detected by using the 1% ferric chloride, 10% lead acetate, and ammoniacal copper sulphate reagent tests. Flavonoids were tested with the 1% sodium hydroxide reagent. The triterpenes were detected with sulphuric acid and the coumarins with sodium hydroxide solution. The identification of proteins was performed by the xanthoprotein test and that of reducing compounds by the Fehling test (Eke *et al.*, 2014; Yadav *et al.*, 2014).

Quantitative Phytochemical Assays

Quantification of Total Phenols by Calibration with Gallic Acid

In each test tube were introduced 200 μ L of the sample solution, either gallic acid or extract (1 mg/mL), and 500 μ L of Folin Ciocalteu's reagent (diluted to 1/2 in distilled water). After a reaction time of 5 min, 500 μ L of sodium carbonate (20 g/L) was added to the previous mixture and then supplemented with 4 mL of distilled water. After homogenization, the solutions thus prepared were incubated for 30 min at laboratory temperature and in the dark. The absorbance of each solution was read at 760 nm with a UV-visible/METASH spectrophotometer (UV-5200 PC) against a blank. The calibration curve (Fig. 2) was established with a gallic acid solution (200 mg/L). The result was expressed as mg Gallic Acid Equivalent per gram (GAE/g) of dry extract.

Assay of Proanthocyanidins or Condensed Tannins

In a test tube, 0.2 mL of an Ammoniacal Iron Sulphate solution ($\text{NH}_4\text{Fe}(\text{SO}_4)_2$: 20 g/L), 7 mL of a mixture of Bu-OH/H-Cl (95-5%: v/v), and 0.2 mL of an aqueous extract solution (30 mg/mL) were successively introduced. After

incubation of the mixture in a water bath at a temperature of around 95°C for 40 min, a pink or red coloration was obtained. The proanthocyanidin concentration of an unknown sample was estimated by reading the Optical Density (OD) of the solution at 550 nm with a UV-visible/METASH spectrophotometer (UV-5200 PC). The Content of proanthocyanidins (C_{Pro}) in a sample, expressed as mg Catechin Equivalent (CE)/g of extract, was calculated using the relationship established by Aksamit-Stachurska *et al.* (2008) as follows:

$$C_{Pro} = \frac{DO}{0.280}$$

With: $OD = 0.280$ which is equivalent to 1% Catechin Equivalent (CE) and using catechin as the standard molecule.

Assessment of the Antioxidant Activity of Total Extracts

A volume of 1 mL of total extract (30 mg/mL) was added to 9 mL of phosphomolybdate reagent. The previous solution was heated in a water bath at 95°C for 90 min and then allowed to cool to room temperature. Optical densities were measured at 695 nm with a UV-visible/METASH spectrophotometer (UV-5200 PC). The calibration curve (Fig. 3) was established with ascorbic acid solution and the result was expressed as mg Ascorbic Acid Equivalent (EAA) per gram of dry extract.




Fig. 2: Gallic acid calibration curve




Fig. 3: Ascorbic acid calibration curve

Statistical Analysis of Data

The data collected on animal weight, feed consumption, and NEPG were statistically processed using Graph Pad Prism 5 software. The one-way ANOVA test was applied to discriminate the mean values calculated statistically, at the 5% probability threshold. The results, presented as means plus or minus Standard Errors of Means (SEM), were compared with each other by applying the TUKEY test.

Results

Zootechnical Parameters of the Cockerels

Feed Consumption

The Daily Feed Consumption (DFC) for cockerels is shown in Fig. 4. No significant differences are observed for the different groups treated. However, the values observed in groups T- and EX3 appear to be slightly higher than those of the other groups.

Cockerel Weight Progression

Live Weight

Figure 5 shows the Live Weight (LW) of the cockerels. The LW values of all of the groups presented a similar evolution between 12th and 17th weeks. However, from the 18th week onwards, group EX1 had the highest growth rate towards the end of the experiment, i.e., around week 22.

Average Daily Gain (ADG)

Figure 6 shows the ADG values of the cockerels. The value obtained by the group EX1 is significantly higher than that of the group T-. However, this value was not significantly different from those of the other groups (EX2, EX3, and T+).

Index of Consumption (IC)

The data displayed in Fig. 7 clearly indicate the IC of the cockerels. The cumulative IC value of the group T- during the trial is the highest, but between groups T- and EX3, the difference is not so significant.

Parasitic Load Evolution

Number of Eggs Per Gram (NEPG) Variation

As a result of the papaya seed treatment, groups EX1 and EX2 showed remarkably lower NEPG than that of group T-. The parasite load of group EX3, although appearing to be higher than that of group T+, remained significantly lower than that of group T- (Fig. 8).

NEPG Reduction Rate

The data presented in Table 1 shows that there was a reduction of NEPG between the cockerels before and after their treatments. The different treatments with papaya seed powder resulted in an average reduction NEPG equal to: 66.77, 58.15 and 43.43%, respectively for the groups EX1, EX2, EX3, and 51.93% for the T+ Generally, the

best rate of reduction in NEPG was obtained in group EX1 after two days of treatment.




Fig. 4: Cockerel daily feed consumption




Fig. 5: Evolution of LW of cockerels as a function of age




Fig. 6: Cockerel average daily gain




Fig. 7: Consumption index of the cockerels

Table 1: NEPG variation before and after treatment

Age (weeks)	Deadline and discount rate		EX1	EX2	EX3	T-	T+
S15	<i>First treatment</i>						
	NEPG before treatment	J0	425	2025	1500	21750	1250
		J2	175				
	NEPG after treatment	J3		900			
		J4			363	26600	625
	Corresponding reduction rate (%)		58.82	55.55	75.80	-22.29	50.00
S19	<i>Second treatment</i>						
	NEPG before treatment	J0	1950		4700	5450	14250
		J2	775				
	NEPG after treatment	J3		1525			
		J4			3500	49375	8000
	Corresponding reduction rate (%)		60.25	67.55	35.77	-246.49	28.88
S23	<i>Third treatment</i>						
	NEPG before treatment	J0	400	925	3475	8100	325
		J2	75				
	NEPG after treatment	J3		450			
		J4			2825	4025	75
	Corresponding reduction rate (%)		81.25	51.35	18.700	50.31	76.92
S15-S23	Average reduction rates (%)		66.77	58.15	43.430	-72.82	51.93




Fig. 8: Variation in NEPG in treated groups according to the age of the cockerels

Autopsy and Intestinal Smear Results

The autopsy showed hemorrhagic areas of the intestinal mucosa throughout the digestive tract, indicative of the presence of *Coccidia* (Fig. 9).

The hemorrhagic areas were more intense in group T- compared to the treated groups. This trend was confirmed by the intestinal mucosal smear, which also showed non-spore-forming oocysts in groups T- and EX3 (Fig. 10). In addition, group T- subjects had more bloody stools than group EX3 subjects.

Extraction of intestinal contents indicated the presence of *Eimeria* species such as *Eimeria acervulina*, *Eimeria maxima*, *Eimeria necatrix*, *Eimeria brunetti*, and *Eimeria tenella* in the cockerel gut (Fig. 11) and the four sites of localization in the intestinal tract were: Duodenum, jejunum, ileum, and caecum (Table 2).

Results of Phytochemical Analyses

Cases of Qualitative Phytochemical Analyses

Qualitative tests reveal in the hydroethanolic extract (50-50%) of the papaya seeds used in this study the presence of the secondary metabolites presented in Table 3.

Cases of Quantitative Phytochemical Analysis

The contents of proanthocyanidins (or condensed tannins) and total phenols and antioxidant activity of the hydroethanolic extracts of papaya seeds are presented in Table 4.




Fig. 9: Autopsy results




Fig. 10: Bowel smear results




Fig. 11: Extracted pictures of *Coccidia* species

Table 2: *Coccidia* species identified in the digestive tracts of cockerels

N°	<i>Coccidia</i> species	Intestinal location
1	<i>Eimeria acervulina</i>	Duodenum
2	<i>Eimeria maxima</i>	Jejunum
3	<i>Eimeria necatrix</i>	Jejunum
4	<i>Eimeria brunetti</i>	Ileon
5	<i>Eimeria tenella</i>	Caecum

Table 3: Results of qualitative phytochemical analyses

Investigated phytochemical compounds	Analytical methods	Achieved results
Reducing compounds	Fehling reagents	+
Alkaloids	Mayer reagent	+
	Dragendorff reagent	+
	Wagner reagent	+
Tannins	Ferric chloride 1%	+
	Lead acetate 10%	+
	Ammoniacal copper sulphate	+
Flavonoids	Sodium hydroxide (1%)	+
Coumarins	Sodium hydroxide (10%)	+
Triterpenes and phytosterols	Sulphuric acid (1 M)	+
Saponosides	Vigorous shaking of the solution	-
Proteins	Nitric acid + Ammonia	+

Legend: + = positive test, while - = negative test

Table 4: Results of quantitative phytochemical analyses of hydroethanolic extracts of papaya seeds

Kinds of quantitative analysis	Measured values
Prothocyanidin content (mg CE/g)	12.05
Total phenol content (µg GAE/g)	45150.01
Antioxidant capacity (mg AAE/g)	42.33

Discussion

The feed consumption of cockerels of groups EX3 and T- was higher than that of the other groups. It could be explained by the higher parasite load of oocysts, because

according to Dossou (2008), the feed consumption of an animal depends on several factors, including its LW.

It was noted that *Coccidia* are spoliating gastrointestinal parasites like helminths (Brugère-Picoux and Silim, 1992). Therefore, the low weight gain recorded in group T- (Fig. 6) would be attributable to the impacts caused by these parasites.

In the case of coccidiosis, the effect of parasitism may be to maintain or even exacerbate the hunger, in order to compensate for the deficits in nutritional intake caused by the intestinal lesions. The low weight gain observed in groups T- and EX3 could be explained by the high parasite load, as parasites are qualitative spoliators that attack their host's metabolism and divert essential nutrients such as amino acids, vitamins, and minerals to themselves (Maminiaina, 2017).

According to our results, all the groups treated with *Carica papaya* L. seed powder showed lower CI compared to the group T-. More specifically, groups EX1, EX2, and EX3 showed better feed utilization efficiency compared to group T-. Similar results were also found by Essomba (2003) who, in the course of his investigation, noted that the IC of chickens was higher in *Coccidia*-infested subjects than in non-*Coccidia*-infested subjects. In their approach, (Chrysostome *et al.*, 2010) observed that during the growth phase, cockerels in the tropics have an IC value of around 5.12%.

In the current study, *Carica papaya* seeds induced a decrease in the IC in the cockerels of groups EX1 and EX2, thus showing better use of the feed provided, so the use of *Carica papaya* L. seed powder significantly reduced oocyst excretion in cockerels. Similar results were also obtained by Mpoame *et al.* (2003); Dakpogan *et al.* (2018) in broilers.

In our first treatment, the rate of reduction of NEPG is proportional to the days of treatment with the seeds. However, over time, the effectiveness of the treatment decreases with the days of treatment, thus with the dose of papaya seeds ingested. These results are in contradiction with those of Mpoame *et al.* (2003) who observed an increase in the rate of reduction with increasing doses of the treatment product in broiler chickens infested by *Eimeria tenella*. Perhaps, these authors used aqueous extracts of *Carica papaya* L. seed powder, possibly more concentrated in coccidiostatic active principles against *Eimeria tenella*.

However, in the second treatment (Mpoame *et al.*, 2003) obtained the highest NEPG with the highest dose, which is then in accordance with our results.

The current study showed that the efficacy of seeds decreases over time with only high doses. This hypothesis could be explained either by the indirect effect of anti-nutritional factors contained in the seeds, or by an accumulation of some phytochemical constituents which, beyond a certain quantity, would present harmful effects on the cockerels.

However, the results presented in the current study are in conformity with those obtained by Mpoame and Essomba (2000) in the treatment with aqueous decoction of *Carica papaya* L. seeds, against gastrointestinal parasitosis in naturally local infested chickens. These authors observed a decrease in the rate of reduction of the NEPG with the dose of the treating product.

The efficacy of papaya seeds in the fight against *Coccidia* would vary with the galenic form of use (aqueous extract, ethanolic extract, decoction, or powder); so according to Eke *et al.* (2014), the antibacterial efficacy of *Carica papaya* seeds depends on the extraction solvent.

According to the points of view (Chapman, 2014), the immediate action of *Coccidia* in the intestine is the destruction of enterocytes, accompanied by inflammation, hemorrhages, atrophies of the intestinal villi, abnormal differentiation of epithelial cells, and thickening of the intestine. This leads to a slowing of intestinal transit, increased permeability, and a reduction in the rate of absorption of nutrients. The diversion of nutrients by *coccidia* therefore causes a deficit in nutritional intake for poultry (N'dri, 2009).

The results presented in this study show that by treating the subjects with *Carica papaya* seed powder, the average weight gain as well as the efficiency of dietary use is inversely proportional to the powder dose. These results are consistent with the work done by Bolu *et al.* (2009); Soedji *et al.* (2017). These authors found that poor feed utilization is observed when the dose of seeds is increased due to the accumulation of anti-nutritional factors contained in the seeds. Indeed, some anti-nutritional compounds in plants interact with the nutrients of birds. The disadvantage is that the ingestion of high levels of anti-nutrients leads to a decrease in the bioavailability of minerals (Reddy *et al.*, 1982), with negative consequences for animal health and productivity. For example, in high doses, tannins can damage intestinal cells, interfere with mineral absorption, and even generate a toxic effect (Butler, 1989). Therefore, a high tannin content in bird feed decreases microbial enzyme activity and cellulose digestion (Aletor, 1999). Flavonoids in turn, at high doses, reduce the availability and digestibility of protein and other nutrients such as fiber and starch (Jeroch *et al.*, 1999). Thus, in the EX3 batch with the highest feed conversion ratio, the accumulation of anti-nutritional factors at an intense proportion could be questioned.

Phytochemical screening of *Carica papaya* (Linne) seeds revealed the presence of phytochemical compounds such as Alkaloids, tannins, flavonoids, terpenoids, coumarins, reducing compounds, and proteins (Table 3). These results are comparable to those of (Dada *et al.*, 2016; Loe *et al.*, 2017) who also showed that *Carica papaya* L. seeds contain alkaloids, tannins, flavonoids, steroids, polyphenols, but also saponosides by

phytochemical analysis. However, in this study, saponosides were not found in the hydroethanolic extract of the seeds. However, saponosides absence in hydroethanolic extracts of the current work are in agreement with the work done by Singh *et al.* (2018) on ethanolic and aqueous extracts of *Carica papaya* leaves and roots. The presence or absence of saponosides in *Carica papaya* seed extracts could be either related to the nature of the varieties studied or to the polarity of the extraction solvent. Indeed, the work carried out by Eke *et al.* (2014) thus reported the presence of saponosides in the ethanolic extract while they were absent in the chloroform and benzene extracts of the fruit and seeds.

The anticoccidial activity of *Carica papaya* L. is linked to the phytochemical constituents synthesized in the seeds of this plant. Indeed, among the various secondary metabolites whose existence has been revealed in our extract, phenolic compounds (N'dri, 2009), alkaloids (Edgar and Flanagan, 1979), condensed tannins (Molan *et al.*, 2009), and flavonoids (Muthamilselvan *et al.*, 2016) have been identified as biomolecules endowed with anticoccidial properties. Therefore, the anticoccidial effect of *Carica papaya* L. seeds could then be attributed to the chemical compounds they contain, so according to Kaleem *et al.* (2014), tannins isolated from *Emblca officinalis* induced humoral immune response in chickens against *Coccidia* infection. Conferring some previous work, the anticoccidial action of *Carica papaya* L. seeds can also be explained by the proteolytic action of the papain contained in the seeds (Al-Fifi, 2007; Nghonjuyi *et al.*, 2015).

Conclusion

In fine, the results presented in this study show that the incorporation of *Carica papaya* L. seed into the feed presented toxic effects against avian coccidiosis in cockerels. Consequently, there is a reduction in the Consumption Index (CI) and an increase in the Average Daily Gain (ADG) of chickens, thanks to the significant reduction in *Coccidian* load; hence the good performance of the zootechnical parameters of cockerels. The presence of some biomolecules revealed in the seed, such as alkaloids and phenolic compounds, would be the basis of this anticoccidial activity. Overall, the seed treatment at a dose of 5% for two days achieved the best results.

However, to allow chicken breeders to make more reliable use of the results, it would be appropriate to isolate the active ingredients contained in the seeds in order to test their effectiveness on each species of *Coccidia*; and then determine the mechanism of action of the active principles and finally evaluate the toxicity of the seeds for chickens.

Acknowledgment

The authors would like to thank the authorities of the Centre d'Excellence Régional sur les Sciences Aviaires (CERSA) of Université de Lomé-Togo for providing them with their technical equipment, solvents, and reagents, necessary for carrying out chemical analyses on the seed of *Carica papaya* L.

Funding Information

This project was financially supported by the World Bank for its implementation at CERSA, at Université de Lomé-Togo.

Author's Contributions

Komlan Mawouli Wampah: Carried out the various experiments and acquisition of data, interpreted the results and participated in article writing.

Kosi Mawuéna Novidzro: Coordinated the experiments and interpretations of data and participated in article written critical revision of the article and final approval of the article.

Namponi Dianam: Participated in the experiment's realization and data interpretation.

Abalo Essosimna Kulo: Designed the research plan and organized the study.

Ethics

All procedures adopted in the current study were approved by the scientific ethics committee of the Centre d'Excellence Régional sur les Sciences Aviaires (CERSA) of the université de Lomé authorized the defense of the master's thesis under the number No. 065, publicly supported on 13/09/2019.

References

- Abd Elgadir, M., Salama, M., & Adam, A. (2014). *Carica papaya* as a source of natural medicine and its utilization in selected pharmaceutical applications. *Int J Pharm Sci*, 6(1), 868-871.
<https://www.researchgate.net/publication/287400709>
- Agboola, B. E., Ologhobo, A. D., Adejumo, I. O., & Adeyemo, G. O. (2018). Response of broiler chickens to *Carica papaya* and talinum triangulare leaf meal under normal and subnormal diets. *Annual Research and Review in Biology*, 1-7.
<https://doi.org/10.9734/ARRB/2018/38144>
- Aksamit-Stachurska, A., Korobczak-Sosna, A., Kulma, A., & Szopa, J. (2008). Glycosyltransferase efficiently controls phenylpropanoid pathway. *BMC Biotechnology*, 8, 1-16.
<https://doi.org/10.1186/1472-6750-8-25>

- Aletor, V. A. (1999). Anti-nutritional factors as nature's paradox in food and nutrition securities. <http://196.220.128.81:8080/xmlui/handle/123456789/256>
- Al-Fifi, Z. I. A. (2007). Effect of leaves extract of *Carica papaya*, *vernonia amigdalina* and *azadiratcha indica* on the coccidiosis in free-range chickens. *Asian Journal of Animal Sciences*, 1(1), 26-32.
<https://doi.org/10.3923/ajas.2007.26.32>
- Barbour, E. K., Ayyash, D. B., Iyer, A., Harakeh, S. T. K. A., & Kumosani, T. (2015). A review of approaches targeting the replacement of coccidiostat application in poultry production. *Brazilian Journal of Poultry Science*, 17, 405-418.
<https://doi.org/10.1590/1516-635x1704405-418>
- Bauri, R. K., Tigga, M. N., & Kullu, S. S. (2015). A review on use of medicinal plants to control parasites. *Indian Journal of Natural Products and Resources (IJNPR) [Formerly Natural Product Radiance (NPR)]*, 6(4), 268-277.
<https://doi.org/10.56042/ijnpr.v6i4.8837>
- Bichet, H., Sanaa, M., Dorchies, P. H., & Reperant, J. M. (2003). Mise en évidence de coccidies multi-resistantes chez la poule pondeuse au senegal. *Revue De Médecine Vétérinaire*, 154(6), 439-445.
<https://www.researchgate.net/publication/299063191>
- Bolu, S. A. O., Sola-Ojo, F. E., & Olorunsanya, O. A. (2009). Effect of graded levels of dried pawpaw (*Carica papaya*) seed on the performance, haematology, serum biochemistry. *International Journal of Poultry Science*, 8(9), 905-909.
<https://doi.org/10.3923/ijps.2009.905.909>
- Brugère-Picoux, J., & Silim, A. (1992). Manuel de pathologie aviaire. *Ed. Chaire de Pathologie Médicale du Bétail et des Animaux de Basse-Cour, Ecole Nationale Vétérinaire d'Alfort, Maisons-Alfort (France)*; 381 pages. ISBN-10: 2-87820-001-2.
<https://search.worldcat.org/fr/title/manuel-de-pathologie-aviaire/oclc/29245019>
- Buldgen, A., R. Parent, P. Steyaert & Legrand. D. (1996). Aviculture semi-industrielle en climat subtropical: guide pratique. *Gembloux: Les Presses Agronomiques*, 122.
<https://koha.uac.bj/cgi-bin/koha/opac-detail.pl?biblionumber=28222>
- Butler, L. G. (1989). Effects of condensed tannin on animal nutrition. *In Chemistry and Significance of Condensed Tannins*, 391-402. Boston, MA: Springer US. https://doi.org/10.1007/978-1-4684-7511-1_24
- Chapman, H. D., Jeffers, T. K., & Williams, R. B. (2010). Forty years of monensin for the control of coccidiosis in poultry. *Poultry Science*, 89(9), 1788-1801.
<https://doi.org/10.3382/ps.2010-00931>

- Chapman, H. D. (2014). Milestones in avian coccidiosis research: A review. *Poultry Science*, 93(3):501-11. <https://doi.org/10.3382/ps.2013-03634>
- Chrysostome, C. A. A. M., Houndonougbo, M. F., & Kpomassè, C. C. (2010). Stratégie alimentaire des coquelets en zone tropicale: Ration complète comparée au système cafétéria. *Livestock Research for Rural Development*, 22(11).
- Dada, F. A., Nzewuji, F.O. Esan, A.M. Oyeleye, S.I. & Adegbola. V.B. (2016). Phytochemical and antioxidant analysis of aqueous extracts of unripe pawpaw (*Carica papaya* Linn.) fruit's peel and seed. *International Journal of Research and Reviews in Applied Sciences*, 27 (3): 68-71.
- Dakpogan, H. B., & Salifou, S. (2013). Coccidiosis prevalence and intensity in litter based high stocking density layer rearing system of Benin. *J. Anim. Plant Sci*, 17(2), 2522-2526. <http://www.m.elewa.org/JAPS>
- Dakpogan, H. B., Mensah, S., Attindehou, S., Chysostome, C., Aboh, A., Naciri, M., & Mensah, G. A. (2018). Anticoccidial activity of *Carica papaya* and *vernonia amygdalina* extract. *International Journal of Biological and Chemical Sciences*, 12(5), 2101-2108. <https://doi.org/10.4314/ijbcs.v12i5.12>
- Dossou, A. D. (2008). Effets du tourteau de Neem (*Azadirachta indica*) sur les coccidioses aviaires. *Thèse de Doctorat d'état en Médecine Vétérinaire de l'Université Cheikh Anta Diop*, Dakar, Sénégal. <https://beep.ird.fr/greenstone/collect/eismv/index/assoc/TD08-27.dir/TD08-27.pdf>
- Edgar, S. A., & Flanagan, C. (1979). Efficacy of Stenorol®(Halofuginone): II. Plus roxarsone or bacitracin MD against selected strains of chicken *Eimeria*. *Poultry Science*, 58(6), 1476-1482. <https://doi.org/10.3382/ps.0581476>
- Eke, O. N., Augustine, A. U., & Ibrahim, H. F. (2014). Qualitative analysis of phytochemicals and antibacterial screening of extracts of *Carica papaya* fruits and seeds. *International Journal of Modern Chemistry*, 6(1), 48-56. ISSN-10: 2165-0128.
- Essomba, L. I. (2003). Amélioration des productions avicoles par l'utilisation de la pharmacopée traditionnelle dans la lutte contre la coccidiose aviaire au Cameroun. *Mémoire de Diplôme d'Études Approfondies de Productions Animales de l'Université Cheikh Anta Diop*. Dakar, Sénégal. <https://beep.ird.fr/greenstone/collect/eismv/index/assoc/MEM03-2.dir/MEM03-2.pdf>
- Fall, O. (2010). Effets de l'incorporation du tourteau de neem (*Azadirachta indica* a. juss) a 2% dans l'aliment et dans la litière sur les performances zootechniques et les maladies parasitaires chez la poule pondeuse. *Thèse en Médecine Vétérinaire de l'Université Cheikh Anta Diop*. Dakar, Sénégal. <https://beep.ird.fr/greenstone/collect/eismv/index/assoc/TD10-23.dir/TD10-23.pdf>
- Gilbert, E. R., Cox, C. M., Williams, P. M., McElroy, A. P., Dalloul, R. A., Ray, W. K., ... & Webb Jr, K. E. (2011). *Eimeria* species and genetic background influence the serum protein profile of broilers with coccidiosis. *PLOS One*, 6(1), e14636. <https://doi.org/10.1371/journal.pone.0014636>
- Heise, H., Crisan, A., & Theuvsen, L. (2015). The poultry market in Nigeria: Market structures and potential for investment in the market. *International Food and Agribusiness Management Review*, 18(1030-2016-83098), 197-222. <https://doi.org/10.22004/ag.econ.207011>
- Hema, S., Arun, T., Senthilkumar, B., Senbagam, D., & Sureshkumar, M. (2015). *In vivo* anticoccidial effects of *Azadirachta indica* and *Carica papaya* L. with salinomycin drug as a dietary feed supplement in broiler chicks. *Pak. J. Pharm. Sci*, 28(4), 1409-1415. <https://www.researchgate.net/publication/277566806>
- Jeroch, H., Drochner, W. & Simon. O. (1999). Ernährung landwirtschaftlicher nutztiere. UTB: Stuttgart, Germany. ISBN-10: 9783825281809.
- Joyner, L. & Long. P. (2008). The specific characters of the *Eimeria*, with special reference to the *Coccidia* of the fowl. *Pathologie Aviaire*, 3(3), 145-157. <https://doi.org/10.1080/03079457409353827>
- Kaleem, Q. M., Akhtar, M., Awais, M. M., Saleem, M., Zafar, M., Iqbal, Z., ... & Anwar, M. I. (2014). Studies on emblica officinalis derived tannins for their immunostimulatory and protective activities against coccidiosis in industrial broiler chickens. *The Scientific World Journal*, 2014. <https://doi.org/10.1155/2014/378473>
- Kouakou, S. K., Toure, A., Ouattara, K., & N'Guessan, J. D. (2010). Activité anticoccidienne *in vivo* de l'extrait aqueux des inflorescences de *thonningia sanguinea* (Balanophoraceae) chez la poule pondeuse. *International Journal of Biological and Chemical Sciences*, 4(4). <https://doi.org/10.4314/ijbcs.v4i4.62969>
- Loe, G. E., Yinyang, J., Ebongue, C. O., Makondo, B. V., Ngaba, G. P., Mpondo, E. M., & Dibong, S. D. (2017). Étude de la toxicité aigue et subaigüe de l'extrait au vin des graines de *Carica papaya* Linn. *Journal of Applied Biosciences*, 120, 12077-12085. <https://www.ajol.info/index.php/jab/article/view/168745>
- Maminiaina, O.F. (2017). Expérimentation sur les moyens de lutte contre les maladies parasitaires internes et la maladie de newcastle dans un milieu contrôlé à kianjasoa. *Technical Report*. <https://doi.org/10.13140/RG.2.2.20465.79201>.
- Molan, A. L., Liu, Z., & De, S. (2009). Effect of pine bark (*Pinus radiata*) extracts on sporulation of *Coccidian* oocysts. *Folia Parasitologica (Prague)*, 56(1), 1. <https://doi.org/10.14411/fp.2009.001>

- Mpoame, M., & Essomba, L. I. (2000). Essai de traitement contre des parasitoses gastro-intestinales du poulet avec des décoctions aqueuses de graines de papaye (*Carica papaya*). *Revue D'élevage Et De Médecine Vétérinaire des Pays Tropicaux*, 53(1), 23-25. <https://doi.org/10.19182/remvt.9761>
- Mpoame, M., Teguia, A., & Akoa Etoa, J. M. (2003). Evaluation de l'efficacité des extraits aqueux de graines de papaye (*Carica papaya* L.) dans le traitement de la cocci-diose caecale à *Eimeria tenella* chez le poulet de chair. *Tropicultura*, 21(3), 153-156. <https://www.researchgate.net/publication/45266366>
- Muazu, U., & Aliyu-Paiko, M. (2020). Evaluating the potentials of *Carica papaya* seed as phytobiotic to improve feed efficiency, growth performance and serum biochemical parameters in broiler chickens. *IOSR Journal of Biotechnology and Biochemistry*, 6(1), 8-18. <https://doi.org/10.9790/264X-0601010818>
- Muthamilselvan, T., Kuo, T. F., Wu, Y. C., & Yang, W. C. (2016). Herbal remedies for coccidiosis control: A review of plants, compounds and anticoccidial actions. *Evidence-Based Complementary and Alternative Medicine: ECAM*, 2016. <https://doi.org/10.1155/2016/2657981>
- N'dri, M. K. (2009). Etude comparée de la résistance à la coccidiose aviaire chez différentes races de poule. *Thèse en Pharmacie et d'Odonto-Stomatologie de l'Université Cheikh Anta Diop*. Dakar, Sénégal.
- Nghonjuyi, N.W., Tiambo, C. K., Kimbi, H. K., Manka'a, C. N., Juliano, R. S., & Lisita, F. (2015). Efficacy of ethanolic extract of *Carica papaya* leaves as a substitute of Sulphanomide for the control of coccidiosis in kabir chickens in Cameroon. *Journal of Animal Health and Production*, 3 (1): 21-27. <https://doi.org/10.14737/journal.jahp/2015/3.1.21.27>
- Quiroz-Castañeda, R. E., & Dantán-González, E. (2015). Control of avian coccidiosis: Future and present natural alternatives. *BioMed Research International*, 2015. <https://doi.org/10.1155/2015/430610>
- Reddy, N. R., Sathe, S. K., & Salunkhe, D. K. (1982). Phytates in legumes and cereals. *Advances in Food Research*, 28, 1-92. [https://doi.org/10.1016/S0065-2628\(08\)60110-X](https://doi.org/10.1016/S0065-2628(08)60110-X)
- Singh, P., Tanwar, N., Saha, T., Gupta, A., & Verma, S. (2018). Phytochemical screening and analysis of *Carica papaya*, agave Americana and piper nigrum. *Int J Curr Microbiol Appl Sci*, 7(2), 1786-1794. <https://doi.org/10.20546/ijcmas.2018.702.216>
- Soedji, K., Teteh, A., Gbeassor, M., Tona, K., Nideou, D., & Decuypere, E. (2017). Effect of *Carica papaya* seeds on gastro-intestinal parasites of pullet and production parameters. *International Journal of Probiotics and Prebiotics*, 12(2).
- Williams, R. B. (1994). Safety of the attenuated anticoccidial vaccine 'paracox' in broiler chickens isolated from extraneous coccidial infection. *Veterinary Research Communications*, 18, 189-198. <https://doi.org/10.1007/BF01839268>
- Yadav, M., Chatterji, S., Gupta, S. K., & Watal, G. (2014). Preliminary phytochemical screening of six medicinal plants used in traditional medicine. *Int J Pharm Pharm Sci*, 6(5), 539-42. ISSN-10: 0975-1491.
- Yim, D., Kang, S. S., Kim, D. W., Kim, S. H., Lillehoj, H. S., & Min, W. (2011). Protective effects of aloe vera-based diets in *Eimeria maxima*-infected broiler chickens. *Experimental Parasitology*, 127(1), 322-325. <https://doi.org/10.1016/j.exppara.2010.08.010>