

Association between Infective Endocarditis and Premalignant Colorectal Lesions

Carles Falces

Department of Cardiology, Cardiovascular Institute, Hospital Clinic-IDIBAPS, University of Barcelona, Barcelona, Spain

The association between Infective Endocarditis (IE) and Colorectal Cancer (CRC) has been known for more than 50 years. The microorganisms in which this association has been described are *Streptococcus gallolyticus* (formerly *Streptococcus bovis*) (Corredoira-Sánchez *et al.*, 2012; Boleij *et al.*, 2011), in different subspecies and, more recently, *Enterococcus faecalis* (Pericàs *et al.*, 2016; 2017; Escolà-Vergé *et al.*, 2020; Pericàs *et al.*, 2021).

Colonoscopy is the gold standard for the screening and diagnosis of colorectal cancer. American and European IE guidelines recommend the systematic performance of colonoscopy in patients with *S. gallolyticus* bacteremia and high suspicion of IE (Baddour *et al.*, 2015; Habib *et al.*, 2015). Previous studies have shown that colorectal neoplasms in patients with IE due to *S. gallolyticus* are much more frequent than in the general population, so colonoscopy is mandatory (Corredoira-Sánchez *et al.*, 2012). Boleij *et al.* (2011) systematic review and meta-analysis reported an association between infection from *S. gallolyticus* and CRC in 65% of cases. Although the association between *S. gallolyticus* IE and CRC is robust, the association with Colorectal Adenomas (CRA) is less conclusive, because CRA are also very frequent in the general population. Recent studies have shown that, like *S. gallolyticus* IE, *E. faecalis* IE is associated with Colorectal Neoplasms (CRN) and thus colonoscopy should also be mandatory in these patients (Pericàs *et al.*, 2016; 2017; Escolà-Vergé *et al.*, 2020; Pericàs *et al.*, 2021).

The association between IE (*S. gallolyticus* or *E. faecalis*) and CRA is highly relevant in clinical practice. In these patients, the diagnosis of severe infectious disease may allow the early diagnosis and treatment of a premalignant colorectal neoplasm.

In the present issue, Vilardell *et al.* analyze the relationship between streptococcal infective endocarditis and pre-neoplastic colorectal lesions in a retrospective single-center study of a Mediterranean population. In this study, 71 patients with IE due to any microorganism underwent colonoscopy as part of the extension analysis during hospitalization: 49 patients (69%) had IE without

colorectal lesions, 14 patients (20%) had IE with dysplastic adenomas and eight (11%) had IE with CRC. *S. gallolyticus* was the microorganism most frequently associated with colorectal disease, especially with pre-neoplastic lesions (50%). Not only high-degree but also low-degree dysplastic adenomas presented this association. The authors conclude that *S. gallolyticus* IE is associated with pre-cancerous colorectal lesions, including low-degree dysplastic adenomas. The results support the current recommendation to perform colonoscopy for the screening of colorectal cancer and pre-neoplastic lesions in patients with *S. gallolyticus* IE.

The relationship between *S. gallolyticus* and colorectal cancer is not well clarified. Colorectal lesions may simply be the gateway for the microorganism, but it may be that it is both a passenger and a cancer-promoting bacterium. For *S. gallolyticus* to facilitate the development of cancer it needs premalignant conditions, like CRA. Therefore, *S. gallolyticus* is not the main cause of CRC, but an auxiliary factor that accelerates its development (Pasquereau-Kotula *et al.*, 2018).

S. gallolyticus is not the only microorganism associated with CRN. Several studies have found that *E. faecalis* might bear carcinogenetic properties contributing to CRN development. Pericàs *et al.* (2016; 2017) reported a 17-fold higher prevalence of CRN in patients with *E. faecalis* IE with an unknown source of the infection undergoing colonoscopy than in the general population. Escolà-Vergé *et al.* (2020) also found similar rates of CRN in patients with a known source. More recently, a study from the GAMES cohort added to previous evidence suggesting a much higher rate of CRN among patients with *E. faecalis* IE than in the general population of the same age and sex (Pericàs *et al.*, 2021).

Conclusion

The association between *S. gallolyticus* IE and CRN (cancer and pre-malignant lesions) is well established. The indication for colonoscopy in these patients should be mandatory. In light of current data, colonoscopy

should also be recommended in patients with *E. faecalis* IE until further studies are available.

References

- Baddour, L. M., Wilson, W. R., Bayer, A. S., Fowler Jr, V. G., Tleyjeh, I. M., Rybak, M. J., ... & American Heart Association Committee on Rheumatic Fever, Endocarditis and Kawasaki Disease of the Council on Cardiovascular Disease in the Young, Council on Clinical Cardiology, Council on Cardiovascular Surgery and Anesthesia and Stroke Council. (2015). Infective endocarditis in adults: diagnosis, antimicrobial therapy and management of complications: a scientific statement for healthcare professionals from the American Heart Association. *Circulation*, 132(15), 1435-1486. <https://doi.org/10.1161/CIR.0000000000000296>
- Boleij, A., van Gelder, M. M., Swinkels, D. W., & Tjalsma, H. (2011). Clinical Importance of *Streptococcus gallolyticus* infection among colorectal cancer patients: systematic review and meta-analysis. *Clinical Infectious Diseases*, 53(9), 870-878. <https://doi.org/10.1093/cid/cir609>
- Corredoira-Sánchez, J., García-Garrote, F., Rabuñal, R., López-Roses, L., García-País, M. J., Castro, E., ... & Varela, J. (2012). Association Between Bacteremia Due to *Streptococcus gallolyticus* subsp. *gallolyticus* (*Streptococcus bovis* I) and Colorectal Neoplasia: A Case-Control Study. *Clinical Infectious Diseases*, 55(4), 491-496. <https://doi.org/10.1093/cid/cis434>
- Escolà-Vergé, L., Peghin, M., Givone, F., Pérez-Rodríguez, M. T., Suárez-Varela, M., Meije, Y., ... & Fernández-Hidalgo, N. (2020). Prevalence of colorectal disease in enterococcus faecalis infective endocarditis: results of an observational multicenter study. *Revista Española de Cardiología (English Edition)*, 73(9), 711-717. <https://doi.org/10.1016/j.recesp.2019.07.016>
- Habib, G., Lancellotti, P., Antunes, M. J., Bongiorni, M. G., Casalta, J. P., Del Zotti, F., ... & Zamorano, J. L. (2015). 2015 ESC guidelines for the management of infective endocarditis: the task force for the management of infective endocarditis of the European Society of Cardiology (ESC) endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the European Association of Nuclear Medicine (EANM). *European Heart Journal*, 36(44), 3075-3128. <https://doi.org/10.1093/eurheartj/ehv319>
- Pasquereau-Kotula, E., Martins, M., Aymeric, L., & Dramsi, S. (2018). Significance of *Streptococcus gallolyticus* subsp. *gallolyticus* association with colorectal cancer. *Frontiers in Microbiology*, 9, 614. <https://doi.org/10.3389/fmicb.2018.00614>
- Pericàs, J. M., Ambrosioni, J., Muñoz, P., de Alarcón, A., Kestler, M., Mari-Hualde, A., ... & Belaustegui, M. C. (2021, January). Prevalence of Colorectal Neoplasms Among Patients With Enterococcus faecalis Endocarditis in the GAMES Cohort (2008-2017). *Mayo Clinic Proceedings*, 96, 132-146. <https://doi.org/10.1016/j.mayocp.2020.06.056>
- Pericàs, J. M., Corredoira, J., & Miró Meda, J. M. (2016). Colorectal adenomas. *New England Journal of Medicine*, 375, 387-390. <https://doi.org/10.1056/NEJMc1604867>
- Pericàs, J. M., Corredoira, J., Moreno, A., García-País, M. J., Falces, C., Rabuñal, R., ... & Miró, J. M. (2017). Relationship between Enterococcus faecalis infective endocarditis and colorectal neoplasm: preliminary results from a cohort of 154 patients. *Revista Española de Cardiología (English Edition)*, 70(6), 451-458. <https://doi.org/10.1016/j.recesp.2016.09.055>