

Estimations of Neutrino and Graviton Masses by a Phenomenological Mass Relation for Stable Particles

Dimitar T. Valev

Space Research and Technology Institute, Bulgarian Academy of Sciences, 6000 Stara Zagora, Bulgaria

Article history

Received: 23-08-2014

Revised 30-01-2015

Accepted 07-05-2015

Abstract: The ratio between the proton and electron masses was shown to be close to the ratio between the shortest lifetimes of particles, decaying by the electromagnetic and strong interactions. The inherent property of each fundamental interaction is defined, namely the Minimal Lifetime of the Interaction (MLTI). The rest mass of the Lightest Free Massive Stable Particle (LFMSP), acted upon by a particular interaction, is shown to be inversely proportional to MLTI. The found mass relation unifies the masses of four stable particles of completely different kinds (proton, electron, electron neutrino and graviton) and covers an extremely wide range of values, exceeding 40 orders of magnitude. On the basis of this mass relation, the electron neutrino and graviton masses have been approximately estimated to $6.5 \times 10^{-4} \text{ eV}$ and $\hbar H/c^2 \approx 1.5 \times 10^{-33} \text{ eV}$, respectively. Besides, the last value has been obtained independently by dimensional analysis by means of three fundamental constants, namely the speed of light in vacuum (c), reduced Planck constant (\hbar) and Hubble constant (H). The presence of an exceptionally small, yet nonzero mass of the graviton implies Yukawa potential of the gravitational field and a finite range of the gravity close to the Hubble distance cH^{-1} . Therefore, the graviton mass determines a finite size of gravitationally connected (observable) universe for an arbitrary observer. It was shown that the rest energy of LFMSP, acted upon by a particular interaction, is close to Breit-Wigner's energy width of the shortest living state, decaying by the respective interaction.

Keywords: Neutrino Mass Limit, Graviton Mass, Mass Relation, Yukawa Potential, Dimensional Analysis

Introduction

Although the neutrino and the graviton belong to different particle kinds (neutral lepton and quantum of the gravitation, respectively), they have some similar properties. Both particles are not acted upon by the strong and the electromagnetic interactions, which makes their detection and investigation exceptionally difficult. Besides, both have masses that are many orders of magnitude lighter than the masses of the rest particles and they are generally accepted to be massless.

Decades after the experimental detection of the neutrino (Reines and Cowan, 1953), it was generally accepted that the neutrino rest mass m_{ov} is rigorously zero. The first experiment, hinting that the neutrino probably possesses a mass, is dated back to the 60-ies (Davis *et al.*, 1968). The total flux of neutrinos from the Sun is about 3 times lower than the one, predicted by theoretical solar models. This discrepancy can be

explained if some of the electron neutrinos transform into another neutrino flavor. Later, the experimental observations showed that the ratio between the atmospheric ν_μ and ν_e fluxes was less than the theoretical predictions (Hirata *et al.*, 1988; Casper *et al.*, 1991). Again the discrepancy could be explained by the neutrino oscillations. The crucial experiments with the 50 Kton neutrino detector Super-Kamiokande found strong evidence for oscillations (and hence-mass) in the atmospheric neutrinos (Fukuda *et al.*, 1998).

The direct neutrino measurements allow to limit the neutrino mass. The upper limit for the mass of the lightest neutrino flavor ν_e was obtained from experiments for measurement of the high-energy part of the tritium β -spectrum and recent experiments yield 2 eV upper limit (Weinheimer *et al.*, 1999; Lobashev *et al.*, 1999). As a result of the recent experiments, the upper mass limits of ν_μ and ν_τ were found to be 170 KeV (Assamagan *et al.*, 1996) and 18.2 MeV (Barate *et al.*, 1998), respectively.

The Solar and atmospheric neutrino experiments allow to find the square mass differences $\Delta m_{12}^2 = m_2^2 - m_1^2$ and $\Delta m_{23}^2 = m_3^2 - m_2^2$, but not the absolute values of the neutrino masses. The astrophysical constraint of the neutrino mass is $\Sigma m_\nu < 2.2 \text{ eV}$ (Bahcall, 1996). The recent extensions of the Standard model lead to non-zero neutrino masses, which are within the large range of $10^{-6} \div 10 \text{ eV}$.

Similarly to the case with the neutrino before 1998, the prevailing current opinion is that the quantum of the gravitation (graviton) is massless. This opinion is connected with Einstein's theory of General Relativity, where the gravitation is described by a massless field of spin 2 in a generally covariance manner. Although the massive gravity meets deep hardships, the interest in massive graviton considerably increases nowadays, especially relating to possibilities of revealing of dark mass and dark energy phenomena (Damour *et al.*, 2002; Feng *et al.*, 2003; Dubovsky *et al.* 2005; Lue, 2006; Babichev *et al.*, 2009; Mukohyama, 2009). The nonzero graviton mass leads to a finite gravitation range $r_g \sim \lambda_g = \hbar / (m_g c)$, where λ_g is Compton wavelength of the graviton. The lowest astrophysical limit of the graviton mass is obtained by rich galactic clusters $m_g < 2 \times 10^{-29} \text{ h}^{-1} \text{ eV}$ (Goldhaber and Nietto, 1974), where $h \approx 0.70$ is a dimensionless Hubble constant. In this case no difference was observed between Yukawa potential for the massive graviton and Newtonian potential for the massless graviton.

It has been obtained a value of the graviton mass $m_g \sim 4.3 \times 10^{-34} \text{ eV}$ for an infinite stationary universe (Woodward *et al.*, 1975), although the expansion of the Universe is a fact, long ago established. The mass $m_{i\min}$ of the Lightest Free Massive Stable Particle (LFMSP), acted upon by a particular interaction, is shown to be proportional to the coupling constant of the respective interaction at extremely low energy (Valev, 2008). The graviton and electron neutrino masses have been estimated by this approach to $m_g \sim 2.3 \times 10^{-34} \text{ eV}$ and $m_{\nu e} \sim 2.1 \times 10^{-4} \text{ eV}$, respectively.

Minimal Lifetime of a Particular Interaction

Among the multitude of particles, several free particles are notable, which are stable or at least their lifetimes are longer than the age of the Universe—the proton (p), electron (e), neutrino (ν) (three flavors), graviton (g) and photon (γ). Only *free massive stable* particles are examined in this study. Quarks and gluons are bound in hadrons by confinement and they cannot be immediately detected in the experiments and the photon is massless. Therefore, these particles are not a subject of this study.

A measure for the interaction strength is a dimensionless quantity—the coupling constant of the interaction (α_i), which is determined from the cross section of the respective processes. Generally, it is known that the

bigger the strength (coupling constant) of an interaction, the quicker (with shorter duration τ) are the processes, ruled by this interaction. Actually, the typical lifetime of resonances, decaying by the strong interaction (τ_s) is from 10^{-24} to 10^{-23} s , the time of the radiative decay (τ_e) of particles and excited stages of nuclei is from 10^{-21} to 10^{-12} s and, the lifetime (τ_w) of particles decaying by the weak interaction is from 10^{-12} to 10^3 s . Since “the age of the Universe” is $H^{-1} \sim 4.33 \times 10^{17} \text{ s} \approx 1.37 \times 10^{10} \text{ years}$, the lifetime of particles decaying by the gravitational interaction is $\tau_g \geq H^{-1}$.

The fastest process by the strong interaction is the f_0 (400-1200) resonance decay, having $\tau_{s\min} \approx 8 \times 10^{-25} \text{ s}$ (Groom, 2000). The fastest process by the electromagnetic interaction is the radiative decay of the super hot nuclei $\tau_{e\min} \sim \lambda_e / c \approx 1.3 \times 10^{-21} \text{ s}$, where λ_e is Compton wavelength of the electron (Nakamura, 2010). The fastest decay by the weak interaction is flavor transformation of the bottom and charmed quarks with $\tau_{w\min} \sim 10^{-12} \text{ s}$. The minimal lifetime $\tau_{g\min}$ of particles, decaying by the gravitational interaction is unknown, therefore we suggest that $\tau_{g\min} \sim H^{-1} \approx 4.3 \times 10^{17} \text{ s}$. Thus, the cosmological expansion of the Universe is considered a manifestation of the gravitational decay. Therefore, the minimal lifetime ($\tau_{i\min}$) of particles, decaying by a particular interaction, appears a *unique inherent property* of each interaction, below named Minimal Lifetime of The Interaction (MLTI).

Mass Relation for Free Massive Stable Particles

The ratio between the proton and electron masses is $m_p/m_e \approx 1836$. On the other side, the ratio between the minimal lifetimes of electromagnetic and strong interactions is $\tau_{e\min}/\tau_{s\min} \sim 1625$. The two ratios differentiate by less than 13%. Therefore, the ratio between the proton and electron masses is close to the ratio between the minimal lifetimes of the electromagnetic and strong interactions:

$$\frac{m_p}{m_e} \sim \frac{\tau_{e\min}}{\tau_{s\min}} \quad (1)$$

The proton and electron are the Lightest Free Massive Stable Particles (LFMSP), acted upon by the strong and electromagnetic interactions, respectively. This relation is remarkable since it connects the masses of LFMSP, acted upon by the strong and electromagnetic interactions and the respective MLTI. The relation (1) suggests that the mass of LFMSP, acted upon by the strong (or the electromagnetic) interaction is inversely proportional to the respective MLTI, i.e., $m_p \sim k/\tau_{s\min}$ and $m_e \sim k/\tau_{e\min}$, where k is a constant. Therefore, it is interesting to examine

whether this rule will be valid both for the weak interaction, whose MLTI is several orders of magnitude longer than the minimal lifetime of the electromagnetic interaction and for the gravity, whose minimal lifetime is dozens orders of magnitude longer than minimal lifetime of the weak interaction. LFMSP acted upon by the weak interaction is the electron neutrino and LFMSP acted upon by the gravity most probably appears the hypothetical graviton. Although the rest masses of the two particles are still unknown, the direct neutrino mass experiments and the theoretical models suggest that the ν_e mass is between 10^{-6} eV and 2 eV, i.e., ν_e is several orders of magnitude lighter than the electron. Again, the astrophysical constraints allow to find the upper limits of the graviton mass and according to these constraints, if the graviton really exists, its mass would be less than 3×10^{-29} eV, i.e., dozens orders of magnitude lighter than ν_e . Table 1 presents MLTI, as well as the masses of LFMSP, acted upon by the respective interaction. The experimental upper limits of the electron neutrino and graviton masses are also presented. Table 1 shows that the mass of LFMSP acted upon by a particular interaction decreases with the increase of MLTI.

The data from Table 1 have been presented in a double-logarithmic scale in Fig. 1, which shows that the trend is clearly expressed.

The points in Fig. 1, corresponding to the electron and proton masses and to the upper limit masses of the electron neutrino and graviton, are approximated by the least squares with a power law:

$$\log m_{\min} = -0.90 \log \tau_{\min} - 12.20 \quad (2)$$

Although this approximation is only on 4 points, the found correlation is close and the correlation coefficient reaches $r = 0.998$, which supports the power law. The modulus of the slope (S) is little smaller than one and that is why it can be said that the regression is close to a linear one. In addition, it should be reminded that instead of the electron neutrino and graviton masses, their upper limit values are used, which produce a certain underestimation of the S value. This approximation shows that the mass of LFMSP, acted upon by a particular interaction, increases with the decrease of the respective MLTI by a power law with $S \sim -1$, i.e., close to the inverse linear one. The inverse proportionality of the proton mass to the strong coupling constant and of the electron mass to the fine structure constant also support inverse linear dependence (*without intercept*). Thus, the experimental data suggest inverse linear dependence ($S = -1$) between the mass of LFMSP acted upon by a particular interaction and MLTI:

$$\log m_{\min} = -\log \tau_{\min} - k_0 \quad (3)$$

where, k_0 is a constant.

Table 1. MLTI and the masses of LFMSP acted upon by the respective interaction

Interaction	MLTI (s)	LFMSP	Mass (eV)
Strong	8×10^{-25}	Proton (p)	9.38×10^8
Electromagnetic	1.3×10^{-21}	Electron (e)	5.11×10^5
Weak	10^{-12}	Neutrino (ν_e)	$0 < m < 2$
Gravitational	4.3×10^{17}	Graviton (g)	$< 3 \times 10^{-29}$




Fig. 1. Dependence between the mass of LFMSP acted upon by a particular interaction and MLTI. The dashed line represents the approximation (2) of e , p and the upper limit masses of g and ν_e . The solid line represents the strict inverse linear approximation ($S = -1$)

The expression (3) transforms into:

$$m_{\min} \tau_{\min} = 10^{-k_0} = k \quad (4)$$

In this way the experimental data and constraints suggest that the mass of LFMSP, acted upon by a particular interaction, is inversely proportional to the respective MLTI:

$$m_{i \min} = \frac{k}{\tau_{i \min}} \quad (5)$$

where, $k = m_e \tau_{e \min} \approx 6.54 \times 10^{-16}$ eV s $= 1.16 \times 10^{-51}$ kg s is a constant, $i = 1, 2, 3, 4$ -index for each interaction and LFMSP acted upon by the respective interaction.

In consideration of $\tau_{e \min} \sim \lambda_e / c$ and $\lambda_e = \hbar / (m_e c)$, the mass relation (5) would be transformed in the equivalent mass formula:

$$m_{i \min} = \frac{m_e \tau_{e \min}}{\tau_{i \min}} \sim \frac{\hbar}{c^2 \tau_{i \min}} \quad (6)$$

Neutrino and Graviton Mass Estimations

The found mass relation (5) and equivalent mass formula (6), could be examined by the strong interaction because the proton mass is measured with

high precision. The application of the mass relation on the strong interaction predicts the lightest stable hadron mass $m_p \approx 819 \text{ MeV}$. Thus, the proton mass value obtained by the mass relation (5) is only 12.7% lower than the experimental value of m_p . This result confirms the reliability of the found mass relation and shows that this relation possesses heuristic power. The application of the mass relation (5) on the weak interaction allows to evaluate the mass of the electron neutrino $m_{\nu_e} \approx 6.5 \times 10^{-4} \text{ eV}$. This value is in order of magnitude of the estimation of the electron neutrino mass, found from Valev (2008).

The above obtained value $m_{\nu_e} \approx 6.5 \times 10^{-4} \text{ eV}$ and the results from the solar and atmospheric neutrino experiments allow to estimate the masses of the heavier neutrino flavor ν_μ and ν_τ . The results from the Super Kamiokande experiment lead to square mass difference $\Delta m_{23}^2 \sim 2.7 \times 10^{-3} \text{ eV}^2$ (Gonzalez-Garcia and Pena-Garay, 2003). Recent results on solar neutrinos provide hints that the Large Mixing Angle (LMA) of Mikheyev-Smirnov-Wolfenstein (MSW) solution is more probable than the Small Mixing Angle (SMA) (Krastev and Smirnov, 2002). The LMA leads to $\Delta m_{12}^2 \sim 7 \times 10^{-5} \text{ eV}^2$ (Maltoni *et al.*, 2003) and the SMA leads to $\Delta m_{12}^2 \sim 6 \times 10^{-6} \text{ eV}^2$ (Albright, 2001). In this way both solutions yield $m_{\nu_\tau} \sim 0.05 \text{ eV}$. The most appropriate LMA yields $m_{\nu_\mu} \approx 8.4 \times 10^{-3} \text{ eV}$ and the SMA leads to $m_{\nu_\mu} \approx 2.5 \times 10^{-3} \text{ eV}$. Thus, the obtained values of the neutrino masses support the normal hierarchy case. These values are close to the predictions of the simple SO(10) model for the neutrino masses (Dermisek, 2004).

In consideration of $\tau_g \sim H^{-1}$, the mass formula (6) allows to estimate the graviton mass:

$$m_g \sim \frac{\hbar}{c^2 \tau_g} \sim \frac{\hbar H}{c^2} \approx 1.5 \times 10^{-33} \text{ eV} \quad (7)$$

From Equation 7 we find amazing fact that the reduced Compton wavelength λ_g of graviton mass is equal to the Hubble distance cH^{-1} :

$$\lambda_g = \frac{\hbar}{m_g c} = cH^{-1} \sim 1.3 \times 10^{26} \text{ m} \quad (8)$$

The nonzero mass of the graviton, implies Yukawa type potential of the gravitational field $V(r) = -\frac{Gm}{r} \exp(-r/\lambda_g)$ and a finite range of gravitational interaction close to the Hubble distance cH^{-1} . Therefore, the mass of the graviton determines a finite size of gravitationally connected (observable) universe for an arbitrary observer. Thus, the gravitationally connected universe coincides with

Hubble sphere that is a 3-dimensional homogeneous isotropic sphere having radius $cH^{-1} \approx 1.38 \times 10^{10}$ light years and centered on the observer (Harrison, 2003). Beyond the Hubble sphere the galaxies recede from that observer at a rate greater than the speed of light due to the expansion of the Universe and they are not gravitationally connected with the observer.

The predicted masses of four LFMSF are presented in Table 2, where it can be seen that the fitting of the predicted values and the experimental data is satisfactory.

The exceptionally small graviton rest mass seriously impedes its experimental determination. Yet, it can be expected that appropriate astrophysical or laboratory experiments would be conducted for this aim. Probably, the investigations of the large-scale structure of the universe and the microwave background radiation would contribute to the astrophysical estimation of the graviton mass. The massive graviton might turn of considerable importance for the description of the processes in the nuclei of the active galaxies and quasars, the gravitational collapse as well as for the improvement of the cosmological models.

Besides, the formula (7) for graviton mass could be obtained *independently* by dimensional analysis. Actually, by means of three fundamental constants, namely the speed of light in vacuum (c), reduced Planck constant (\hbar) and Hubble constant (H), a mass dimension quantity m_x could be constructed:

$$m_x = kc^\alpha \hbar^\beta H^\gamma \quad (9)$$

where, k is dimensionless parameter of the order of magnitude of one and α , β and γ are unknown exponents, which will be determined by dimensional analysis below.

Dimensional analysis has been successfully used from Valev (2014) for estimation of mass of the observable universe by means of following fundamental constants-the speed of light (c), universal gravitational constant (G) and Hubble constant (H).

The dimensions of the left and right sides of the Equation 9 must be equal:

$$[m_x] = [c]^\alpha [\hbar]^\beta [H]^\gamma \quad (10)$$

Taking into account the dimensions of quantities in formula (10) we obtain:

$$L^0 T^0 M^1 = (LT^{-1})^\alpha (ML^2 T^{-1})^\beta (T^{-1})^\gamma = L^{\alpha+2\beta} T^{-\alpha-\beta-\gamma} M^\beta \quad (11)$$

where, L , T and M are dimensions of length, distance and mass, respectively.

Table 2. Experimental and predicted values of LFMSP masses (eV)

Particle	Experimental mass	Predicted mass (This paper)	Predicted mass (Valev, 2008)
p	9.38×10^8	8.19×10^8	9.80×10^8
e	5.11×10^5	5.11×10^5	5.11×10^5
ν_e	$0 < m < 2$	6.5×10^{-4}	2.1×10^{-4}
g	$< 3 \times 10^{-29}$	1.5×10^{-33}	2.3×10^{-34}

From (11) we obtain the system of linear equation:

$$\begin{aligned} \alpha + 2\beta &= 0 \\ -\alpha - \beta - \gamma &= 0 \\ \beta &= 1 \end{aligned} \quad (12)$$

Solving the system we find the exponents $\alpha = -2, \beta = 1, \gamma = 1$. Replacing the obtained values of the exponents in Equation 9 we find the formula (13) for the graviton mass:

$$m_x \sim \frac{\hbar H}{c^2} \quad (13)$$

Although this formula has been found by totally different approach, it coincides with formula (7), which reinforces the found phenomenological mass relation (6).

According Big Bang cosmology (Hubble, 1929), Hubble constant decrease with age of the universe, therefore the found graviton mass (7) slowly decrease with time. On the other hand, according Tired Light model (Zwicky, 1929) and Steady State theory (Bondi and Gold, 1948), the Hubble constant H is truly a constant not only in all directions, but at all time. Therefore, the graviton mass is truly constant in the framework of Tired Light model and Steady State theory.

Discussion and Conclusion

From mass formula (6) we obtain:

$$m_{i\min} c^2 \sim \frac{\hbar}{\tau_{i\min}} \approx \Gamma_{i\min} \quad (14)$$

where, $\Gamma_{i\min}$ is Breit-Wigner's energy width of the shortest living state, decaying by the respective interaction.

Therefore, the rest energy of LFMSP acted upon by a particular interaction is close to Breit-Wigner's energy width of the shortest living state, decaying by the respective interaction. It should be reminded that here $\tau_{i\min}$ isn't the lifetime of the particle (it is stable) but $\tau_{i\min}$ is the respective MLTI.

The mass formula (6) would be written in the form:

$$m_{i\min} c^2 \tau_{i\min} \sim \hbar \quad (15)$$

The mass (m_i) of each free massive stable particle, acted upon by a particular interaction $m_i \geq m_{i\min}$ and the lifetime (τ_i)

of particles decaying by the respective interaction $\tau_i \geq \tau_{i\min}$. As a result, the inequality (16) is obtained:

$$m_i c^2 \tau_i \geq \hbar \quad (16)$$

The comparison of (16) and the Uncertainty Principle $\Delta E \Delta t = \Delta(m c^2) \Delta t \geq \hbar$ shows that the inequality (16), which results from the mass formula (6), is related to the Uncertainty Principle. Thus, Equation 15 appears a boundary case of the Uncertainty Principle at minimal allowed values of the rest energy and lifetime of the real particles. In this case, however, a more general interpretation of the Uncertainty Principle will be necessary since Equation 15 relates the mass ($m_{i\min}$) of LFMSP acted upon by a particular interaction with minimal lifetime ($\tau_{i\min}$) of particles (states), decaying by the respective interaction. The future complete Unified theory of the four interactions would give theoretical explanation of this dependence. Most probably $\tau_{i\min}$ determines inability of the respective interaction to create free massive stable particles, possessing rest mass $m_i \geq m_{i\min} = \hbar / (c^2 \tau_{i\min})$. In other words, the stronger an interaction the smaller is MLTI and the heavier is LFMSP, which it is capable to create.

The mass relation (17) has been obtained in (Valev, 2008) by a similar phenomenological approach:

$$m_{i\min} = \frac{m_e}{\alpha} \alpha_i(0) \quad (17)$$

where, $m_{i\min}$ is the mass of LFMSP, acted upon by a particular interaction, $\alpha_i(0)$ is the coupling constant of a particular interaction at extremely low energy $E \sim m_e c^2$ and α is the fine structure constant.

From (6) and (17) we obtain:

$$\alpha_i(0) = \frac{\hbar \alpha}{m_e c^2 \tau_{i\min}} = \frac{\alpha \tilde{\lambda}_e}{c \tau_{i\min}} \sim \alpha \frac{\tau_{e\min}}{\tau_{i\min}} \approx \frac{\tau_n}{\tau_{i\min}} \quad (18)$$

where, τ_n is the nuclear time.

Equation 18 supports the natural suggestion that the coupling constant of a particular interaction at extremely low energy $\alpha_i(0)$ is inversely proportional to MLTI and it determines from the ratio $\tau_n / \tau_{i\min}$.

It is worth noting that each LFMSP, acted upon by a particular interaction also appears the lightest free massive particle, possessing the respective *universal*

conserving quantity-baryon number, electric charge, lepton number and mass. Actually, p , e , ν_e and g are LFMSP, acted upon by the strong, electromagnetic, weak and gravitational interaction, respectively and they also appear lightest free massive particles, possessing baryon number, electric charge, lepton number and mass. It should also be mentioned that all four lightest free massive particles, possessing universal conserving quantity are stable or, at least their lifetimes are longer than the age of the Universe.

The massive graviton rises severe challenges before the modern unified theories. Among them are van Dam-Veltman-Zakharov (νDVZ) discontinuity (van Dam and Veltman, 1970; Zakharov, 1970) and the violation of the gauge invariance and the general covariance. There are, however, already encouraging attempts to solve νDVZ discontinuity in Anti de Sitter (AdS) background (Dvali *et al.*, 2000; Kogan *et al.*, 2001).

It was found that the ratio between the proton and electron masses is close to the ratio between the shortest lifetime of particles, decaying by the electromagnetic and strong interaction. The inherent property of each fundamental interaction is defined, namely the Minimal Lifetime of The Interaction (MLTI). Inverse proportionality has been found between MLTI as well as the rest mass of the LFMSP acted upon by the respective interaction $m_{i\min} = k / \tau_{i\min} \sim \hbar / (c^2 \tau_{i\min})$.

The rest mass of the electron neutrino has been obtained by this approach to $m_{\nu_e} \approx 6.5 \times 10^{-4} eV$. The masses of the heavier neutrino flavors have been estimated by the results of the solar and atmospheric neutrino experiments. The mass of ν_τ is estimated to $0.05 eV$ and the mass of ν_μ is estimated to $8.4 \times 10^{-3} eV$ for LMA and $2.5 \times 10^{-3} eV$ for SMA . The graviton rest mass has been estimated by this approach $m_g \sim \hbar H / c^2 \approx 1.5 \times 10^{-33} eV$. Besides, the last value has been obtained independently by dimensional analysis by means of the fundamental constants c , \hbar and H . The presence of a small nonzero mass of the graviton implies Yukawa potential of the gravitational field $V(r) = -\frac{Gm}{r} \exp(-r / \lambda_g)$ and a finite range of the gravity close to the Hubble distance cH^{-1} . Therefore, the Hubble distance $cH^{-1} \approx 1.38 \times 10^{10}$ light years appears the size of gravitationally connected (observable) universe for an arbitrary observer.

It has been found that the rest energy of LFMSP, acted upon by a particular interaction, is close to Breit-Wigner's energy width of the shortest living state, decaying by the respective interaction. It has been shown that the mass formula for free massive stable particles $m_{i\min} = \hbar / (c^2 \tau_{i\min})$ probably involves the Uncertainty Principle.

Acknowledgment

I would like to thank Prof. Ivan G. Koprnikov and Dr. Andrew W. Beckwith for encouraging discussions.

Ethics

This article is original and contains unpublished material. The corresponding author confirms that all of the other authors have read and approved the manuscript and no ethical issues involved.

References

- Albright, C., 2001. Overview of grand unified models and their predictions for neutrino oscillations. Nucl. Instr. Meth. Phys. Res. A, 472: 359-363. DOI: 10.1016/S0168-9002(01)01271-2
- Assamagan, K., C. Brönnimann, M. Daum, H. Forrer and R. Frosch *et al.*, 1996. Upper limit of the muon-neutrino mass and charged-pion mass from momentum analysis of a surface muon beam. Phys. Rev. D, 53: 6065-6077. DOI: 10.1103/PhysRevD.53.6065
- Babichev, E., C. Deffayet and R. Ziour, 2009. Recovering general relativity from massive gravity. Phys. Rev. Lett., 103: 201102-201102. DOI: 10.1103/PhysRevLett.103.201102
- Bahcall, J., 1996. Solar neutrinos-where we are, where we are going. Astrophys. J., 467: 475-484. DOI: 10.1086/177624
- Barate, R., D. Buskulic, D. Decamp, J.B. Hansen and J.D. Hansen *et al.*, 1998. An upper limit on the tau neutrino mass from three-prong and five-prong tau decays. Eur. Phys. J. C, 2: 395-406. DOI: 10.1007/s100520050149
- Bondi, H. and T. Gold, 1948. The steady-state theory of the expanding universe. Mon. Not. Roy. Astron. Soc., 108: 252-270. DOI: 10.1093/mnras/108.3.252.
- Casper, D., R. Becker-Szendy, C.B. Bratton, D.R. Cady and R. Claus *et al.*, 1991. Measurement of atmospheric neutrino composition with the IMB-3 detector. Phys. Rev. Lett., 66: 2561-2564. DOI: 10.1103/PhysRevLett.66.2561
- Damour, T., I.I. Kogan and A. Papazoglou, 2002. Nonlinear bigravity and cosmic acceleration. Phys. Rev. D, 66: 104025-104025. DOI: 10.1103/PhysRevD.66.104025
- Davis, R., D. Harmer and K. Hoffman, 1968. Search for neutrinos from the Sun. Phys. Rev. Lett., 20: 1205-1209. DOI: 10.1103/PhysRevLett.20.1205
- Dermisek, R., 2004. Neutrino masses and mixing, quark-lepton symmetry and strong right-handed neutrino hierarchy. Phys. Rev. D, 70: 073016-073016. DOI: 10.1103/PhysRevD.70.073016

- Dubovsky, S.L., P.G. Tinyakov and I.I. Tkachev, 2005. Massive graviton as a testable cold-dark-matter candidate. *Phys. Rev. Lett.*, 94: 181102-181102. DOI: 10.1103/PhysRevLett.94.181102
- Dvali, G., G. Gabadadze and M. Porrati, 2000. 4D gravity on a brane in 5D Minkowski space. *Phys. Lett. B*, 485: 208-214. DOI: 10.1016/S0370-2693(00)00669-9
- Feng, J.L., A. Rajaraman and F. Takayama, 2003. Graviton cosmology in universal extra dimensions. *Phys. Rev. D*, 68: 085018-085018. DOI: 10.1103/PhysRevD.68.085018
- Fukuda, Y., T. Hayakawa, E. Ichihara, K. Inoue and K. Ishihara *et al.*, 1998. Evidence for oscillation of atmospheric neutrinos. *Phys. Rev. Lett.*, 81: 1562-1567. DOI: 10.1103/PhysRevLett.81.1562
- Goldhaber, A. and M. Nietto, 1974. Mass of the graviton. *Phys. Rev. D*, 9: 1119-1121. DOI: 10.1103/PhysRevD.9.1119
- Gonzalez-Garcia, M. and C. Pena-Garay, 2003. Three-neutrino mixing after the first results from K2K and KamLAND. *Phys. Rev. D*, 68: 093003-093003. DOI: 10.1103/PhysRevD.68.093003
- Groom, D., 2000. Light unflavored mesons. *Eur. Phys. J. C*, 15: 405-405.
- Harrison, E., 2003. *Masks of the Universe*. 1st Edn., Cambridge University Press, ISBN-10: 0521773512, pp: 342.
- Hirata, K.S., T. Kajita, M. Koshiba, M. Nakahata and S. Ohara *et al.*, 1988. Experimental study of the atmospheric neutrino flux. *Phys. Lett. B*, 205: 416-420. DOI: 10.1016/0370-2693(88)91690-5
- Hubble, E., 1929. Relation between distance and radial velocity among extra-galactic nebulae. *Proc. Nat. Acad. Sci.*, 15: 168-173. DOI: 10.1073/pnas.15.3.168
- Kogan, I.I., S. Mouslopoulos and A. Papazoglou, 2001. The $m \rightarrow 0$ limit for massive graviton in dS_4 and AdS_4 -How to circumvent the van Dam-Veltman-Zakharov discontinuity. *Phys. Lett. B*, 503: 173-180. DOI: 10.1016/S0370-2693(01)00209-X
- Krastev, P. and A. Smirnov, 2002. Global analysis with SNO: Toward the solution of the solar neutrino problem. *Phys. Rev. D*, 65: 073022-073022. DOI: 10.1103/PhysRevD.65.073022
- Lobashev, V.M., V.N. Aseev, A.I. Belesev, A.I. Berlev and E.V. Geraskin *et al.*, 1999. Direct search for mass of neutrino and anomaly in the tritium beta spectrum. *Phys. Lett. B*, 460: 227-235. DOI: 10.1016/S0370-2693(99)00781-9
- Lue, A., 2006. The phenomenology of Dvali Gabadadze Porrati cosmologies. *Physics Reports*, 423: 1-48. DOI: 10.1016/j.physrep.2005.10.007
- Maltoni, M., T. Schwetz, M.A. Tortola and J.W.F. Valle, 2003. Status of three-neutrino oscillations after the SNO-salt data. *Phys. Rev. D*, 68: 113010-113010. DOI: 10.1103/PhysRevD.68.113010
- Mukohyama, S., 2009. Dark matter as integration constant in Horava-Lifshitz gravity. *Phys. Rev. D*, 80: 064005-064005. DOI: 10.1103/PhysRevD.80.064005
- Nakamura, K., 2010. Review of particle physics. *J. Phys. G*, 37: 075021-075021. DOI: 10.1088/0954-3899/37/7A/075021
- Reines, F. and C. Cowan, 1953. Detection of the free neutrino. *Phys. Rev.*, 92: 830-830. DOI: 10.1103/PhysRev.92.830
- Valev, D., 2008. Neutrino and graviton rest mass estimations by a phenomenological approach. *Aerospace Res. Bulg.*, 22: 68-82.
- Valev, D., 2014. Estimations of total mass and energy of the observable universe. *Phys. Int.*, 5: 15-20. DOI: 10.3844/pisp.2014.15.20
- van Dam, H. and M.J. Veltman, 1970. Massive and mass-less Yang-Mills and gravitational fields. *Nucl. Phys. B*, 22: 397-411. DOI: 10.1016/0550-3213(70)90416-5
- Weinheimer, C., B. Degenndag, A. Bleile, J. Bonn and L. Bornschein *et al.*, 1999. High precision measurement of the tritium beta spectrum near its endpoint and upper limit on the neutrino mass. *Phys. Lett. B*, 460: 219-226. DOI: 10.1016/S0370-2693(99)00780-7
- Woodward, J.F., R.J. Crowley and W. Yourgrau, 1975. Mach's principle and the rest mass of the graviton. *Phys. Rev. D* 11, 1371-1374. DOI: 10.1103/PhysRevD.11.1371
- Zakharov, V., 1970. Linearized gravitation theory and the graviton mass. *JETP Lett.*, 12: 312-314.
- Zwicky, F., 1929. On the red shift of spectral lines through interstellar space. *Proc. Nat. Acad. Sci.*, 15: 773-779. DOI: 10.1073/pnas.15.10.773